

THE SET OF INVARIANT STAR PRODUCTS ON NON-DEGENERATE TRIANGULAR LIE BIALGEBRAS OBTAINED IN THE ETINGOF-KAZHDAN QUANTIZATION THEORY

CARLOS MORENO AND JOANA TELES

ABSTRACT: Let $(\mathfrak{a}, [\cdot, \cdot]_{\mathfrak{a}}, \varepsilon_{\mathfrak{a}} = d_c r_1)$ be a non-degenerate triangular finite dimensional Lie bialgebra over \mathbb{R} . When a Lie associator Φ is fixed and for any formal solution $r_{\hbar} = r_1 + \hbar r_2 + \dots \in (\mathfrak{a} \wedge \mathfrak{a})[[\hbar]]$ of Yang Baxter Equation (YBE), the theory of Etingof-Kazhdan allows us to obtain an Invariant Star Product (ISP), $\tilde{J}_{r_{\hbar}} \in (\mathcal{U}\mathfrak{a} \otimes \mathcal{U}\mathfrak{a})[[\hbar]]$ on $(\mathfrak{a}, [\cdot, \cdot]_{\mathfrak{a}}, \varepsilon_{\mathfrak{a}} = d_c r_1)$. Every ISP F determines an element r_{\hbar} such that F and $\tilde{J}_{r_{\hbar}}$ are equivalent. We define in this way a bijective mapping between the set of classes of ISPS modulo equivalence and the co-homology space $\hbar H^2(\mathfrak{a})[[\hbar]]$.

KEYWORDS: Quantization of Lie bialgebras, Star Products, Quantized Universal Enveloping algebras.

AMS SUBJECT CLASSIFICATION (2000): 16W30; 17B62; 17B37; 46L65; 53D55.

1. Introduction

1) Let $(\mathfrak{a}, [\cdot, \cdot]_{\mathfrak{a}})$ be the Lie algebra of a Lie group \mathbf{G} . Let (\mathbf{G}, β_1) be a connected and simply connected Lie group endowed with an invariant symplectic structure $\beta_1 \in \mathfrak{a}^* \wedge \mathfrak{a}^*$ and let $r_1 \in \mathfrak{a} \wedge \mathfrak{a}$ be the corresponding solution to the Yang-Baxter-Equation (YBE) on $(\mathfrak{a}, [\cdot, \cdot]_{\mathfrak{a}})$. Let $(\mathfrak{a}, [\cdot, \cdot]_{\mathfrak{a}}, \varepsilon_{\mathfrak{a}} = d_c r_1)$ be the corresponding non-degenerate triangular Lie bialgebra. In [5] Drinfeld obtains all the Invariant Star Products (ISPS) on (\mathbf{G}, β_1) (equivalently on $(\mathfrak{a}, [\cdot, \cdot]_{\mathfrak{a}}, \varepsilon_{\mathfrak{a}} = d_c r_1)$) and a theorem showing that under equivalence [10] the classifying set of all those ISPS is the Chevalley space $\beta_1 + \hbar \cdot \sum H^2(\mathfrak{a}, \mathbb{R})[[\hbar]]$.

2) The aim of this paper is to obtain a classification theorem for all the ISPS on any non-degenerate triangular finite dimensional Lie bialgebra over \mathbb{R} when they are obtained following the Etingof-Kazhdan [9] theory of quantization of Lie bialgebras. This theory is very different from the one considered in [5] but the classifying set for the set of ISPS is again $\beta_1 + \hbar \cdot \sum H^2(\mathfrak{a}, \mathbb{R})[[\hbar]]$. In case we consider the Knizhnik-Zamolodchikov associator [7], over the field

Received January 16, 2008.

The second author has been partially supported by CMUC and FCT - project POCI/MAT/58452/2004.

\mathbb{C} , we have seen [22] that the ISP \tilde{J}_{r_1} obtained here and the ISP $F(r_1)$ obtained by Drinfeld in [5] coincide modulo \hbar^3 . In two particular cases of dimensions 2 and 4 for \mathfrak{a} , \tilde{J}_{r_1} and $F(r_1)$ coincide modulo \hbar^4 .

3) The adjoint representation of \mathbf{G} induces a representation on the Chevalley complex $H^*(\mathfrak{a}, \mathbb{R})$ which is trivial. This classical theorem contains the idea for the proof of our classification theorem 6.11. We may compare this with the proofs in [19, 20] for the similar theorem in the quantization context of [5].

We present here the proofs of most of the results obtained.

4) Similar results as in the Abstract can be obtained on a *non-degenerate triangular deformation Lie bialgebra*, [6, 7], $(\mathfrak{a}_t \equiv \mathfrak{a} \otimes_{\mathbb{K}} \mathbb{K}[[t]], [,]_{\mathfrak{a}_t}, \varepsilon_{\mathfrak{a}_t} = d_c(t)r_1(t))$ over $\mathbb{K}[[t]]$ of the non-degenerate triangular Lie bialgebra $(\mathfrak{a}, [,]_{\mathfrak{a}}, \varepsilon_{\mathfrak{a}} = d_c r_1)$ over a field of characteristic zero \mathbb{K} . In this case we need to observe that a) $\mathbb{K}[[t]]$ is a local ring and a Principal Ideal Domain; b) the symmetric algebra of the $\mathbb{K}[[t]]$ -module $\mathfrak{a}[[t]]$ is the algebra of divided powers $\Gamma(\mathfrak{a}_t)$, in the sense of [3], over the $\mathbb{K}[[t]]$ -module \mathfrak{a}_t ; c) this symmetric algebra is isomorphic to the algebra of symmetric tensors $TS(\mathfrak{a}[[t]])$ [4]; d) the Hochschild cohomology $H^*(\mathcal{U}\mathfrak{a}[[t]])$ of the coalgebra $\mathcal{U}\mathfrak{a}[[t]]$ over $\mathbb{K}[[t]]$ is $\wedge(\mathfrak{a}[[t]])$ [3].

2. Some notations

1) Definitions and notations are those of [5, 7, 15, 16]. A finite dimensional Lie bialgebra over \mathbb{R} is denoted by the symbol $(\mathfrak{a}, [,]_{\mathfrak{a}}, \varepsilon_{\mathfrak{a}})$ where $\varepsilon_{\mathfrak{a}}$ is a 1-cocycle in the Chevalley cohomology on $(\mathfrak{a}, [,]_{\mathfrak{a}})$ with respect to the $ad_{\mathfrak{a}}$ -representation. When it is quasitriangular we write $\varepsilon_{\mathfrak{a}} = d_c r_1$, where d_c is the coboundary in the above cohomology, $r_1 \in \mathfrak{a} \otimes \mathfrak{a}$ is a solution to CYBE, $[r_1, r_1] = 0$ on $(\mathfrak{a}, [,]_{\mathfrak{a}})$ and $r_1 + \sigma(r_1)$ is $ad_{\mathfrak{a}}$ -invariant where σ is the permutation (12). In case r_1 is skew-symmetric, it is a triangular Lie bialgebra and if moreover $\det(r_1) \neq 0$ we call it a non-degenerate triangular Lie bialgebra. If $(\mathfrak{a}, [,]_{\mathfrak{a}}, \varepsilon_{\mathfrak{a}})$ is a Lie bialgebra we denote its quasitriangular double Lie bialgebra as the set $(\mathfrak{a} \oplus \mathfrak{a}^*, [,]_{\mathfrak{a} \oplus \mathfrak{a}^*}, \varepsilon_{\mathfrak{a} \oplus \mathfrak{a}^*} = d_c r)$, where $r \in (\mathfrak{a} \oplus \mathfrak{a}^*)^{\otimes 2}$ is the invariant canonical element [2]. The element $\Omega = r + \sigma(r)$ is symmetric and $ad_{\mathfrak{a} \oplus \mathfrak{a}^*}$ -invariant.

2) The symbol $(\mathfrak{a}[[\hbar]], [,]_{\mathfrak{a}[[\hbar]]})$ will denote the Lie algebra over the ring $\mathbb{R}[[\hbar]]$ obtained from the Lie algebra $(\mathfrak{a}, [,]_{\mathfrak{a}})$ over \mathbb{R} by the extension of scalars $\mathbb{R} \rightarrow \mathbb{R}[[\hbar]]$. $(\mathfrak{a} \oplus \mathfrak{a}^*)[[\hbar]]$ will denote the Lie algebra (bialgebra) over $\mathbb{R}[[\hbar]]$ which is the extension of the Lie bialgebra $(\mathfrak{a} \oplus \mathfrak{a}^*, [,]_{\mathfrak{a} \oplus \mathfrak{a}^*}, \varepsilon_{\mathfrak{a} \oplus \mathfrak{a}^*} = d_c r)$ over \mathbb{R} . It is obvious that $\Omega = r + \sigma(r) \in (\mathfrak{a} \oplus \mathfrak{a}^*)^{\otimes 2}[[\hbar]]$ is $ad_{\mathfrak{a}[[\hbar]]}$ invariant

and that these Lie-algebras over $\mathbb{R}[[\hbar]]$ are deformations algebras [7] of their corresponding Lie-algebras over \mathbb{R} .

3) Let $r_t = r_1 + r_2 t + r_3 t^2 + \dots \in \mathfrak{a} \wedge \mathfrak{a}$ be an analytic function on a neighborhood of $0 \in \mathbb{R}$ defining a non-degenerate, $\text{rank}(r_1) = \dim \mathfrak{a}$, solution of the CYBE $[r_t, r_t] = 0$ on the Lie algebra $(\mathfrak{a}, [,]_{\mathfrak{a}})$ over \mathbb{R} . For each t , $(\mathfrak{a}, [,]_{\mathfrak{a}}, \varepsilon_{\mathfrak{a}} = d_c r_t)$ is then a non-degenerate triangular Lie bialgebra on $M\mathbb{R}$. The symbol $(\mathfrak{a} \oplus \mathfrak{a}_{r_t}^*, [,]_{\mathfrak{a} \oplus \mathfrak{a}_{r_t}^*}, \varepsilon_{\mathfrak{a} \oplus \mathfrak{a}_{r_t}^*} = d_c r)$ will denote the corresponding quasitriangular double Lie bialgebra on \mathbb{R} . Let $\mu_{r_t} : \wedge^2 \mathfrak{a} \longrightarrow \wedge^2 \mathfrak{a}^*$ be the linear isomorphism defined by the Poisson cocycle r_t . It induces an isomorphism between Poisson and Chevalley cohomology spaces [14]. Let (\mathbf{G}, β_t) be the corresponding connected and simply-connected Lie group endowed with the invariant symplectic structure $\beta_t = \mu_{r_t}(r_t)$.

4) Let $r_{\hbar} = r_1 + r_2 \hbar + r_3 \hbar^2 + \dots \in (\mathfrak{a} \wedge \mathfrak{a})[[\hbar]]$ be an invertible element solution of the CYBE $[r_{\hbar}, r_{\hbar}] = 0$ on $(\mathfrak{a}[[\hbar]], [,]_{\mathfrak{a}[[\hbar]]})$. The set $(\mathfrak{a}[[\hbar]], [,]_{\mathfrak{a}[[\hbar]]}, \varepsilon_{\mathfrak{a}[[\hbar]]} = d_{cr_{\hbar}})$ will denote the corresponding triangular non-degenerate Lie bialgebra over $\mathbb{R}[[\hbar]]$ and its quasitriangular double Lie bialgebra will be denoted by $(\mathfrak{a} \oplus \mathfrak{a}_{r_{\hbar}}^*[[\hbar]], [,]_{(\mathfrak{a} \oplus \mathfrak{a}_{r_{\hbar}}^*)}, \varepsilon_{\mathfrak{a} \oplus \mathfrak{a}_{r_{\hbar}}^*} = d_{cr})$. In this situation, let $\mu_{r_{\hbar}}$ be the isomorphism similar to μ_{r_t} in 3).

5) We fix [9] a Lie associator $\Phi = \exp P(\hbar t_{12}, \hbar t_{23})$ over \mathbb{R} .

3. Finite dimensional Etingof-Kazhdan quantization theory.

3.1. Quantization of the pair $(\mathfrak{a} \oplus \mathfrak{a}^*; r)$. From theorem A'' in [7], we can deduce the following:

Theorem 3.1. [7] Let $(\mathfrak{a}, [,]_{\mathfrak{a}}, \varepsilon_{\mathfrak{a}})$, $(\mathfrak{a} \oplus \mathfrak{a}^*, [,]_{\mathfrak{a} \oplus \mathfrak{a}^*}, \varepsilon_{\mathfrak{a} \oplus \mathfrak{a}^*} = d_c r)$ and $\Omega = r + \sigma(r)$ be as in section 2, 1). Let $\Phi = \exp P(\hbar \Omega_{12}, \hbar \Omega_{23}) \in \mathcal{U}(\mathfrak{a} \oplus \mathfrak{a}^*)^{\otimes 3}[[\hbar]]$. Let $(\mathcal{U}(\mathfrak{a} \oplus \mathfrak{a}^*)[[\hbar]], \cdot, \Delta_0, \epsilon_0, S_0)$ be the usual Hopf universal enveloping algebra. Write $\Phi = \sum_i X_i \otimes Y_i \otimes Z_i$ and $c = \sum_i X_i \cdot S_0(Y_i) \cdot Z_i$. Then the set

$$\left(\mathcal{U}(\mathfrak{a} \oplus \mathfrak{a}^*)[[\hbar]]; \cdot; 1; \Delta_0; \epsilon_0; \Phi; S_0; \alpha = c^{-1}; \beta = 1; R = e^{\frac{\hbar}{2}\Omega} \right)$$

is a quasitriangular quasi-Hopf QUE-algebra whose classical limit [7] is $(\mathfrak{a} \oplus \mathfrak{a}^*; \Omega)$.

The existence of the antipode follows from Theorem 1.6 in [7]. From Propositions 1.1 and 1.3 in [7] and from [11] it can be taken, [22], as the triple $(S_0; \alpha = c^{-1}; \beta = 1)$.

To quantize the pair $(\mathfrak{a} \oplus \mathfrak{a}^*; r)$ is to obtain a quasitriangular-Hopf QUE-algebra over $\mathbb{R}[[\hbar]]$ such that its classical limit is that pair. A main theorem in this direction is the following (see also [8]):

Theorem 3.2. [9] *Let $(\mathfrak{a}, [,]_{\mathfrak{a}}, \varepsilon_{\mathfrak{a}})$, $(\mathfrak{a} \oplus \mathfrak{a}^*, [,]_{\mathfrak{a} \oplus \mathfrak{a}^*}, \varepsilon_{\mathfrak{a} \oplus \mathfrak{a}^*} = d_c r)$ and $\Omega = r + \sigma(r)$ be as in section 2, 1). Let M_{\pm} be the $\mathfrak{a} \oplus \mathfrak{a}^*$ -modules with one generator 1_{\pm} and defined as follows: $M_+ = \mathcal{U}\mathfrak{a}^* \cdot 1_+$, $\mathcal{U}\mathfrak{a} \cdot 1_+ = 0$ and $M_- = \mathcal{U}\mathfrak{a} \cdot 1_-$, $\mathcal{U}\mathfrak{a}^* \cdot 1_- = 0$. Then*

1) *The equalities $i_{\pm}(1_{\pm}) = 1_{\pm} \otimes 1_{\pm}$ define unique $\mathfrak{a} \oplus \mathfrak{a}^*$ -module morphisms $i_{\pm} : M_{\pm} \longrightarrow M_{\pm} \otimes M_{\pm}$.*

2) *The equality $\phi(1) = 1_+ \otimes 1_-$ defines a unique $\mathfrak{a} \oplus \mathfrak{a}^*$ -module morphism $\phi : \mathcal{U}(\mathfrak{a} \oplus \mathfrak{a}^*) \longrightarrow M_+ \otimes M_-$. ϕ is an isomorphism.*

3) *There exists an element $J = \sum u_i \otimes v_i \in (\mathcal{U}(\mathfrak{a} \oplus \mathfrak{a}^*)[[\hbar]])^{\hat{\otimes} 2}$ with $(id \otimes \varepsilon_0)J = 1 = (\varepsilon_0 \otimes id)J$ such that when twisting [6, 7] the quasitriangular quasi-Hopf algebra of theorem 3.1 via J^{-1} one obtains a quasitriangular Hopf QUE-algebra, $(\mathcal{U}(\mathfrak{a} \oplus \mathfrak{a}^*)[[\hbar]]; \cdot; 1; \Delta; \varepsilon_0; S; R)$, which is a quantization of pair $(\mathfrak{a} \oplus \mathfrak{a}^*; r)$. The element J is*

$$J = (\phi^{-1} \otimes \phi^{-1}) \left(\Phi_{1,2,3,4}^{-1} \circ \Phi_{2,3,4} \circ \sigma_{23} \circ e^{\frac{\hbar}{2}\Omega_{23}} \circ \Phi_{2,3,4}^{-1} \circ \Phi_{1,2,3,4}(i_+ \otimes i_-)(\phi(1)) \right),$$

and when writing $Q = \sum S_0(u_i) \cdot v_i$, $u \in \mathfrak{a} \oplus \mathfrak{a}^*$ it is

$$\Delta(u) = J^{-1} \cdot \Delta_0(u) \cdot J, \quad S(u) = Q^{-1} \cdot S_0(u) \cdot Q, \quad R = \sigma(J^{-1}) \cdot e^{\frac{\hbar}{2}\Omega} \cdot J$$

and Φ verifies the following equalities

$$\Phi \cdot (\Delta_0 \otimes id)(J) \cdot (J \otimes 1) = (1 \otimes \Delta_0)(J) \cdot (1 \otimes J), \quad R = 1 \otimes 1 + \hbar r \quad \text{mod } \hbar^2.$$

This quasitriangular Hopf QUE-algebra will be denoted by $A_{(\mathfrak{a} \oplus \mathfrak{a}^*)[[\hbar]], \Omega, J^{-1}}$.

3.2. Quantization of quasitriangular Lie bialgebras. 1) *Let $(\mathfrak{a}, [,]_{\mathfrak{a}}, \varepsilon_{\mathfrak{a}} = d_c r_1)$ be a quasitriangular Lie bialgebra. Consider the following linear isomorphisms $\theta^+, \theta^- : \mathfrak{a} \otimes \mathfrak{a} \longrightarrow Hom(\mathfrak{a}^*, \mathfrak{a})$ defined by*

$$\theta^+(x \otimes y)(x^*) = x \cdot x^*(y) \quad \theta^-(x \otimes y)(x^*) = x^*(x) \cdot y.$$

Write $\mathfrak{a}_{\pm} = Im \theta^{\pm}(r_1) \subseteq \mathfrak{a}$. The notion of the rank of r_1 allows us to assert:

a) *the subspaces $Im \theta^{\pm}(r_1) = \mathfrak{a}_{\pm}$ associated with $r_1 \in \mathfrak{a} \otimes \mathfrak{a}$ are canonically isomorphic to $(Im \theta^{\mp}(r_1))^* \cong \mathfrak{a}_{\mp}^*$.*

b) *$r_1 \in \mathfrak{a}_+ \otimes \mathfrak{a}_-$ and $r_1 = \sum_i a_i \otimes b_i$, where $a_i \in \mathfrak{a}_+$ and $b_i \in \mathfrak{a}_-$, $\forall i = \{1, 2, \dots, rank(r_1)\}$.*

c) the mapping $\chi_{r_1} : \mathfrak{a}_+^* \rightarrow \mathfrak{a}_-$ defined by $\chi_{r_1}(x^*) = \sum_i x^*(a_i) \cdot b_i$ is an isomorphism.

As r_1 is a solution of the CYBE, \mathfrak{a}_+ and \mathfrak{a}_- are Lie subalgebras of \mathfrak{a} , a result from [12].

Using the isomorphism χ_{r_1} it is possible to define a Lie algebra structure on \mathfrak{a}_+^* by $[\cdot, \cdot]_{\mathfrak{a}_+^*} = \chi_{r_1}^{-1} \circ [\cdot, \cdot]_{\mathfrak{a}_-} \circ (\chi_{r_1} \otimes \chi_{r_1})$. Then we have

Theorem 3.3. *The Lie algebra structures $[\cdot, \cdot]_{\mathfrak{a}_+^*}$ on \mathfrak{a}_+^* and $[\cdot, \cdot]_{\mathfrak{a}_+}$ on \mathfrak{a}_+ are compatible in the sense of Drinfeld (see [15, 16]). The mapping $\varepsilon_{\mathfrak{a}_+} = \phi^t : \mathfrak{a}_+ \rightarrow \mathfrak{a}_+ \otimes \mathfrak{a}_+$ where $\phi(\xi_1 \otimes \xi_2) = [\xi_1; \xi_2]_{\mathfrak{a}_+^*}$ is then a 1-cocycle on $(\mathfrak{a}_+, [\cdot, \cdot]_{\mathfrak{a}_+})$. The set $(\mathfrak{a}_+, [\cdot, \cdot]_{\mathfrak{a}_+}, \varepsilon_{\mathfrak{a}_+} = \phi^t)$ is a Lie bialgebra whose quasitriangular double Lie bialgebra is $(\mathfrak{a}_+ \oplus \mathfrak{a}_+^*, [\cdot, \cdot]_{\mathfrak{a}_+ \oplus \mathfrak{a}_+^*}, \varepsilon_{\mathfrak{a}_+ \oplus \mathfrak{a}_+^*} = d_c r_+)$, where r_+ is the invariant canonical element.*

From [12, 9] it follows

Proposition 3.4. *The mapping $\tilde{\pi} : \mathfrak{a}_+ \oplus \mathfrak{a}_+^* \rightarrow \mathfrak{a}$, defined as $\tilde{\pi}(x; \xi) = x + \chi_{r_1}(\xi)$ is a Lie-bialgebra-morphism. That is, a Lie-algebra morphism verifying $d_c r_1 \circ \tilde{\pi} = (\tilde{\pi} \otimes \tilde{\pi}) \circ d_c r_+$. Moreover $(\tilde{\pi} \otimes \tilde{\pi})r_+ = r_1$. The symbol $\tilde{\pi}$ will also denote the unique algebra morphism $\tilde{\pi} : \mathcal{U}(\mathfrak{a}_+ \oplus \mathfrak{a}_+^*) \rightarrow \mathcal{U}(\mathfrak{a})$ defined by the Lie algebra morphism $\tilde{\pi}$.*

2) From theorems 3.1, 3.3 and proposition 3.4 it is possible to obtain a quantization of the pair $(\mathfrak{a}, r_1 + \sigma(r_1))$. With the obvious notations we have

Theorem 3.5. *Let $(\mathfrak{a}_+ \oplus \mathfrak{a}_+^*, [\cdot, \cdot]_{\mathfrak{a}_+ \oplus \mathfrak{a}_+^*}, \varepsilon_{\mathfrak{a}_+ \oplus \mathfrak{a}_+^*} = d_c r_+)$ be the quasitriangular Lie bialgebra in theorem 3.3. Let $(\mathcal{U}\mathfrak{a}[[\hbar]], \cdot, 1, \Delta_{\mathfrak{a}}, \epsilon_{\mathfrak{a}}, S_{\mathfrak{a}})$ be the usual Hopf universal enveloping algebra. Let*

$$\left(\mathcal{U}(\mathfrak{a}_+ \oplus \mathfrak{a}_+^*)[[\hbar]], \cdot, 1, \Delta_0^+, \epsilon_0^+, \Phi^+, S_0^+, \alpha^+ = (c^+)^{-1}, \beta^+ = 1, R_0^+ = e^{\frac{\hbar}{2}\Omega_+} \right),$$

be the quasitriangular quasi-Hopf algebra in theorem 3.1 whose classical limit is the pair $(\mathfrak{a}_+ \oplus \mathfrak{a}_+^*, \Omega_+)$. Then we have the equalities : $(\tilde{\pi} \otimes \tilde{\pi}) \circ \Delta_0^+ = \Delta_{\mathfrak{a}} \circ \tilde{\pi}$; $\tilde{\pi} \circ S_0^+ = S_{\mathfrak{a}} \circ \tilde{\pi}$ and putting $\tilde{\Phi}^+ = (\tilde{\pi} \otimes \tilde{\pi})\Phi^+$, $R_{\mathfrak{a}} = (\tilde{\pi} \otimes \tilde{\pi})R_0^+$, the set $(\mathcal{U}\mathfrak{a}[[\hbar]], \cdot, 1, \Delta_{\mathfrak{a}}, \epsilon_{\mathfrak{a}}, \tilde{\Phi}^+, S_{\mathfrak{a}}, \tilde{\alpha} = (\tilde{\pi}(c^+))^{-1}, \tilde{\beta} = 1, R_{\mathfrak{a}})$ is a quasitriangular quasi-Hopf QUE-algebra whose classical limit is the pair $(\mathfrak{a}, r_1 + \sigma(r_1))$.

The above results allow us to obtain a quantization of the pair (\mathfrak{a}, r_1) . With the obvious notations the result can be stated as follows [9] (see also [22]):

Theorem 3.6. *Let $(\mathfrak{a}, [\cdot, \cdot]_{\mathfrak{a}}, \varepsilon_{\mathfrak{a}} = d_c r_1)$ be a quasitriangular Lie bialgebra. Let $(\mathfrak{a}_+ \oplus \mathfrak{a}_+^*, [\cdot, \cdot]_{\mathfrak{a}_+ \oplus \mathfrak{a}_+^*}, \varepsilon_{\mathfrak{a}_+ \oplus \mathfrak{a}_+^*} = d_c r_+)$ be the quasitriangular Lie bialgebra in theorem 3.3. Let $(\mathcal{U}(\mathfrak{a}_+ \oplus \mathfrak{a}_+^*)[[\hbar]], \cdot, 1; \varepsilon_0^+; \Delta^+, S^+, R^+)$ be the quasitriangular Hopf QUE algebra obtained as in theorem 3.2, 3). Write $\tilde{J}_{r_1}^+ = (\tilde{\pi} \otimes \tilde{\pi})J^+$. We have the equality*

$$\tilde{\Phi}_{1,2,3}^+ \cdot (\Delta_{\mathfrak{a}} \otimes id) \tilde{J}_{r_1}^+ \cdot (\tilde{J}_{r_1}^+ \otimes 1) = (id \otimes \Delta_{\mathfrak{a}}) \tilde{J}_{r_1}^+ \cdot (1 \otimes \tilde{J}_{r_1}^+).$$

Write again $\tilde{J}_{r_1} = \sum p_i \otimes q_i$; $a \in \mathcal{U}\mathfrak{a}$, $\tilde{Q} = \sum S_{\mathfrak{a}}(p_i) \cdot q_i$. The set $(\mathcal{U}\mathfrak{a}[[\hbar]], \cdot, 1, \tilde{\Delta}, \tilde{\varepsilon} = \varepsilon_{\mathfrak{a}}, \tilde{S}, \tilde{R})$ where

$$\tilde{\Delta}(a) = (\tilde{J}_{r_1}^+)^{-1} \cdot \Delta_{\mathfrak{a}}(a) \cdot \tilde{J}_{r_1}^+; \varepsilon_{\mathfrak{a}}; \tilde{R} = (\tilde{\pi} \otimes \tilde{\pi}) R^+; \tilde{S}(a) = \tilde{Q}^{-1} \cdot S_{\mathfrak{a}}(a) \cdot \tilde{Q}$$

is a quasitriangular Hopf QUE-algebra which is a quantization of the pair $(\mathfrak{a}; r_1)$, and has been obtained by a twist, [6], via the element $(\tilde{J}_{r_1}^+)^{-1}$ from the quasitriangular quasi-Hopf algebra in theorem 3.5.

4. Quantization of non-degenerate triangular Lie bialgebras

1) In case of a non-degenerate triangular Lie bialgebra $(\mathfrak{a}, [\cdot, \cdot]_{\mathfrak{a}}, \varepsilon_{\mathfrak{a}} = d_c r_1)$, we have $\text{rank}(r_1) = \dim \mathfrak{a}$, $r_1 \in \wedge^2(\mathfrak{a})$. Also $\mathfrak{a}_+ = \mathfrak{a}_- = \mathfrak{a}$ as Lie algebras, $(\tilde{\pi} \otimes \tilde{\pi})\Omega_+ = (\tilde{\pi} \otimes \tilde{\pi})(r_1 + \sigma(r_1)) = 0$ and we get the equality $\varepsilon_{\mathfrak{a}_+} = \varepsilon_{\mathfrak{a}} = d_c r_1$.

Definition 4.1. [1, 5, 19, 17]) *An ISP on a non-degenerate triangular Lie bialgebra $(\mathfrak{a}, [\cdot, \cdot]_{\mathfrak{a}}, \varepsilon_{\mathfrak{a}} = d_c r_1)$ is any element $F = \sum_0^{\infty} F_k \cdot \hbar^k \in \mathcal{U}\mathfrak{a}^{\otimes 2}[[\hbar]]$ verifying the following equalities:*

- 1) $(\varepsilon_{\mathfrak{a}} \otimes id)F = (id \otimes \varepsilon_{\mathfrak{a}})F = 1 \otimes 1$;
- 2) $F - \sigma(F) = r_1 \hbar \text{ mod } \hbar^2$;
- 3) $(\Delta_{\mathfrak{a}} \otimes 1)F \cdot (F \otimes 1) = (1 \otimes \Delta_{\mathfrak{a}})F \cdot (1 \otimes F)$.

Theorem 3.6 allows us to obtain an ISP on any non-degenerate triangular Lie bialgebra. In fact it allows us to obtain, modulo equivalence, all of them, as we will see in section 6.

Theorem 4.2. *Suppose in Theorem 3.6 that $(\mathfrak{a}, [\cdot, \cdot]_{\mathfrak{a}}, \varepsilon_{\mathfrak{a}} = d_{c r_1})$ is a non-degenerate triangular Lie bialgebra. Then*

- $$\begin{aligned} \tilde{\Phi} &= (\tilde{\pi} \otimes \tilde{\pi} \otimes \tilde{\pi})\Phi = 1 \otimes 1 \otimes 1. \\ 1) \quad &(\varepsilon_{\mathfrak{a}} \otimes id)\tilde{J}_{r_1} = (id \otimes \varepsilon_{\mathfrak{a}})\tilde{J}_{r_1} = 1 \otimes 1. \\ 2) \quad &\tilde{J}_{r_1} = 1 \otimes 1 + \frac{1}{2}r_1\hbar + \dots \\ 3) \quad &(\Delta_{\mathfrak{a}} \otimes 1)\tilde{J}_{r_1} \cdot (\tilde{J}_{r_1} \otimes 1) = (1 \otimes \Delta_{\mathfrak{a}})\tilde{J}_{r_1} \cdot (1 \otimes \tilde{J}_{r_1}). \\ 4) \quad &\tilde{R} = (\tilde{\pi} \otimes \tilde{\pi})R = \sigma(\tilde{J}_{r_1}^{-1}) \cdot (1 \otimes 1) \cdot \tilde{J}_{r_1} = 1 \otimes 1 + r_1\hbar + \dots \end{aligned}$$

In particular \tilde{J}_{r_1} is an ISP on $(\mathfrak{a}, [\cdot, \cdot]_{\mathfrak{a}}, \varepsilon_{\mathfrak{a}} = d_{c r_1})$.

The triangular Hopf QUE algebra $(\mathcal{U}\mathfrak{a}[[\hbar]], \cdot, 1, \tilde{\Delta}, \tilde{\varepsilon} = \varepsilon_{\mathfrak{a}}, \tilde{S}, \tilde{R})$, denoted by $A_{\mathfrak{a}[[\hbar]], \tilde{J}_{r_1}^{-1}}$, which is obtained by a twist via $\tilde{J}_{r_1}^{-1}$ from the trivial triangular Hopf QUE algebra $(\mathcal{U}\mathfrak{a}[[\hbar]], \cdot, 1, \Delta_{\mathfrak{a}}, \varepsilon_{\mathfrak{a}}, S_{\mathfrak{a}}, R_{\mathfrak{a}} = 1 \otimes 1)$ is a quantization of the pair $(\mathfrak{a}; r_1)$.

2) The above proposition shows how to obtain an ISP on (\mathbf{G}, β_t) as in 3) Section 2. The following proposition will show that if we put \hbar in place of t in this star product we have again an ISP but this time on the Lie group (\mathbf{G}, β_1) . In this way we don't get all the ISPS on (\mathbf{G}, β_1) but if we now replace the above r_{\hbar} coming from r_t by any element in $(\mathfrak{a} \wedge \mathfrak{a})[[\hbar]]$ of the form $r_{\hbar} = r_1 + \dots \in (\mathfrak{a} \wedge \mathfrak{a})[[\hbar]]$ we obtain, up to equivalence, all the (ISP's) on (\mathbf{G}, β_1) .

Proposition 4.3. *Let $\left((\mathfrak{a} \oplus \mathfrak{a}_{r_{\hbar}}^*)[[\hbar]], [\cdot, \cdot]_{\mathfrak{a} \oplus \mathfrak{a}_{r_{\hbar}}^*}, \varepsilon_{\mathfrak{a} \oplus \mathfrak{a}_{r_{\hbar}}^*} = d_{c r} \right)$ and $r_{\hbar} \in (\mathfrak{a} \wedge \mathfrak{a})[[\hbar]]$ be as in Section 2, 4) . For any $N \in \mathbb{N}$ there exists an analytic function in a neighborhood of $t = 0$, $r_t^N = r_1 + r_2 t + r_3 t^2 + \dots \in \mathfrak{a} \wedge \mathfrak{a}$ and a solution of YBE on $(\mathfrak{a}, [\cdot, \cdot]_{\mathfrak{a}})$, such that when replacing t by \hbar in the above series expansion r_t^N and r_{\hbar} coincide up to order N .*

Proof:

In the Poisson cohomology on the Lie algebra $(\mathfrak{a}[[\hbar]], [\cdot, \cdot]_{\mathfrak{a}[[\hbar]]})$ the element $\beta_{\hbar} = \beta_1 + \beta_2 \hbar + \beta_3 \hbar^2 + \dots \in (\mathfrak{a}^ \wedge \mathfrak{a}^*)[[\hbar]]$ defined as $\beta_{\hbar} = \mu_{r_{\hbar}}(r_{\hbar})$ is a Chevalley 2-cocycle. The polynomial $\beta_t^N = \beta_1 + \beta_2 t + \beta_3 t^2 + \dots + \beta_{N-1} t^{N-1} \in \mathfrak{a}^* \wedge \mathfrak{a}^*$ is also a Chevalley 2-cocycle on $(\mathfrak{a}, [\cdot, \cdot]_{\mathfrak{a}})$. The corresponding element $r_t^N = r_1 + r_2 t + \dots$ satisfies YBE on $(\mathfrak{a}, [\cdot, \cdot]_{\mathfrak{a}})$ and when we replace \hbar in place of t it coincides with r_{\hbar} up to term N . ■*

3) From theorem 3.1 we may deduce, in the obvious notations,

Theorem 4.4. *Let $(\mathfrak{a} \oplus \mathfrak{a}_{r_t}^*, [,]_{\mathfrak{a} \oplus \mathfrak{a}_{r_t}^*}, \varepsilon_{\mathfrak{a} \oplus \mathfrak{a}_{r_t}^*} = d_c r)$ over \mathbb{R} be as in Section 2, 3). The set*

$$(\mathcal{U}(\mathfrak{a} \oplus \mathfrak{a}_{r_t}^*[[\hbar]], \cdot_t, 1, \Delta_0^t, \epsilon_0^t, \Phi_{r_t}, S_0^t, \alpha^t = c_t^{-1}, \beta^t = 1, R_0^t = e^{\frac{\hbar}{2}\Omega_t}),$$

where $c_t = \sum_i X_i \cdot_t S_0^t(Y_i) \cdot_t Z_i$, with $\Phi_{r_t} = \sum_i X_i^t \otimes Y_i^t \otimes Z_i^t$, is a quasitriangular quasi-Hopf QUE algebra over $\mathbb{R}[[\hbar]]$ with the pair $(\mathfrak{a} \oplus \mathfrak{a}_{r_t}^*, \Omega_t)$ as its classical limit.

From theorem 4.4 and proposition 4.3 we can prove

Theorem 4.5. *Let $(\mathfrak{a} \oplus \mathfrak{a}_{r_{\hbar}}^*[[\hbar]], [,]_{\mathfrak{a} \oplus \mathfrak{a}_{r_{\hbar}}^*}, \varepsilon_{\mathfrak{a} \oplus \mathfrak{a}_{r_{\hbar}}^*} = d_c r)$ over $\mathbb{R}[[\hbar]]$ be as in Section 2, 4). The set*

$$(\mathcal{U}(\mathfrak{a} \oplus \mathfrak{a}_{r_{\hbar}}^*[[\hbar]], \cdot_{\hbar}, 1, \Delta_0^{\hbar}, \epsilon_0, \Phi_{r_{\hbar}}, S_0^{\hbar}, \alpha^{\hbar} = c_{\hbar}^{-1}, \beta^{\hbar} = 1, R_0^{\hbar} = e^{\frac{\hbar}{2}\Omega_{\hbar}})$$

is then a quasitriangular quasi-Hopf QUE algebra over $\mathbb{R}[[\hbar]]$, where $c_{\hbar} = \sum_i X_i^{\hbar} \cdot_{\hbar} S_0^{\hbar}(Y_i^{\hbar}) \cdot_{\hbar} Z_i^{\hbar}$ with $\Phi_{r_{\hbar}} = \sum_i X_i^{\hbar} \otimes Y_i^{\hbar} \otimes Z_i^{\hbar}$ and S_0^{\hbar} is the antipode of $\mathcal{U}(\mathfrak{a} \oplus \mathfrak{a}_{r_{\hbar}}^*[[\hbar]])$.

Proof:

This theorem follows from Theorem A'' in [7]. In view of the next sections we want to obtain it from theorem 4.4 by quantizing first the Lie groups (\mathbf{G}, β_t) . All the elements in the above set in the theorem are well defined with the corresponding meanings on $\mathcal{U}(\mathfrak{a} \oplus \mathfrak{a}_{r_{\hbar}}^*[[\hbar]])$ and can be seen as those obtained from the corresponding ones in theorem 4.4 if we use the full-meaning trick of putting \hbar in place of t . To prove that this set defines a quasitriangular quasi-Hopf QUE algebra over $\mathbb{R}[[\hbar]]$ we need to prove the equalities which define this structure [6, 7]. These equalities are satisfied in the case of r_t . This means that for each one of them and when an ordered basis is used in $\mathcal{U}(\mathfrak{a} \oplus \mathfrak{a}_{r_t}^*)$ an infinite set of polynomials in the components of r_t are zero. But the set of these components is characterized just by the algebraic equations characterizing a solution of YBE. As a consequence we can see that the corresponding equalities are also satisfied if we replace everywhere t by \hbar , and of course also in the products of elements in the above basis, that is r_t by the corresponding solution r_{\hbar} of YBE on $(\mathfrak{a}[[\hbar]], [,]_{\mathfrak{a}[[\hbar]]})$. Then applying Proposition 4.3 we get the theorem. \blacksquare

4) Theorem 3.2 allows us to write

Theorem 4.6. *Let $(\mathfrak{a} \oplus \mathfrak{a}_{r_t}^*, [,]_{\mathfrak{a} \oplus \mathfrak{a}_{r_t}^*}, \varepsilon_{\mathfrak{a} \oplus \mathfrak{a}_{r_t}^*} = d_c r)$ over \mathbb{R} be as in Section 2, 3). Write $J_{r_t} = \sum_i u_i \otimes v_i \in \mathcal{U}(\mathfrak{a} \oplus \mathfrak{a}_{r_t}^*)^{\otimes 2}[[\hbar]]$ where*

$$J_{r_t} = (\phi_t^{-1} \otimes \phi_t^{-1}) \left((\Phi_{r_t}^{-1})_{1,2,3,4} \circ (\Phi_{r_t})_{2,3,4} \circ \sigma_{23} \circ e^{\frac{\hbar}{2}(\Omega_t)_{23}} \circ (\Phi_{r_t}^{-1})_{2,3,4} \circ (\Phi_{r_t})_{1,2,3,4} \circ (i_+ \otimes i_-) \circ \phi_t(1) \right)$$

is the corresponding element to the one introduced in theorem 3.2, part 3). In the present case we have written $\phi_t : \mathcal{U}(\mathfrak{a} \oplus \mathfrak{a}_{r_t}^) \longrightarrow M_+^{r_t} \otimes M_-^{r_t}$ for the morphism ϕ . The element J_{r_t} satisfies the equalities $(id \otimes \epsilon_0^t) J_{r_t} = 1 = (\epsilon_0^t \otimes id) J_{r_t}$ and when twisting the quasitriangular quasi-Hopf QUE-algebra in theorem 4.4 via $J_{r_t}^{-1}$ one obtains a quasitriangular Hopf QUE-algebra $(\mathcal{U}(\mathfrak{a} \oplus \mathfrak{a}_{r_t}^*)[[\hbar]], \cdot_t, 1, \Delta_t, \epsilon_t \equiv \epsilon_0^t, S_t, R_t)$ over $\mathbb{R}[[\hbar]]$ whose classical limit is the pair $(\mathfrak{a} \oplus \mathfrak{a}_{r_t}^*; r_t)$ and which will be denoted by $A_{\mathfrak{a} \oplus \mathfrak{a}_{r_t}^*[[\hbar]], \Omega_t, J_{r_t}^{-1}}$. If we write $Q_t = \sum_i S_0^t(u_i) \cdot_t v_i$ the above defining elements are*

$$\Delta_t(u) = J_{r_t}^{-1} \cdot_t \Delta_0^t(u) \cdot_t J_{r_t}; \quad S_t(u) = Q_t^{-1} \cdot_t S_0^t(u) \cdot_t Q_t; \quad R_t = \sigma(J_{r_t})^{-1} \cdot_t e^{\frac{\hbar}{2}\Omega_t} \cdot_t J_{r_t},$$

and Φ_{r_t} satisfies the following equalities

$$\Phi_{r_t} \cdot_t (\Delta_0^t \otimes id)(J_{r_t}) \cdot_t (J_{r_t} \otimes 1) = (1 \otimes \Delta_0^t)(J_{r_t}) \cdot_t (1 \otimes J_{r_t}); \quad R_t = 1 \otimes 1 + \hbar r_t \quad \text{mod } \hbar^2.$$

The next theorem can be proved from theorem 4.6 in a similar way as theorem 4.5. See also Lemma 4.8.

Theorem 4.7. *The set $(\mathcal{U}(\mathfrak{a} \oplus \mathfrak{a}_{r_{\hbar}}^*)[[\hbar]], \cdot_{\hbar}, 1, \Delta_{\hbar}, \epsilon_0, S_{\hbar}, R_{\hbar})$ is a quasitriangular Hopf QUE algebra over $\mathbb{R}[[\hbar]]$. Its defining elements are the following*

$$\begin{aligned} \Delta_{\hbar}(a) &= J_{r_{\hbar}}^{-1} \cdot_{\hbar} \Delta_0^{\hbar}(a) \cdot_{\hbar} J_{r_{\hbar}}, & S_{\hbar}(a) &= Q_{\hbar}^{-1} \cdot_{\hbar} S_0^{\hbar}(a) \cdot_{\hbar} Q_{\hbar}, \\ R_{\hbar} &= \sigma(J_{r_{\hbar}}^{-1}) \cdot_{\hbar} e^{\frac{\hbar}{2}\Omega} \cdot_{\hbar} J_{r_{\hbar}}, \end{aligned}$$

where $Q_{\hbar} = \sum_i S_0^{\hbar}(p_i) \cdot_{\hbar} q_i$, $J_{r_{\hbar}} = \sum_i p_i \otimes q_i$, $a \in \mathcal{U}(\mathfrak{a} \oplus \mathfrak{a}_{r_{\hbar}}^)[[\hbar]]$ and $\Phi_{r_{\hbar}}$ satisfies the following equalities*

$$\Phi_{r_{\hbar}} \cdot_{\hbar} (\Delta_0^{\hbar} \otimes id)(J_{r_{\hbar}}) \cdot_{\hbar} (J_{r_{\hbar}} \otimes 1) = (1 \otimes \Delta_0^{\hbar})(J_{r_{\hbar}}) \cdot_{\hbar} (1 \otimes J_{r_{\hbar}}); \quad R_{\hbar} = 1 \otimes 1 + \hbar r_{\hbar} \quad \text{mod } \hbar^2.$$

This quasitriangular Hopf QUE algebra over $\mathbb{R}[[\hbar]]$ is therefore obtained from the quasitriangular quasi-Hopf algebra in theorem 4.5 via the element $J_{r_{\hbar}}^{-1} \in \mathcal{U}(\widehat{\mathfrak{a} \oplus \mathfrak{a}_{r_{\hbar}}^}) \hat{\otimes} \mathcal{U}(\widehat{\mathfrak{a} \oplus \mathfrak{a}_{r_{\hbar}}^*})$. Its classical limit is the quasitriangular Lie bialgebra $(\mathfrak{a} \oplus \mathfrak{a}_{r_1}^*, [,]_{\mathfrak{a} \oplus \mathfrak{a}_{r_1}^*}, \varepsilon = d_c r)$. We denote it by $A_{\mathfrak{a} \oplus \mathfrak{a}_{r_{\hbar}}^*, \Omega, J_{r_{\hbar}}^{-1}}$.*

5) The following two lemmas will be needed.

Choose an ordered basis $\{e_a\}$ in \mathfrak{a} , and its dual basis $\{e^a\}$ in \mathfrak{a}^* . Then we can construct ordered bases in $\mathfrak{a} \oplus \mathfrak{a}_{r_t}^*$, $\mathcal{U}\mathfrak{a}$, $\mathcal{U}\mathfrak{a}^*$, $\mathcal{U}(\mathfrak{a} \oplus \mathfrak{a}_{r_t}^*)^{\otimes 2}$. We can prove

Lemma 4.8. [22] *Let $r_t = r_1 + r_2 t + r_3 t^2 + \dots \in \mathfrak{a} \wedge \mathfrak{a}$ be as in Section 2, 3). The element $J_{r_t} \in \mathcal{U}(\mathfrak{a} \oplus \mathfrak{a}_{r_t}^*)^{\otimes 2}[[\hbar]]$ in Theorem 4.6 can be written as*

$$J_{r_t} = 1 \otimes 1 + \frac{1}{2} r \hbar + \sum_{k \geq 2} \left(r_t^{i_1 j_1} \dots r_t^{i_{l(k)} j_{l(k)}} Q_{i_1, \dots, i_{l(k)}, j_1, \dots, j_{l(k)}, k} \right) \hbar^k,$$

where $Q_{i_1, \dots, i_{l(k)}, j_1, \dots, j_{l(k)}, k} \in \mathcal{U}(\mathfrak{a} \oplus \mathfrak{a}_{r_t}^*)^{\otimes 2}$ is a (finite) linear combination of tensor products of elements in the above ordered basis. r_t is manifested in every element of the ordered basis through the product in $\mathcal{U}(\mathfrak{a} \oplus \mathfrak{a}_{r_t}^*)$ but it does not appear in the coefficients defining $Q_{i_1, \dots, i_{l(k)}, j_1, \dots, j_{l(k)}, k}$.

Proof:

From the expression of J on theorem 3.2 and because $\Phi = 1 \otimes 1 \otimes 1 + O(\hbar^2)$ we have

$$\begin{aligned} J_{r_t} &= (\phi_t^{-1} \otimes \phi_t^{-1}) \left((1 + \frac{\hbar}{2} \Omega_{23})(1_+ \otimes 1_- \otimes 1_+ \otimes 1_-) \right) \text{ mod } \hbar^2 \\ &= ((\phi_t^{-1} \otimes \phi_t^{-1})(1_+ \otimes 1_- \otimes 1_+ \otimes 1_-) + \\ &\quad + \frac{1}{2} (\phi_t^{-1} \otimes \phi_t^{-1})(1_+ \otimes r_{12}(1_- \otimes 1_+) \otimes 1_-) \hbar) \text{ mod } \hbar^2 \\ &= \left(1 \otimes 1 + \frac{1}{2} r \hbar \right) \text{ mod } \hbar^2. \end{aligned}$$

In the expression for J_{r_t} the coefficient of \hbar^k , for $k \geq 2$, is an element in $\mathcal{U}(\mathfrak{a} \oplus \mathfrak{a}_{r_t}^*) \otimes \mathcal{U}(\mathfrak{a} \oplus \mathfrak{a}_{r_t}^*)$ which depends on r_t (by the brackets on Φ_{r_t} and by the products on $\mathcal{U}(\mathfrak{a} \oplus \mathfrak{a}_{r_t}^*)$), so is of the form

$$r_t^{i_1 j_1} \dots r_t^{i_{l(k)} j_{l(k)}} Q_{i_1, \dots, i_{l(k)}, j_1, \dots, j_{l(k)}, k},$$

where $Q_{i_1, \dots, i_{l(k)}, j_1, \dots, j_{l(k)}, k} \in \mathcal{U}(\mathfrak{a} \oplus \mathfrak{a}_{r_t}^*) \otimes \mathcal{U}(\mathfrak{a} \oplus \mathfrak{a}_{r_t}^*)$ only depends on r_t through the products on the enveloping algebra $\mathcal{U}(\mathfrak{a} \oplus \mathfrak{a}_{r_t}^*)$ in the ordered basis. \blacksquare

As the product in $\mathcal{U}\mathfrak{a}$ is independent of r_t , applying $\tilde{\pi}_t \otimes \tilde{\pi}_t$ to J_{r_t} and then putting $t = \hbar$, we can also prove

Lemma 4.9. [22] *Let $r_t = r_1 + r_2 t + r_3 t^2 + \dots \in \wedge^2(\mathfrak{a})$ be as in Section 2, 3). Write $r_l = r_l^{ab} e_a \otimes e_b$, $r_l^{ab} + r_l^{ba} = 0$, $l = 1, 2, 3, \dots$. Let $\tilde{\pi}_t$ be the Lie*

bialgebra morphism defined in proposition 3.4 and define $\tilde{J}_{r_t} = (\tilde{\pi}_t \otimes \tilde{\pi}_t)J_{r_t}$, $\tilde{J}_{r_{\hbar}} = (\tilde{J}_{r_t})|_{t \rightarrow \hbar} \in (\mathcal{U}\mathfrak{a})^{\otimes 2}[[\hbar]]$. Write it as a formal power series in \hbar . The element $\tilde{J}_{r_{\hbar}}$ can be written as

$$\tilde{J}_{r_{\hbar}} = 1 \otimes 1 + \frac{1}{2}r_1\hbar + \sum_{R=2}^{\infty} \left(\frac{1}{2}r_R + \sum_{i_2, j_2, \dots, i_R, j_R} \left(\sum_{A_{i_2, j_2}(R), \dots, A_{i_R, j_R}(R)} (r_1^{i_2 j_2})^{A_{i_2, j_2}(R)} \dots (r_{R-1}^{i_R j_R})^{A_{i_R, j_R}(R)} H_{i_2, \dots, i_R, j_2, \dots, j_R, A_{i_2, j_2}(R), \dots, A_{i_R, j_R}(R), R} \right) \right) \hbar^R,$$

where $H_{i_2, \dots, i_R, j_2, \dots, j_R, A_{i_2, j_2}(R), \dots, A_{i_R, j_R}(R), R} \in (\mathcal{U}\mathfrak{a})^{\otimes 2}$ is a (finite) linear combination of tensor products of elements in the above ordered basis and it is independent of r_l , $l = 1, 2, 3, \dots$

6) From lemma 4.9 we obtain the following proposition and corollary which will be applied in the next section.

Proposition 4.10. [22] *Let $r_t = r_1 + r_2 t + r_3 t^2 + \dots \in \Lambda^2(\mathfrak{a})$ be as in Section 2, 3). Let $(\mathfrak{a}, [\cdot, \cdot]_{\mathfrak{a}}, \varepsilon_{\mathfrak{a}} = d_c r_t)$ be the non-degenerate triangular Lie bialgebra defined by r_t . Let $(\mathcal{U}\mathfrak{a}[[\hbar]], \cdot, \tilde{\Delta}_t, \tilde{S}_t, \tilde{R}_t)$ be the triangular Hopf QUE-algebra whose classical limit is the pair $(\mathfrak{a}; r_t)$ and was obtained in theorem 4.2 from the usual triangular Hopf algebra $(\mathcal{U}\mathfrak{a}[[\hbar]], \cdot, \Delta_{\mathfrak{a}}, S_{\mathfrak{a}}, R_{\mathfrak{a}} = 1)$ by a twist via $(\tilde{J}_{r_t})^{-1}$. Consider, as before, the element $\tilde{J}_{r_{\hbar}} = (\tilde{J}_{r_t})|_{t \rightarrow \hbar} \in \mathcal{U}\mathfrak{a}[[\hbar]] \hat{\otimes} \mathcal{U}\mathfrak{a}[[\hbar]]$. Then the following equalities hold:*

- (a) $(\Delta_{\mathfrak{a}} \otimes id) \tilde{J}_{r_{\hbar}} \cdot_{\hbar} (\tilde{J}_{r_{\hbar}} \otimes 1) = (1 \otimes \Delta_{\mathfrak{a}}) \tilde{J}_{r_{\hbar}} \cdot_{\hbar} (id \otimes \tilde{J}_{r_{\hbar}})$;
- (b) $\tilde{J}_{r_{\hbar}} = 1 \otimes 1 + \frac{1}{2}r_1 \hbar + 0(\hbar^2)$.

Let us define $\Delta(a) = (\tilde{J}_{r_{\hbar}})^{-1} \cdot_{\hbar} \Delta_{\mathfrak{a}}(a) \cdot_{\hbar} \tilde{J}_{r_{\hbar}}$, $R = (\sigma \tilde{J}_{r_{\hbar}})^{-1} \cdot_{\hbar} (1 \otimes 1) \cdot_{\hbar} \tilde{J}_{r_{\hbar}}$ and $S(a) = Q^{-1} \cdot_{\hbar} S_{\mathfrak{a}}(a) \cdot_{\hbar} Q$, where $Q = \sum S_{\mathfrak{a}}(a_i) \cdot_{\hbar} b_i$, $\tilde{J}_{r_{\hbar}} = \sum a_i \otimes b_i$; $a_i, b_i \in \mathcal{U}\mathfrak{a}[[\hbar]]$. The set $(\mathcal{U}\mathfrak{a}[[\hbar]], \cdot_{\hbar}, \Delta, S, R)$ is then a triangular Hopf QUE algebra obtained twisting the usual triangular Hopf algebra $(\mathcal{U}\mathfrak{a}[[\hbar]], \cdot, \Delta_{\mathfrak{a}}, S_{\mathfrak{a}}, R_{\mathfrak{a}} = 1 \otimes 1)$ via the element $(\tilde{J}_{r_{\hbar}})^{-1} \in \mathcal{U}\mathfrak{a}[[\hbar]] \hat{\otimes} \mathcal{U}\mathfrak{a}[[\hbar]]$. We write it as $A_{\mathfrak{a}[[\hbar]], (\tilde{J}_{r_{\hbar}})^{-1}}$.

Corollary 4.11. *Let $r'_t = r_1 + r_2 t + r_3 t^2 + \dots + r_{k-1} t^{k-2} + (r_k + s_k) t^{k-1} + \dots \in \Lambda^2(\mathfrak{a})$ be another element. Let $\tilde{J}'_{r_{\hbar}}$ be the star product determined by r'_t in the similar way as $\tilde{J}_{r_{\hbar}}$ was from r_t . Then $\tilde{J}_{r_{\hbar}}$ and $\tilde{J}'_{r_{\hbar}}$ coincide up to order $k - 1$ and*

$$\left(\tilde{J}'_{r_{\hbar}} \right)_k - \left(\tilde{J}_{r_{\hbar}} \right)_k = \frac{1}{2} s_k.$$

5. ISP F determines $r_{\hbar} \in (\mathfrak{a} \otimes \mathfrak{a})[[\hbar]]$. $\tilde{J}_{r_{\hbar}}^{\Phi}$ and F are equivalent.

Let $F \in \mathcal{U}\mathfrak{a}[[\hbar]] \hat{\otimes} \mathcal{U}\mathfrak{a}[[\hbar]]$ be an ISP on the non-degenerate triangular Lie bialgebra $(\mathfrak{a}, [\cdot, \cdot]_{\mathfrak{a}}, \varepsilon_{\mathfrak{a}} = d_c r_1)$. Let $A_{\mathfrak{a}[[\hbar]], F^{-1}}$ be the triangular Hopf QUE algebra obtained by a twist via F^{-1} from the trivial triangular Hopf QUE algebra $(\mathcal{U}(\mathfrak{a})[[\hbar]], \cdot, 1, \Delta_{\mathfrak{a}}, S_{\mathfrak{a}}, R_{\mathfrak{a}} = 1 \otimes 1)$. It is then a quantization of the pair (\mathfrak{a}, r_1) .

The following proposition does not depend on any specific context of quantization [19] but only on the notion of deformation of associative algebras [10], the fact that Hochschild cohomology spaces of coalgebra $(\mathcal{U}\mathfrak{a}, \mathbb{R})$ are $H^k(\mathcal{U}\mathfrak{a}) = \Lambda^k \mathfrak{a}$, $k \in \mathbb{N}$, [3], and the Hochschild cohomological [18] interpretation of the Quantum Yang Baxter equation.

Proposition 5.1. [18] *Let $F = \sum_i^{\infty} F_i \hbar^i$ and $F' = \sum_i^{\infty} F'_i \hbar^i$ be ISP on $(\mathfrak{a}, [\cdot, \cdot]_{\mathfrak{a}}, \varepsilon_{\mathfrak{a}} = d_c r_1)$. Let $A_{\mathfrak{a}[[\hbar]], F^{-1}}$ and $A_{\mathfrak{a}[[\hbar]], F'^{-1}}$ be as before in this section. Suppose that F and F' coincide up to order k , i.e. $F'_l = F_l$, $l = 1, 2, \dots, k$. Then: a) there exist $h_{k+1} \in \mathfrak{a} \wedge \mathfrak{a}$ and $E_{k+1} \in \mathcal{U}\mathfrak{a}$ such that $F'_{k+1} - F_{k+1} = h_{k+1} + d_H E_{k+1}$ where d_H is the coboundary operator in the Hochschild cohomology of $\mathcal{U}\mathfrak{a}$; b) h_{k+1} is not only a Hochschild 2-cocycle but also a Poisson 2-cocycle relative to the invariant Poisson structure defined by the element $r_1 \in \mathfrak{a} \wedge \mathfrak{a}$.*

Again, the above Hochschild cohomology spaces and proposition 5.1 play a central role in the proof of the next theorem. In the context of quantification in [9] the next theorem corresponds to a main theorem by Drinfeld in the context of quantification in [5] and in [19, 20] there is a proof of this Drinfeld theorem. See the References in [20] for a similar theorem about Star Products on general symplectic manifolds and [13] on Poisson manifolds.

Theorem 5.2. *Fix a Lie associator Φ . Let $A_{\mathfrak{a}[[\hbar]], F^{-1}}$ be as defined at the beginning of this section. We have:*

(a) *There exist elements $r_{\hbar} = r_1 + r_2 \hbar + r_3 \hbar^2 + \dots \in (\wedge^2 \mathfrak{a})[[\hbar]]$ and $E^{r_{\hbar}} = 1 + E_1^{r_{\hbar}} \hbar + \dots + E_n^{r_{\hbar}} \hbar^n + \dots \in \mathcal{U}\mathfrak{a}[[\hbar]]$ such that*

$$F = \Delta_{\mathfrak{a}}((E^{r_{\hbar}})^{-1}) \cdot_{\hbar} \tilde{J}_{r_{\hbar}}^{\Phi} \cdot_{\hbar} (E^{r_{\hbar}} \otimes E^{r_{\hbar}});$$

i.e., F and $\tilde{J}_{r_{\hbar}}^{\Phi}$ are equivalent ISPS over the non-degenerate triangular Lie bialgebra $(\mathfrak{a}, [\cdot, \cdot]_{\mathfrak{a}}, \varepsilon_{\mathfrak{a}} = d_c r_1)$.

(b) The triangular Hopf QUE algebras $A_{\mathfrak{a}[[\hbar]],F^{-1}}$ and $A_{\mathfrak{a}[[\hbar]],(\tilde{J}_{r\hbar}^\Phi)^{-1}}$ are isomorphic.

Proof:

(a) We construct an analytic function

$$r_t(n) = r_1 + \sum_{l=2}^{\infty} r_l(n)t^{l-1} \in \wedge^2(\mathfrak{a})$$

which is a non-degenerate solution of CYBE and we prove that $\tilde{J}_{r\hbar(n)}^\Phi$ and F are equivalent at order n , $n \in \mathbb{N}$, by induction on the order of equivalence. The results comes from that equivalence.

(i) Let $\tilde{J}_{r\hbar(1)}$ be the star product obtained from the E-K quantization (proposition 4.10), determined by the CYBE solution $r_t(1) = r_1$.

As $A_{\mathfrak{a}[[\hbar]],F^{-1}}$ is a quantization of the given Lie bialgebra $R = 1 \otimes 1 + r_1\hbar + \dots$, and because it is the twist of the usual triangular Hopf QUE algebra ($R_{\mathfrak{a}} = 1 \otimes 1$) we have

$$F_1 - \sigma F_1 = r_1 \tag{5.1}$$

and, by construction,

$$(\tilde{J}_{r\hbar(1)})_1 = \frac{1}{2}r_1. \tag{5.2}$$

The associative property at order 1 for F and $\tilde{J}_{r\hbar(1)}$ is $d_H F_1 = 0$ and $d_H(\tilde{J}_{r\hbar(1)})_1 = 0$, d_H being the Hochschild cohomology operator. Therefore, $d_H((\tilde{J}_{r\hbar(1)})_1 - F_1) = 0$, i.e., $(\tilde{J}_{r\hbar(1)})_1 - F_1$ is a Hochschild 2-cocycle. The 2-cocycle condition implies that there exist $h_1 \in \mathfrak{a} \wedge \mathfrak{a}$ and $E_1 \in \mathcal{U}\mathfrak{a}$ such that

$$(\tilde{J}_{r\hbar(1)})_1 - F_1 = h_1 + d_H E_1. \tag{5.3}$$

On the other hand, from (5.1) and (5.2), and because $(\tilde{J}_{r\hbar(1)})_1$ is skew-symmetric, we conclude that $F_1 - (\tilde{J}_{r\hbar(1)})_1 = \sigma(F_1 - (\tilde{J}_{r\hbar(1)})_1)$, i.e., $F_1 - (\tilde{J}_{r\hbar(1)})_1$ is symmetric. As h_1 is the skew-symmetric part in (5.3), we get $h_1 = 0$ and

$$(\tilde{J}_{r\hbar(1)})_1 - F_1 = d_H E_1. \tag{5.4}$$

$\tilde{J}_{r\hbar(1)}$ and F are equivalent to order 1 ($E^{r\hbar(1)} = 1 + E_1\hbar$).

(ii) $\tilde{J}_{r\hbar(1)}$ and F being equivalent to order 1, we know (Gerstenhaber [10]) that $(\tilde{J}_{r\hbar(1)})_2 - F_2 + G_2(E_1, (\tilde{J}_{r\hbar(1)})_1, F_1)$ is a Hochschild 2-cocycle. Then, there

exist $h_2 \in \mathfrak{a} \wedge \mathfrak{a}$ and $E_2 \in \mathcal{U}\mathfrak{a}$ such that

$$(\tilde{J}_{r_h(1)})_2 - F_2 + G_2(E_1, (\tilde{J}_{r_h(1)})_1, F_1) = \frac{1}{2}h_2 + d_H E_2. \quad (5.5)$$

Put $E^{r_h(2)} = 1 + E_1\hbar + E_2\hbar^2 \in \mathcal{U}\mathfrak{a}[[\hbar]]$ and consider the star product, equivalent to F , $F'^{(1)} = \Delta_{\mathfrak{a}}((E^{r_h(2)})^{-1}) \cdot F \cdot (E^{r_h(2)} \otimes E^{r_h(2)})$.

At first order, the equivalence condition is

$$F_1'^{(1)} - F_1 = d_H E_1, \quad (5.6)$$

and, from (5.4), we conclude that $F_1'^{(1)} = (\tilde{J}_{r_h(1)})_1$.

At order 2, the equivalence can be written in the form

$$F_2'^{(1)} - F_2 + G_2(E_1, F_1'^{(1)}, F_1) = d_H E_2 \quad (5.7)$$

but, as $F_1'^{(1)} = (\tilde{J}_{r_h(1)})_1$, we get $G_2(E_1, F_1'^{(1)}, F_1) = G_2(E_1, (\tilde{J}_{r_h(1)})_1, F_1)$ and, comparing (5.5) with (5.7), we obtain

$$F_2'^{(1)} = (\tilde{J}_{r_h(1)})_2 - \frac{1}{2}h_2. \quad (5.8)$$

As $(\tilde{J}_{r_h(1)})_1 = F_1'^{(1)}$, h_2 is a Poisson 2-cocycle ([19]), so that $\beta_2 = \mu_{r_1}(h_2)$ is an invariant De Rham (or Chevalley) 2-cocycle, where μ_{r_1} is the isomorphism defined before. We consider $\beta_1 = \mu_{r_1}(r_1) \in \mathfrak{a}^* \wedge \mathfrak{a}^*$ (equivalent to $(r_1)^{ab}(\beta_1)_{ac} = \delta_c^b$) and define $\beta_t(2) = \beta_1 + \beta_2 t$ ($d_c \beta_t = 0$ and β_1 is non-degenerate). Let us define $r_t(2) = r_1(2) + \sum_{k \geq 2} r_k(2)t^{k-1} \in \mathfrak{a} \wedge \mathfrak{a}[[t]]$, by $\mu_{r_t(2)}^{-1}(\beta_t(2)) = r_t(2)$. Then, $r_1(2) = r_1 = r_1(1)$ and $r_2(2) = -\mu_{r_1}^{-1}(\beta_2) = -h_2$ and $[r_t(2), r_t(2)] = 0$, $r_t(2)$ is a solution of the CYBE.

Let $\tilde{J}_{r_h(2)}$ be the star product determined following Etingof-Kazhdan by the element $r_t(2)$, after $t = \hbar$, as in proposition 4.10. We have, also from proposition 4.10, that $(\tilde{J}_{r_h(2)})_1 = (\tilde{J}_{r_h(1)})_1$ and $(\tilde{J}_{r_h(2)})_2 = (\tilde{J}_{r_h(1)})_2 + \frac{1}{2}r_2(2) = (\tilde{J}_{r_h(1)})_2 - \frac{1}{2}h_2$. Thus, $(\tilde{J}_{r_h(2)})_1 = F_1'^{(1)} = (\tilde{J}_{r_h(1)})_1$, and from (5.8), we have $(\tilde{J}_{r_h(2)})_2 = F_2'^{(1)}$. Replacing these equalities in the expressions (5.6) and (5.7), we get

$$\begin{aligned} (\tilde{J}_{r_h(2)})_1 - F_1 &= d_H E_1 \\ (\tilde{J}_{r_h(2)})_2 - F_2 + G_2(E_1, (\tilde{J}_{r_h(2)})_1, F_1) &= d_H E_2. \end{aligned}$$

So, F is equivalent to $\tilde{J}_{r_h(2)}$, to order 2, $\tilde{J}_{r_h(2)}$ being the star product determined by $r_t(2)$, with $\mu_{r_t(2)}^{-1}(\beta_t(2)) = r_t(2)$ and $\beta_t(2) = \beta_1 + \mu_{r_1}(h_2)t$.

(iii) Using the induction hypothesis, and following the same steps, we conclude the proof.

(b) this part follows from part a) and [7] page 841, Remark 2. \blacksquare

As a consequence we have the following isomorphisms (see also [21]):

Corollary 5.3. *Let Φ, Φ' be two Lie associators. Let $A_{\mathfrak{a}[[\hbar]], F^{-1}}$ be given as in the theorem. Let $r_{\hbar}, r'_{\hbar} \in (\wedge^2 \mathfrak{a})[[\hbar]]$ the elements respectively determined in the theorem by the pairs $(\Phi; A_{\mathfrak{a}[[\hbar]], F^{-1}})$ and $(\Phi'; A_{\mathfrak{a}[[\hbar]], F^{-1}})$. Then we have*

$$A_{\mathfrak{a}[[\hbar]], F^{-1}} \stackrel{isom}{\approx} A_{\mathfrak{a}[[\hbar]], (\tilde{J}_{r_{\hbar}}^{\Phi})^{-1}} \stackrel{isom}{\approx} A_{\mathfrak{a}[[\hbar]], (\tilde{J}_{r'_{\hbar}}^{\Phi'})^{-1}}$$

6. Invariant star products on $(\mathfrak{a}, [,]_{\mathfrak{a}}, \varepsilon_{\mathfrak{a}} = d_c r_1)$

1) We now develop what we wrote in 3) at the Introduction. We need the following proposition:

Proposition 6.1. [22] *Let Γ be a set. Let $(\mathfrak{a}, [,]_{\mathfrak{a}}, \varepsilon = d_c r_s)$ be a non-degenerate triangular Lie bialgebra, $s \in \Gamma$. Let $\varphi_s^1 : \mathfrak{a} \rightarrow \mathfrak{a}$ a Lie algebra isomorphism $\forall s \in \Gamma$. Let r'_s be the element in $\mathfrak{a} \wedge \mathfrak{a}$ defined by $r'_s = (\varphi_s^1 \otimes \varphi_s^1) r_s$.*

- a) *The set $(\mathfrak{a}, [,]_{\mathfrak{a}}, \varepsilon'_a = d_c r'_s)$ is a non-degenerate triangular Lie bialgebra.*
- b) *The transposed map $(\varphi_s^1)^t : \mathfrak{a}_{r'_s}^* \rightarrow \mathfrak{a}_{r_s}^*$ is a Lie algebra isomorphism.*
- c) *The pair $(\varphi_s^1; \varphi_s^2 = ((\varphi_s^1)^t)^{-1})$, $s \in \Gamma$, defines a Lie bialgebra isomorphism between the Lie bialgebra $\mathfrak{a} \oplus \mathfrak{a}_{r_s}^*$ (the classical double of the Lie bialgebra $(\mathfrak{a}, [,]_{\mathfrak{a}}, \varepsilon_{\mathfrak{a}} = d_c r_s)$) and the Lie bialgebra $\mathfrak{a} \oplus \mathfrak{a}_{r'_s}^*$ (the classical double of the Lie bialgebra $(\mathfrak{a}, [,]_{\mathfrak{a}}, \varepsilon'_a = d_c r'_s)$). Furthermore, this isomorphism sends the canonical element r into itself.*

Corollary 6.2. [22] a) *Under the hypothesis of the proposition let $\beta_s = \mu_{r_s}(r_s)$, $\beta'_s = \mu_{r'_s}(r'_s)$. Then $(\varphi_s^2 \otimes \varphi_s^2) \beta_s = \beta'_s$.*

b) *Conversely, let $(\mathfrak{a}, [,]_{\mathfrak{a}}, \varepsilon = d_c r_s)$ and $(\mathfrak{a}, [,]_{\mathfrak{a}}, \varepsilon'_a = d_c r'_s)$ be non-degenerate triangular Lie bialgebras. Let β_s and β'_s be as in a). Let $\varphi_s^1 : \mathfrak{a} \rightarrow \mathfrak{a}$ be a Lie algebra isomorphism and $\varphi_s^2 = ((\varphi_s^1)^t)^{-1}$. Suppose that $(\varphi_s^2 \otimes \varphi_s^2) \beta_s = \beta'_s$. Then, $(\varphi_s^1 \otimes \varphi_s^1) r_s = r'_s$.*

2) Let $(\mathfrak{a}, [,]_{\mathfrak{a}})$ be a Lie algebra over \mathbb{R} . Consider the following Lie-algebra isomorphisms: $\varphi_t^1 = \exp(t \cdot ad_{X_t})$ where $X_t = X_1 + X_2 t + X_3 t^2 + \dots \in \mathfrak{a}$ is an analytic function in a neighborhood of $0 \in \mathbb{R}$. Then $\varphi_t^2 = \exp(-t \cdot ad_{X_t}^t) = \exp(t \cdot ad_{X_t}^*)$. Our interest is in the map $\varphi_t^2 \otimes \varphi_t^2 = \exp(ad_{X_t}^* \otimes 1 + 1 \otimes ad_{X_t}^*)^{\otimes 2}$. We have $\varphi_t^2 \otimes \varphi_t^2 = \exp(ad_{X_t}^* \otimes 1 + 1 \otimes ad_{X_t}^*)$ and then

Proposition 6.3. *Let $\beta_t = \beta_1 + \beta_2 t + \beta_3 t^2 + \dots \in \wedge^2(\mathfrak{a}^*)$ be an analytic function in a neighborhood of $0 \in \mathbb{R}$ defining a non-degenerate 2-cocycle $\forall t$. $\beta_1, \beta_2, \dots \in \wedge^2(\mathfrak{a}^*)$ are then 2-cocycles and β_1 is non-degenerate. Let X_t be as before. Then we obtain*

$$\exp(\text{ad}_{tX_t}^*)^{\otimes 2}(\beta_t) = \exp(\text{ad}_{tX_t}^* \otimes 1 + 1 \otimes \text{ad}_{tX_t}^*)(\beta_t) = \beta_t + d_c \alpha_t,$$

where $\alpha_t = \alpha_1 t + \alpha_2 t^2 + \alpha_3 t^3 + \dots \in \mathfrak{a}^*$ is an analytic function in a neighborhood of $0 \in \mathbb{R}$ given by

$$\alpha_k = \sum_{j=1}^k \left(\sum_{i=1}^{k-j+1} \frac{1}{i!} \sum_{\substack{a_1 + \dots + a_i = k-j+1 \\ a_1, \dots, a_i \geq 1}} ((-1)^{i+1} (i_{X_{a_1}} \beta_j) \cdot \text{ad}_{X_{a_2}} \cdots \text{ad}_{X_{a_i}}) \right).$$

Proof:

The first terms of the series are:

$$\begin{aligned} \exp(\text{ad}_{tX_t}^* \otimes 1 + 1 \otimes \text{ad}_{tX_t}^*)(\beta_1) &= \beta_1 + (1 \otimes \text{ad}_{X_1}^* + \text{ad}_{X_1}^* \otimes 1)(\beta_1)t + \\ &+ \left((1 \otimes \text{ad}_{X_2}^* + \text{ad}_{X_2}^* \otimes 1) + \frac{1}{2!} (1 \otimes \text{ad}_{X_1}^* + \text{ad}_{X_1}^* \otimes 1)^2 \right) (\beta_1)t^2 + \dots, \end{aligned}$$

because

$$\begin{aligned} \exp(\text{ad}_{tX_t}^* \otimes 1 + 1 \otimes \text{ad}_{tX_t}^*) &= \\ &= 1 \otimes 1 + \sum_{p \geq 1} \left(\sum_{i=1}^p \frac{1}{i!} \sum_{\substack{a_1 + \dots + a_i = p \\ a_1, \dots, a_i \geq 1}} \prod_{j=1}^i (1 \otimes \text{ad}_{X_{a_j}}^* + \text{ad}_{X_{a_j}}^* \otimes 1) \right) t^p. \end{aligned} \quad (6.9)$$

For any $e_a, e_b \in \mathfrak{a}$ elements in a basis, we have

$$\begin{aligned} \langle (1 \otimes \text{ad}_{X_1}^* + \text{ad}_{X_1}^* \otimes 1)(\beta_1); e_a \otimes e_b \rangle &= -\langle \beta_1; (1 \otimes \text{ad}_{X_1} + \text{ad}_{X_1} \otimes 1)e_a \otimes e_b \rangle \\ &= -\langle \beta_1; e_a \otimes [X_1, e_b] + [X_1, e_a] \otimes e_b \rangle \\ &= -X_1^i C_{ib}^t(\beta_1)_{at} - X_1^i C_{ia}^k(\beta_1)_{kb} \\ &= X_1^i C_{ab}^k(\beta_1)_{ki} \\ &= (-i_{X_1} \beta_1)([e_a, e_b]) \\ &= \langle d_c(i_{X_1} \beta_1); e_a \otimes e_b \rangle, \end{aligned}$$

where we used that β_1 is a 2-cocycle ($\beta_1([x, y], z) + \beta_1([y, z], x) + \beta_1([z, x], y) = 0$, $x, y, z \in \mathfrak{a}$), C_{ab}^k are the structure constants of the Lie algebra \mathfrak{a} in a basis $\{e_i\}$ and $d_c \alpha(e_a \otimes e_b) = -\alpha([e_a, e_b])$.

So, we obtain

$$(1 \otimes ad_{X_1}^* + ad_{X_1}^* \otimes 1)(\beta_1) = d(i_{X_1}\beta_1), \quad (6.10)$$

for any cocycle β_1 and any $X_1 \in \mathfrak{a}$.

With a similar computation, for any $X_1, X_2 \in \mathfrak{a}$, we get

$$(1 \otimes ad_{X_2}^* + ad_{X_2}^* \otimes 1) \cdot (1 \otimes ad_{X_1}^* + ad_{X_1}^* \otimes 1)(\beta_1) = d_c(-(i_{X_1}\beta_1) \cdot ad_{X_2}). \quad (6.11)$$

By (6.10) and (6.11), we have

$$\begin{aligned} \exp(ad_{tX}^* \otimes 1 + 1 \otimes ad_{tX}^*)(\beta_1) &= \beta_1 + d_c(i_{X_1}\beta_1)t + \\ &\quad + \left(d_c(i_{X_2}\beta_1) + \frac{1}{2!}d_c(-(i_{X_1}\beta_1) \cdot ad_{X_1}) \right) t^2 + \dots \end{aligned}$$

By induction it is possible to prove that, for any $n, X_1, X_2, \dots, X_n \in \mathfrak{a}$ and any cocycle β_1 ,

$$\begin{aligned} (1 \otimes ad_{X_n}^* + ad_{X_n}^* \otimes 1) \cdots (1 \otimes ad_{X_2}^* + ad_{X_2}^* \otimes 1) \cdot (1 \otimes ad_{X_1}^* + ad_{X_1}^* \otimes 1)\beta_1 &= \\ = d_c \left((-1)^{n+1} (i_{X_1}\beta_1) \cdot ad_{X_2} \cdots \cdots ad_{X_{n-1}} \cdot ad_{X_n} \right). \end{aligned}$$

With this result, and from (6.9), it is clear that

$$\exp(ad_{tX_t}^* \otimes 1 + 1 \otimes ad_{tX_t}^*)(\beta_1) = \beta_1 + d_c\gamma_t,$$

where $\gamma_t = \gamma_1 t + \gamma_2 t^2 + \dots$ is the 1-cochain given by

$$\gamma_k = \sum_{i=1}^k \frac{1}{i!} \sum_{\substack{a_1 + \dots + a_i = k \\ a_1, \dots, a_i \geq 1}} \left((-1)^{i+1} (i_{X_{a_1}}\beta_1) \cdot ad_{X_{a_2}} \cdots \cdots ad_{X_{a_i}} \right).$$

Since, for each $t, X_t \in \mathfrak{a}$ and $\exp(ad_{tX_t}) : \mathfrak{a} \longrightarrow \mathfrak{a}$, we have $\exp(ad_{tX_t}^*) : \mathfrak{a}^* \longrightarrow \mathfrak{a}^*$, i.e., $\beta_1 + d_c\gamma_t \in \mathfrak{a}^*$, so $d_c\gamma_t \in \mathfrak{a}^*$, for each t .

Applying this result to the cocycles $\beta_1, \beta_2, \beta_3, \dots$, we get the expression given. ■

A converse of proposition 6.3 is:

Proposition 6.4. *Let $\beta_t = \beta_1 + \beta_2 t + \beta_3 t^2 + \dots \in \wedge^2(\mathfrak{a}^*)$ as in proposition 6.3. Let $\alpha_t = \alpha_1 t + \alpha_2 t^2 + \alpha_3 t^3 + \dots \in \mathfrak{a}^*$ be an analytic function. Define $\beta'_t = \beta_t + d_c\alpha_t$. Then, there exists a unique $X_t = X_1 + X_2 t + X_3 t^2 + \dots \in \mathfrak{a}$, analytic,*

such that $\exp(\text{ad}_{tX_t}^*)^{\otimes 2}(\beta_t) = \beta'_t$, where $X_p = -(\omega_p \otimes 1)r_1 = -\chi_{r_1}(\omega_p)$, $r_1 = \mu_{r_1}^{-1}(\beta_1)$, $\omega_1 = \alpha_1$ and, for $p \geq 2$,

$$\begin{aligned} \omega_p = & \alpha_p - \sum_{i=2}^p \frac{1}{i!} \sum_{\substack{a_1+\dots+a_i=p \\ a_1, \dots, a_i \geq 1}} ((-1)^{i+1} (i_{X_{a_1}} \beta_1) \cdot \text{ad}_{X_{a_2}} \cdots \text{ad}_{X_{a_i}}) \\ & - \sum_{j=2}^p \left(\sum_{i=1}^{p-j+1} \frac{1}{i!} \sum_{\substack{a_1+\dots+a_i=p-j+1 \\ a_1, \dots, a_i \geq 1}} ((-1)^{i+1} (i_{X_{a_1}} \beta_j) \cdot \text{ad}_{X_{a_2}} \cdots \text{ad}_{X_{a_i}}) \right). \end{aligned}$$

Proof:

By the previous theorem, we know that, for any $X_t \in \mathfrak{a}$ in the above form, we have

$$\exp(\text{ad}_{tX_t}^*)^{\otimes 2}(\beta_t) = \beta_t + d_c \alpha_t,$$

where $\alpha_t = \alpha_1 t + \alpha_2 t^2 + \cdots \in \mathfrak{a}^*$ is the 1-cochain of theorem 6.3. The question now is to know if it is possible to determine $X_t = X_1 + X_2 t + X_3 t^2 + \dots$ in such a way that this 1-cochain α_t is equal to the 1-cochain given $\alpha_t = \alpha_1 t + \alpha_2 t^2 + \dots$. The elements $X_1, X_2, \dots \in \mathfrak{a}$ must satisfy

$$\begin{aligned} \alpha_1 &= i_{X_1} \beta_1 \\ \alpha_2 &= i_{X_2} \beta_1 - \frac{1}{2!} (i_{X_1} \beta_1) \cdot \text{ad}_{X_1} + i_{X_1} \beta_2. \end{aligned}$$

Since β_1 is non-degenerate, the first equality has a (unique) solution

$$X_1 = -(\alpha_1 \otimes 1)r_1 = -\chi_{r_1}(\alpha_1),$$

where $r_1 = \mu_{r_1}^{-1}(\beta_1)$ and χ_{r_1} is the map defined before.

From the second equality, X_1 is known and β_1 being non-degenerate, we obtain

$$X_2 = -\chi_{r_1} \left(\alpha_2 + \frac{1}{2!} (i_{X_1} \beta_1) \cdot \text{ad}_{X_1} - i_{X_1} \beta_2 \right).$$

Suppose now that we know X_1, X_2, \dots, X_{p-1} . Putting α_p equal to the expression given in theorem 6.3 with $k = p$ and separating some terms of the

sum, we obtain

$$\begin{aligned} \alpha_p = & i_{X_p} \beta_1 + \sum_{i=2}^p \frac{1}{i!} \sum_{\substack{a_1+\dots+a_i=p \\ a_1, \dots, a_i \geq 1}} ((-1)^{i+1} (i_{X_{a_1}} \beta_1) \cdot ad_{X_{a_2}} \cdots ad_{X_{a_i}}) + \\ & + \sum_{j=2}^p \left(\sum_{i=1}^{p-j+1} \frac{1}{i!} \sum_{\substack{a_1+\dots+a_i=p-j+1 \\ a_1, \dots, a_i \geq 1}} ((-1)^{i+1} (i_{X_{a_1}} \beta_j) \cdot ad_{X_{a_2}} \cdots ad_{X_{a_i}}) \right). \end{aligned} \quad (6.12)$$

Consider the 1-cochain

$$\begin{aligned} \omega_p = & \alpha_p - \sum_{i=2}^p \frac{1}{i!} \sum_{\substack{a_1+\dots+a_i=p \\ a_1, \dots, a_i \geq 1}} ((-1)^{i+1} (i_{X_{a_1}} \beta_1) \cdot ad_{X_{a_2}} \cdots ad_{X_{a_i}}) \\ & - \sum_{j=2}^p \left(\sum_{i=1}^{p-j+1} \frac{1}{i!} \sum_{\substack{a_1+\dots+a_i=p-j+1 \\ a_1, \dots, a_i \geq 1}} ((-1)^{i+1} (i_{X_{a_1}} \beta_j) \cdot ad_{X_{a_2}} \cdots ad_{X_{a_i}}) \right), \quad p \geq 2. \end{aligned}$$

We may say that, knowing X_1, X_2, \dots, X_{p-1} , ω_p is determined.

Again, as β_1 is non-degenerate, we can compute X_p , using ω_p and the equality (6.12):

$$X_p = -\chi_{r_1}(\omega_p) = -(\omega_p \otimes 1)r_1.$$

By the bijectivity between β'_t and X_t is easy to see that if one of them is convergent so is the other. \blacksquare

3) We need to relate the Etingof-Kazhdan quantization of classical doubles with the isomorphisms between these doubles. Even if the following proposition could be expected its proof is not trivial.

Proposition 6.5. [22] *Let $(\mathfrak{a}, [,]_{\mathfrak{a}}, \varepsilon_{\mathfrak{a}} = d_{c r_s})$ and $(\mathfrak{a}, [,]_{\mathfrak{a}}, \varepsilon'_{\mathfrak{a}} = d_{c r'_s})$ be non-degenerate triangular Lie bialgebras, $s \in \Gamma$, whose quasitriangular double Lie bialgebras are respectively $(\mathfrak{a} \oplus \mathfrak{a}_{r_s}^*, [,]_{\mathfrak{a} \oplus \mathfrak{a}_{r_s}^*}, \varepsilon_{\mathfrak{a} \oplus \mathfrak{a}_{r_s}^*} = d_{c r})$ and $(\mathfrak{a} \oplus \mathfrak{a}_{r_s}^*, [,]_{\mathfrak{a} \oplus \mathfrak{a}_{r_s}^*}, \varepsilon_{\mathfrak{a} \oplus \mathfrak{a}_{r_s}^*} = d_{c r})$. Let $(\varphi_s^1; \psi_s) : \mathfrak{a} \oplus \mathfrak{a}_{r_s}^* \longrightarrow \mathfrak{a} \oplus \mathfrak{a}_{r'_s}^*$ be a Lie algebra isomorphism such that $\varphi_s^1 : \mathfrak{a} \longrightarrow \mathfrak{a}$ and $\psi_s : \mathfrak{a}_{r_s}^* \longrightarrow \mathfrak{a}_{r'_s}^*$ are Lie algebra isomorphisms. Let $\tilde{\varphi}_s^1, \tilde{\psi}_s$ be the extensions of φ_s^1 and ψ_s to homomorphisms $\mathcal{U}\mathfrak{a}[[\hbar]] \longrightarrow \mathcal{U}\mathfrak{a}[[\hbar]]$ and $\mathcal{U}\mathfrak{a}_{r_s}^*[[\hbar]] \longrightarrow \mathcal{U}\mathfrak{a}_{r'_s}^*[[\hbar]]$ respectively. Let $X \in \mathcal{U}(\mathfrak{a} \oplus \mathfrak{a}_{r_s}^*)^{\otimes 2}$. Let ϕ_{r_s} be the $\mathfrak{a} \oplus \mathfrak{a}_{r_s}^*$ -module isomorphism from $\mathcal{U}(\mathfrak{a} \oplus \mathfrak{a}_{r_s}^*)$ to $M_+^{r_s} \otimes M_-^{r_s}$ such that*

$\phi_{r_s} \left(\mathbf{1}_{U(\mathfrak{a} \oplus \mathfrak{a}_{r_s}^*)} \right) = \mathbf{1}_+^{r_s} \otimes \mathbf{1}_-^{r_s}$ and similarly for $\phi_{r'_s}$. (See the definition of ϕ in theorem 3.2). Then we have

$$\phi_{r'_s}^{-1} \left[\left((\tilde{\varphi}_s^1; \tilde{\psi}_s)^{\otimes 2} (X) \right) \cdot (\mathbf{1}_+^{r'_s} \otimes \mathbf{1}_-^{r'_s}) \right] = \left((\tilde{\varphi}_s^1; \tilde{\psi}_s) \circ \phi_{r_s}^{-1} \right) (X \cdot (\mathbf{1}_+^{r_s} \otimes \mathbf{1}_-^{r_s})).$$

If in the expression of J given in theorem 3.2 we take into account the above proposition and also that a Lie associator determines $\Phi = e^{P(\hbar\Omega_{12}, \hbar\Omega_{23})}$ we arrive [22] to:

Proposition 6.6. *Hypotheses are as in the above proposition and suppose moreover that $(\varphi_s^1; \psi_s) \otimes (\varphi_s^1; \psi_s) \Omega = \Omega$. Denote by $J_{r'_s}$ and J_{r_s} the corresponding elements in theorem 3.2. Then we have the equality $J_{r'_s} = (\tilde{\varphi}_s^1; \tilde{\psi}_s)^{\otimes 2} J_{r_s}$. In particular this proposition is valid for the Lie bialgebra isomorphism $(\varphi_s^1; \varphi_s^2)$ constructed in proposition 6.1 and those in propositions 6.3 and 6.4.*

We can also prove the following:

Proposition 6.7. *Let $(\mathfrak{a}, [,]_{\mathfrak{a}}, \varepsilon_{\mathfrak{a}} = d_c r_s)$ and $(\mathfrak{a}, [,]_{\mathfrak{a}}, \varepsilon'_{\mathfrak{a}} = d_c r'_s)$ be non-degenerate triangular Lie bialgebras, $s \in \Gamma$. Let $\varphi_s^1 : \mathfrak{a} \rightarrow \mathfrak{a}$ be a Lie algebra isomorphism such that $r'_s = (\varphi_s^1 \otimes \varphi_s^1) r_s$ and let $(\varphi_s^1; \varphi_s^2)$ be the Lie bialgebra isomorphism between the corresponding classical doubles constructed in proposition 6.1. Then we have*

$$\tilde{\pi}'_s \circ (\varphi_s^1; \varphi_s^2) = \varphi_s^1 \circ \tilde{\pi}_s,$$

where $\tilde{\pi}$ is defined in Proposition 3.4.

4) Using propositions 6.1, 6.6, 6.7 and corollary 6.2 we can prove:

Theorem 6.8. *Let $\tilde{J}_{r_{\hbar}}$ and $\tilde{J}_{r'_{\hbar}}$ be invariant star products on a non-degenerate triangular Lie bialgebra over \mathbb{R} , $(\mathfrak{a}, [,]_{\mathfrak{a}}, \varepsilon_{\mathfrak{a}} = d_c r_1)$, and determined as in lemma 4.9 and propositions 4.10, respectively by non-degenerate skew-symmetric solutions $r_{\hbar} = r_1 + \dots$ and $r'_{\hbar} = r_1 + \dots$ of YBE on Lie algebra over $\mathbb{R}[[\hbar]]$, $(\mathfrak{a}[[\hbar]], [,]_{\mathfrak{a}[[\hbar]]})$. Let $\mu_{r_{\hbar}}(r_{\hbar}) = \beta_{\hbar} = \beta_1 + \beta_2 \hbar + \dots \in (\mathfrak{a}^* \wedge \mathfrak{a}^*)[[\hbar]]$ and $\mu_{r'_{\hbar}}(r'_{\hbar}) = \beta'_{\hbar} = \beta_1 + \beta'_2 \hbar + \dots \in (\mathfrak{a}^* \wedge \mathfrak{a}^*)[[\hbar]]$. Suppose that the cocycles $\beta_{\hbar} = \mu_{r_{\hbar}}(r_{\hbar}) = \beta_1 + \beta_2 \hbar + \dots$ and $\beta'_{\hbar} = \mu_{r'_{\hbar}}(r'_{\hbar}) = \beta_1 + \beta'_2 \hbar + \dots$ belong to the same cohomological class, i.e., that $\beta'_{\hbar} = \beta_{\hbar} + d_c \alpha_{\hbar}$ for some 1-cochain $\alpha_{\hbar} = \alpha_1 \hbar + \alpha_2 \hbar^2 + \dots \in \mathfrak{a}^*[[\hbar]]$. Then, $\tilde{J}_{r_{\hbar}}$ and $\tilde{J}_{r'_{\hbar}}$ are equivalent ISP.*

Proof:

The elements \tilde{J}_{r_t} and $\tilde{J}_{r'_t}$ are defined by

$$\tilde{J}_{r_t} = (\tilde{\pi}_t \otimes \tilde{\pi}_t) J_{r_t}, \quad \tilde{J}_{r'_t} = (\tilde{\pi}'_t \otimes \tilde{\pi}'_t) J_{r'_t},$$

where J_{r_t} and $J_{r'_t}$ are the elements in $\mathcal{U}(\mathfrak{a} \oplus \mathfrak{a}_{r_t}^*)^{\otimes 2}[[\hbar]]$ and $\mathcal{U}(\mathfrak{a} \oplus \mathfrak{a}_{r'_t}^*)^{\otimes 2}[[\hbar]]$, respectively, defined in theorem 3.2.

Since $\beta_t, \beta'_t \in \wedge^2(\mathfrak{a}^*)$ belong to the same cohomological class, by theorem 6.4, there exists a $X_t = X_1 + X_2 t + \dots \in \mathfrak{a}$ such that

$$\exp(\text{ad}_{tX_t}^*)^{\otimes 2}(\beta_t) \equiv (\varphi_t^2)^{\otimes 2} \beta_t = \beta'_t.$$

By proposition 6.2, $\varphi_t^1 = ((\exp(\text{ad}_{tX_t}^*))^{-1})^t = \exp(\text{ad}_{tX_t})$ is a Lie algebra isomorphism $\mathfrak{a} \longrightarrow \mathfrak{a}$ such that $(\varphi_t^1 \otimes \varphi_t^1)r_t = r'_t$, where $r_t = \mu_{r_t}^{-1}(\beta_t)$ and $r'_t = \mu_{r'_t}^{-1}(\beta'_t)$. By proposition 6.1, $(\varphi_t^1; \varphi_t^2)$ is a Lie bialgebra isomorphism between $\mathfrak{a} \oplus \mathfrak{a}_{r_t}^*$ and $\mathfrak{a} \oplus \mathfrak{a}_{r'_t}^*$ such that $(\varphi_t^1; \varphi_t^2)^{\otimes 2}r = r$, where r is the canonical element in $(\mathfrak{a} \oplus \mathfrak{a}^*) \otimes (\mathfrak{a} \oplus \mathfrak{a}^*)$. So, $(\varphi_t^1; \varphi_t^2)^{\otimes 2}\Omega = \Omega$.

Then, we have

$$\tilde{J}_{r'_t} = (\tilde{\pi}'_t \otimes \tilde{\pi}'_t)J_{r'_t} = (\tilde{\varphi}_t^1 \otimes \tilde{\varphi}_t^1)\tilde{J}_{r_t},$$

using propositions 6.6 and 6.7.

Putting $t = \hbar$, we obtain $\tilde{J}_{r'_\hbar} = (\tilde{\varphi}_\hbar^1 \otimes \tilde{\varphi}_\hbar^1)\tilde{J}_{r_\hbar}$, or, equivalently, $\tilde{J}_{r'_\hbar}^{-1} = (\tilde{\varphi}_\hbar^1 \otimes \tilde{\varphi}_\hbar^1)\tilde{J}_{r_\hbar}^{-1}$. The map $\varphi_\hbar^1 = \exp(\text{ad}_{\hbar X_\hbar}) : \mathfrak{a}[[\hbar]] \longrightarrow \mathfrak{a}[[\hbar]]$ is obviously a morphism (in fact, an isomorphism) of Lie algebras over $\mathbb{R}[[\hbar]]$ (where the Lie algebra structure on $\mathfrak{a}[[\hbar]]$ is the trivial one of $(\mathfrak{a}, \mathbb{R})$ by extension of the ring of the scalars from \mathbb{R} to $\mathbb{R}[[\hbar]]$) and, considering $u = 1$ and the last equality, we may apply proposition 3.9 in [6].

Then, the extension $\tilde{\varphi}_\hbar^1 : \mathcal{U}(\mathfrak{a}[[\hbar]]) \longrightarrow \mathcal{U}(\mathfrak{a}[[\hbar]])$ is a morphism of triangular Hopf QUE algebras $A_{\mathfrak{a}[[\hbar]], \tilde{J}_{r_\hbar}^{-1}} \longrightarrow A_{\mathfrak{a}[[\hbar]], \tilde{J}_{r'_\hbar}^{-1}}$. The pairs $(\varphi_\hbar^1, 1)$ and (λ_1, u_1) determine the same morphism if, and only if, $\lambda_1 = \exp(-\hbar \text{ad}_v) \circ \varphi_\hbar^1$ and $u_1 = e^{\hbar v}$, for some $v \in \mathfrak{a}[[\hbar]]$. Putting $v = X_t|_{t=\hbar}$, we obtain $\lambda_1 = 1$, $u_1 = e^{\hbar X_\hbar}$ and

$$\tilde{J}_{r'_\hbar}^{-1} = (u_1 \otimes u_1) \cdot_\hbar \tilde{J}_{r_\hbar}^{-1} \cdot_\hbar \Delta_{\mathfrak{a}}(u_1)^{-1}.$$

This last equality is equivalent to

$$\tilde{J}_{r'_\hbar} = \Delta_{\mathfrak{a}}(u_1) \cdot_\hbar \tilde{J}_{r_\hbar} \cdot_\hbar (u_1^{-1} \otimes u_1^{-1}),$$

and u_1^{-1} defines an equivalence between the invariant star products \tilde{J}_{r_\hbar} and $\tilde{J}_{r'_\hbar}$ on $(\mathfrak{a}, [\cdot, \cdot]_{\mathfrak{a}}, \varepsilon_{\mathfrak{a}} = d_c r_1)$. ■

Before proving the converse result we need the following lemma.

Lemma 6.9. *Suppose that in theorem 6.8*

$$\beta_{\hbar} = \mu_{r_{\hbar}}(r_{\hbar}) = \beta_1 + \beta_2\hbar + \cdots + \beta_{R-1}t^{R-2} + \beta_R\hbar^{R-1} + \dots$$

$$\beta'_{\hbar} = \mu_{r'_{\hbar}}(r'_{\hbar}) = \beta_1 + \beta_2\hbar + \cdots + \beta_{R-1}\hbar^{R-2} + (\beta_R + d_c\alpha_{R-1})\hbar^{R-1} + \dots,$$

where α_{R-1} is a 1-cochain. This means that β_{\hbar} and β'_{\hbar} are equal except in the term of order $R-1$. Then, $\tilde{J}_{r_{\hbar}}$ and $\tilde{J}_{r'_{\hbar}}$ are equivalent,

$$\tilde{J}_{r'_{\hbar}} = \Delta_{\mathfrak{a}}(E)^{-1} \cdot_{\hbar} \tilde{J}_{r_{\hbar}} \cdot_{\hbar} (E \otimes E),$$

and the element $E = 1 + E_1\hbar + E_2\hbar^2 + \cdots + E_{R-1}\hbar^{R-1} + \dots$ which defines this equivalence verifies

$$E_1 = 0, \quad E_2 = 0, \quad \dots, \quad E_{R-2} = 0, \quad E_{R-1} = \chi_{r_1}(\alpha_{R-1}) = \mu_{r_1}^{-1}(\alpha_{R-1}).$$

Proof:

Consider the elements β_{\hbar} and β'_{\hbar} obeying the above conditions. One of the steps of the proof of theorem 6.8 is to find the element $X_t = X_1 + X_2t + X_3t^2 + \cdots + X_{R-1}t^{R-2} + \dots$ of theorem 6.4. For the elements β_t and β'_t considered, this element will be

$$X_1 = 0, \quad X_2 = 0, \quad \dots, \quad X_{R-2} = 0, \quad X_{R-1} = -\chi_{r_1}(\alpha_{R-1}).$$

Then, the element E will be, in view of the same proof, the following:

$$E = u_1^{-1} = (e^{\hbar X_{\hbar}})^{-1} = 1 + 0\hbar + \cdots + 0\hbar^{R-2} + \chi_{r_1}(\alpha_{R-1})\hbar^{R-1} + \dots$$

■

Lemma 6.9 and Hochschild cohomology properties allow us to prove

Theorem 6.10. *Let $\tilde{J}_{r_{\hbar}}$ and $\tilde{J}_{r'_{\hbar}}$ as in a) Theorem 6.8. Suppose $\tilde{J}_{r_{\hbar}}$ and $\tilde{J}_{r'_{\hbar}}$ are equivalent. Then, β_{\hbar} and β'_{\hbar} belong to the same cohomological class, i.e., there exists a formal 1-cochain $\alpha_{\hbar} = \alpha_1\hbar + \alpha_2\hbar^2 + \cdots \in \mathfrak{a}^*[[\hbar]]$ such that $\beta'_{\hbar} = \beta_{\hbar} + d_c\alpha_{\hbar}$.*

Proof:

If $\tilde{J}_{r_{\hbar}}$ and $\tilde{J}_{r'_{\hbar}}$ are equivalent ISP, there exists $E^{(1)} = 1 + E_1^{(1)}\hbar + E_2^{(1)}\hbar^2 + \cdots \in \mathcal{U}\mathfrak{a}[[\hbar]]$ such that

$$\tilde{J}_{r'_{\hbar}} = \Delta_{\mathfrak{a}}(E^{(1)})^{-1} \cdot_{\hbar} \tilde{J}_{r_{\hbar}} \cdot_{\hbar} (E^{(1)} \otimes E^{(1)}).$$

At order 1, this equivalence may be written as $(\tilde{J}_{r'_{\hbar}})_1 - (\tilde{J}_{r_{\hbar}})_1 = d_H E_1^{(1)}$, where d_H is the Hochschild cohomology operator and $E_1^{(1)} \in \mathcal{U}\mathfrak{a}$. But, in this case,

$(\tilde{J}'_{r'_h})_1 = (\tilde{J}_{r_h})_1 = \frac{1}{2}r_1$. So, we have $d_H E_1^{(1)} = 0$, which implies that $E_1^{(1)} \in \mathfrak{a}$. Therefore, $\Delta_{\mathfrak{a}}(E_1^{(1)}) = E_1^{(1)} \otimes 1 + 1 \otimes E_1^{(1)}$.

At order 2, there exists $E_2^{(1)} \in \mathcal{U}\mathfrak{a}$ such that

$$(\tilde{J}'_{r'_h})_2 - (\tilde{J}_{r_h})_2 + G_2(E_1^{(1)}, (\tilde{J}'_{r'_h})_1, (\tilde{J}_{r_h})_1) = d_H E_2^{(1)}, \quad (6.13)$$

where, for $k = 2, 3, \dots$,

$$\begin{aligned} G_k(E_1, \dots, E_{k-1}, F'_1, \dots, F'_{k-1}, F_1, \dots, F_{k-1}) &= \\ &= \sum_{\substack{i+j=k \\ i,j \geq 1}} (\Delta_{\mathfrak{a}}(E_i) \cdot F'_j - F_i \cdot (1 \otimes E_j) - F_i \cdot (E_j \otimes 1)) - \sum_{\substack{i+j=k \\ i,j \geq 1}} (E_i \otimes 1) \cdot (1 \otimes E_j) - \\ &\quad - \sum_{\substack{i+j+l=k \\ i,j,l \geq 1}} F_i \cdot (E_j \otimes 1) \cdot (1 \otimes E_l). \end{aligned}$$

Since $(\tilde{J}'_{r'_h})_1 = (\tilde{J}_{r_h})_1 = \frac{1}{2}r_1$, we have

$$\begin{aligned} G_2(E_1^{(1)}, (\tilde{J}'_{r'_h})_1, (\tilde{J}_{r_h})_1) &= \\ &= \Delta_{\mathfrak{a}}(E_1^{(1)}) \cdot \frac{1}{2}r_1 - \frac{1}{2}r_1 \cdot (1 \otimes E_1^{(1)} + E_1^{(1)} \otimes 1) - E_1^{(1)} \otimes E_1^{(1)} \\ &= -\frac{1}{2}d_P^{r_1} E_1^{(1)} - E_1^{(1)} \otimes E_1^{(1)}. \end{aligned}$$

The skew-symmetric part of $G_2(E_1^{(1)}, (\tilde{J}'_{r'_h})_1, (\tilde{J}_{r_h})_1)$ is $-\frac{1}{2}d_P^{r_1} E_1^{(1)}$, where $d_P^{r_1}$ is the Poisson cohomology operator. We know that $(\tilde{J}'_{r'_h})_2 - (\tilde{J}_{r_h})_2 = \frac{1}{2}(r'_2 - r_2)$. So, the equality between the skew-symmetric parts of both sides of (6.13) leads to

$$r'_2 = r_2 + d_P^{r_1} E_1^{(1)}. \quad (6.14)$$

Let $\beta_t = \mu_{r_t}(r_t) = \beta_1 + \beta_2 t + \dots$ and $\beta'_t = \beta_t^{(1)} = \mu_{r'_t}(r'_t) = \beta_1 + \beta'_2 t + \beta'_3 t^2 + \dots$. Using (6.14), we get $\beta'_2 - \beta_2 = -\mu_{r_1}(r'_2 - r_2) = -\mu_{r_1}(d_P^{r_1} E_1^{(1)}) = d_c(\mu_{r_1}(E_1^{(1)}))$ (the last equality, using relation $\mu \circ (-d_P) = d_c \circ \mu$). This means that there exists a 1-cochain $\alpha_1 = \mu_{r_1}(E_1^{(1)})$ such that $\beta'_2 = \beta_2 + d_c \alpha_1$.

Consider now the following elements:

$$\beta_t^{(2)} = \beta_1 + \beta_2 t + \beta'_3 t^2 + \beta'_4 t^3 + \dots \quad \text{and} \quad r_t^{(2)} = \mu_{r_t^{(2)}}^{-1}(\beta_t^{(2)}).$$

The elements $\beta_t^{(2)}$ and β'_t belong to the same formal cohomological class. By theorem 6.8, the star products $\tilde{J}_{r_h^{(2)}}$ and $\tilde{J}'_{r'_h}$ are equivalent, and the element

$u^{(2)}$ which defines the equivalence

$$\tilde{J}_{r'_h} = \Delta_{\mathfrak{a}}(u^{(2)})^{-1} \cdot \tilde{J}_{r_h^{(2)}} \cdot (u^{(2)} \otimes u^{(2)}),$$

equivalent also to

$$\tilde{J}_{r_h^{(2)}} = \Delta_{\mathfrak{a}}(u^{(2)}) \cdot \tilde{J}_{r'_h} \cdot ((u^{(2)})^{-1} \otimes (u^{(2)})^{-1})$$

satisfies (by the previous lemma)

$$u^{(2)} = 1 + \mu_{r_1}^{-1}(\alpha_1)\hbar + \dots = 1 + E_1^{(1)}\hbar + \dots .$$

But, as \tilde{J}_{r_h} and $\tilde{J}_{r'_h}$ are also equivalent, we obtain

$$\begin{aligned} \tilde{J}_{r_h^{(2)}} &= \Delta_{\mathfrak{a}}(u^{(2)}) \cdot \Delta_{\mathfrak{a}}(E^{(1)})^{-1} \cdot \tilde{J}_{r_h} \cdot (E^{(1)} \otimes E^{(1)}) \cdot ((u^{(2)})^{-1} \otimes (u^{(2)})^{-1}) \\ &= \Delta_{\mathfrak{a}}(E^{(1)} \cdot (u^{(2)})^{-1})^{-1} \cdot \tilde{J}_{r_h} \cdot ((E^{(1)} \cdot (u^{(2)})^{-1}) \otimes (E^{(1)} \cdot (u^{(2)})^{-1})), \end{aligned}$$

i.e., the element $E^{(2)} = E^{(1)} \cdot (u^{(2)})^{-1} \in \mathcal{U}\mathfrak{a}[[\hbar]]$ defines an equivalence between \tilde{J}_{r_h} and $\tilde{J}_{r_h^{(2)}}$ at any order, and we may compute the first terms of $E^{(2)}$:

$$E^{(2)} = 1 + (E_1^{(1)} - E_1^{(1)})\hbar + E_2^{(2)}\hbar^2 + \dots = 1 + 0\hbar + E_2^{(2)}\hbar^2 + \dots .$$

Since $r_t^{(2)} = r_1 + r_2t + r_3^{(2)}t^2 + \dots$ and $r_t = r_1 + r_2t + r_3t^2 + \dots$, we have

$$\left(\tilde{J}_{r_h^{(2)}}\right)_1 = \left(\tilde{J}_{r_h}\right)_1, \quad \left(\tilde{J}_{r_h^{(2)}}\right)_2 = \left(\tilde{J}_{r_h}\right)_2 \quad \text{and} \quad \left(\tilde{J}_{r_h^{(2)}}\right)_3 - \left(\tilde{J}_{r_h}\right)_3 = \frac{1}{2}(r_3^{(2)} - r_3).$$

But \tilde{J}_{r_h} and $\tilde{J}_{r_h^{(2)}}$ are equivalent. So, they are equivalent at order 2. This means that there exists an element $E_2^{(2)} \in \mathcal{U}\mathfrak{a}$ such that

$$\left(\tilde{J}_{r_h^{(2)}}\right)_2 - \left(\tilde{J}_{r_h}\right)_2 + G_2\left(E_1^{(2)}, \left(\tilde{J}_{r_h^{(2)}}\right)_1, \left(\tilde{J}_{r_h}\right)_1\right) = d_H E_2^{(2)}.$$

Since $E_1^{(2)} = 0$, we have $G_2\left(E_1^{(2)}, \left(\tilde{J}_{r_h^{(2)}}\right)_1, \left(\tilde{J}_{r_h}\right)_1\right) = 0$ and so $d_H E_2^{(2)} = 0$, which implies that $E_2^{(2)}$ belongs to \mathfrak{a} .

The equivalence at order 3 means that there exists $E_3^{(2)} \in \mathcal{U}\mathfrak{a}$ such that

$$\left(\tilde{J}_{r_h^{(2)}}\right)_3 - \left(\tilde{J}_{r_h}\right)_3 + G_3\left(E_1^{(2)}, E_2^{(2)}, \left(\tilde{J}_{r_h^{(2)}}\right)_1, \left(\tilde{J}_{r_h}\right)_1, \left(\tilde{J}_{r_h^{(2)}}\right)_2, \left(\tilde{J}_{r_h}\right)_2\right) = d_H E_3^{(2)}. \quad (6.15)$$

Since $E_1^{(2)} = 0$ and $\left(\tilde{J}_{r_{\hbar}^{(2)}}\right)_1 = \left(\tilde{J}_{r_{\hbar}}\right)_1$, we obtain

$$\begin{aligned} G_3 \left(E_1^{(2)}, E_2^{(2)}, \left(\tilde{J}_{r_{\hbar}^{(2)}}\right)_1, \left(\tilde{J}_{r_{\hbar}}\right)_1, \left(\tilde{J}_{r_{\hbar}^{(2)}}\right)_2, \left(\tilde{J}_{r_{\hbar}}\right)_2 \right) &= \\ &= \Delta_{\mathfrak{a}}(E_2^{(2)}) \cdot \left(\tilde{J}_{r_{\hbar}}\right)_1 - \left(\tilde{J}_{r_{\hbar}}\right)_1 \cdot (1 \otimes E_2^{(2)}) - \left(\tilde{J}_{r_{\hbar}}\right)_1 \cdot (E_2^{(2)} \otimes 1). \end{aligned}$$

The skew-symmetric part of this element is

$$AG_3 \left(E_1^{(2)}, E_2^{(2)}, \left(\tilde{J}_{r_{\hbar}^{(2)}}\right)_1, \left(\tilde{J}_{r_{\hbar}}\right)_1, \left(\tilde{J}_{r_{\hbar}^{(2)}}\right)_2, \left(\tilde{J}_{r_{\hbar}}\right)_2 \right) = -\frac{1}{2}d_P^{r_1}E_2^{(2)}.$$

The equality of the skew-symmetric parts of both sides of (6.15) leads to

$$r_3^{(2)} = r_3 + d_P^{r_1}E_2^{(2)}.$$

Thus,

$$\beta'_3 - \beta_3 = \beta_3^{(2)} - \beta_3 = -\mu_{r_1}(r_3^{(2)} - r_3) = -\mu_{r_1}(d_P^{r_1}E_2^{(2)}) = d_c(\mu_{r_1}(E_2^{(2)})).$$

This means that there exists a 1-cochain $\alpha_2 = \mu_{r_1}(E_2^{(2)})$ such that

$$\beta'_3 = \beta_3 + d_c\alpha_2.$$

Using the induction hypothesis, and following the same steps, we conclude the proof. \blacksquare

Combining the last two theorems we obtain the following result, similar in Etingof-Kazhdan quantization theory to the one by Drinfeld in [5].

Theorem 6.11. *Let $\tilde{J}_{r_{\hbar}}$ and $\tilde{J}_{r'_{\hbar}}$ be as in a) Theorem 6.8. Then, $\tilde{J}_{r_{\hbar}}$ and $\tilde{J}_{r'_{\hbar}}$ are equivalent ISP if, and only if, β_{\hbar} and β'_{\hbar} belong to the same formal cohomological class. In other words, $\tilde{J}_{r_{\hbar}}$ and $\tilde{J}_{r'_{\hbar}}$ are equivalent ISP if, and only if, there exists a formal 1-cochain $\alpha_{\hbar} = \alpha_1\hbar + \alpha_2\hbar^2 + \dots$ such that $\beta'_{\hbar} = \beta_{\hbar} + d_c\alpha_{\hbar}$.*

Theorem 6.11 and Remark 2) in page 841 of [7] allow us to obtain

Theorem 6.12. *Two triangular Hopf QUE-algebras $A_{\mathfrak{a}[[\hbar]], \tilde{J}_{r_{\hbar}}^{-1}}$ and $A_{\mathfrak{a}[[\hbar]], \tilde{J}_{r'_{\hbar}}^{-1}}$ are isomorphic if, and only if, there exists an isomorphism of Lie algebras $\lambda : \mathfrak{a}[[\hbar]] \longrightarrow \mathfrak{a}[[\hbar]]$ over $MR[[\hbar]]$ such that $(\lambda^2 \otimes \lambda^2)\beta_{\hbar}$ and β'_{\hbar} belong to the same cohomological class where $\beta_{\hbar} = \mu_{r_{\hbar}}(r_{\hbar})$, $\beta'_{\hbar} = \mu_{r'_{\hbar}}(r'_{\hbar})$ and $\lambda^2 = (\lambda^{-1})^{\dagger}$.*

Proof:

If the pair (λ, u) defines an isomorphism between the triangular Hopf QUE algebras (see proposition 3.9 in [6]), in particular, the map $Ad(u) \circ \tilde{\lambda}$ satisfies the following equality

$$(Ad(u) \circ \tilde{\lambda}) \otimes (Ad(u) \circ \tilde{\lambda}) R^{r\hbar} = R^{r'\hbar}.$$

We know that $R^{r\hbar} = 1 + r_1\hbar + O(\hbar^2)$ and $R^{r'\hbar} = 1 + r_1\hbar + O(\hbar^2)$. Computing the coefficients of \hbar at both sides of last equality, we obtain

$$(\lambda \otimes \lambda)r_1 = r_1 \text{ mod } \hbar.$$

So, $(\lambda \otimes \lambda)(r_{\hbar}) = r_1 + r_2''\hbar + r_3''\hbar^2 + \dots$. Denote this element by r_{\hbar}'' (it is also a solution of CYBE). We know also that

$$\tilde{J}_{r_{\hbar}'}^{-1} = (u \otimes u) \cdot (\tilde{\lambda} \otimes \tilde{\lambda}) \left(\tilde{J}_{r_{\hbar}}^{-1} \right) \cdot \Delta_{\mathbf{a}}(u)^{-1}. \quad (6.16)$$

Consider the invariant star product $\tilde{J}_{r_{\hbar}''}$ defined through the E-K quantization. By proposition 6.6, since $(\lambda \otimes \lambda)(r_{\hbar}) = r_{\hbar}''$, we have

$$(\lambda, \lambda^2)^{\otimes 2} J_{r_{\hbar}} = J_{r_{\hbar}''},$$

where $\lambda^2 = (\lambda^{-1})^t$. Using proposition 6.7, we obtain

$$(\tilde{\lambda} \otimes \tilde{\lambda}) \tilde{J}_{r_{\hbar}} = \tilde{J}_{r_{\hbar}''}.$$

Taking inverses at both sides of (6.16), we obtain

$$\begin{aligned} \tilde{J}_{r_{\hbar}'} &= \Delta_{\mathbf{a}}(u) \cdot (\tilde{\lambda} \otimes \tilde{\lambda}) \left(\tilde{J}_{r_{\hbar}} \right) \cdot (u^{-1} \otimes u^{-1}) \\ &= \Delta_{\mathbf{a}}(u) \cdot \tilde{J}_{r_{\hbar}''} \cdot (u^{-1} \otimes u^{-1}). \end{aligned}$$

This means that $\tilde{J}_{r_{\hbar}'}$ and $\tilde{J}_{r_{\hbar}''}$ are equivalent star products. By theorem 6.10, $\beta_{\hbar}' = \mu_{r_{\hbar}'}(r_{\hbar}')$ and $\beta_{\hbar}'' = \mu_{r_{\hbar}''}(r_{\hbar}'')$ belong to the same formal cohomological class. Since $(\lambda \otimes \lambda)r_{\hbar} = r_{\hbar}''$, by proposition 6.2, we get $\beta_{\hbar}'' = (\lambda^{-1})^t \otimes (\lambda^{-1})^t \beta_{\hbar} = (\lambda^2 \otimes \lambda^2)\beta_{\hbar}$ and we conclude that β_{\hbar}' and $(\lambda^2 \otimes \lambda^2)\beta_{\hbar}$ belong to the same formal cohomological class.

If $((\varphi^1)^{-1})^t \otimes ((\varphi^1)^{-1})^t \beta_{\hbar} = \beta_{\hbar}''$ and β_{\hbar}' belong to the same cohomological class, then, by theorem 6.8, $\tilde{J}_{r_{\hbar}''}$ and $\tilde{J}_{r_{\hbar}'}$ are equivalent star products, where $r_{\hbar}'' = \mu_{r_{\hbar}''}^{-1}(\beta_{\hbar}'')$. This means that there exists an element $u \in \mathcal{U}\mathbf{a}[[\hbar]]$, $u \equiv 1 \text{ mod } \hbar$, such that

$$\tilde{J}_{r_{\hbar}'} = \Delta_{\mathbf{a}}(u)^{-1} \cdot \tilde{J}_{r_{\hbar}''} \cdot (u \otimes u),$$

equivalent also to

$$\tilde{J}_{r'_h}^{-1} = (u^{-1} \otimes u^{-1}) \cdot \tilde{J}_{r''_h}^{-1} \cdot \Delta_{\mathfrak{a}}(u). \quad (6.17)$$

Since $(\varphi^1 \otimes \varphi^1)r_h = r''_h$ (see proposition 6.2), we have, by propositions 6.6 and 6.7,

$$\tilde{J}_{r''_h} = (\tilde{\varphi}^1 \otimes \tilde{\varphi}^1)\tilde{J}_{r_h}.$$

Substituting this in (6.17), we obtain

$$\tilde{J}_{r'_h}^{-1} = (u^{-1} \otimes u^{-1}) \cdot (\tilde{\varphi}^1 \otimes \tilde{\varphi}^1) \left(\tilde{J}_{r_h}^{-1} \right) \cdot \Delta_{\mathfrak{a}}(u).$$

Considering $\lambda_1 = \varphi^1$ and $u_1 = u^{-1}$, the pair (λ_1, u_1) defines an isomorphism between $A_{\mathfrak{a}[[\hbar]], \tilde{J}_{r'_h}^{-1}}$ and $A_{\mathfrak{a}[[\hbar]], \tilde{J}_{r_h}^{-1}}$ (see proposition 3.9 in [6]). \blacksquare

5) From the above results and Remark 2) in page 841 of [7] we may also prove :

Proposition 6.13. *Let $A_{\mathfrak{a} \oplus \mathfrak{a}_{r_h}^*, \Omega, J_{r_h}^{-1}}$ and $A_{\mathfrak{a} \oplus \mathfrak{a}_{r'_h}^*, \Omega, J_{r'_h}^{-1}}$ be quasitriangular Hopf QUE algebras over $\mathbb{R}[[\hbar]]$ which are quantizations, as in theorem 4.7, of the quasitriangular Lie bialgebra $(\mathfrak{a} \oplus \mathfrak{a}_{r_1}^*, [,]_{\mathfrak{a} \oplus \mathfrak{a}_{r_1}^*}, \varepsilon_{\mathfrak{a} \oplus \mathfrak{a}_{r_1}^*} = d_c r)$. Let $\beta_h = \mu_{r_h}(r_h) = \beta_1 + \beta_2 \hbar + \dots$ and $\beta'_h = \mu_{r'_h}(r'_h) = \beta_1 + \beta'_2 \hbar + \dots$. If β_h and β'_h belong to the same cohomological class, then $A_{\mathfrak{a} \oplus \mathfrak{a}_{r_h}^*, \Omega, J_{r_h}^{-1}}$ and $A_{\mathfrak{a} \oplus \mathfrak{a}_{r'_h}^*, \Omega, J_{r'_h}^{-1}}$ are isomorphic.*

Proof:

If β_t and β'_t belong to the same cohomological class, by theorem 6.4, there exists a unique element $X_t = X_1 + X_2 t + \dots \in \mathfrak{a}$, such that

$$\exp(ad_{X_t}^*)^{\otimes 2} \beta_t = \beta'_t.$$

Using proposition 6.2 with $\varphi_t^2 = \exp(ad_{X_t}^*)$, there exists an isomorphism $\varphi_t^1 = ((\varphi_t^2)^t)^{-1} : \mathfrak{a} \longrightarrow \mathfrak{a}$ of Lie algebras such that $(\varphi_t^1 \otimes \varphi_t^1)r_t = r'_t$. By proposition 6.1, this pair $(\varphi_t^1; \varphi_t^2)$ defines an isomorphism from the Lie bialgebra $\mathfrak{a} \oplus \mathfrak{a}_{r_t}^*$ to the Lie bialgebra $\mathfrak{a} \oplus \mathfrak{a}_{r'_t}^*$ and sends the canonical element r of the vector space $(\mathfrak{a} \oplus \mathfrak{a}^*)^{\otimes 2}$ to itself. Thus, $(\varphi_t^1; \varphi_t^2)$ also will send Ω into Ω , where $\Omega = r_{12} + r_{21}$. Applying now proposition 6.6, we have

$$J_{r'_t} = (\tilde{\varphi}_t^1; \tilde{\varphi}_t^2)^{\otimes 2} J_{r_t},$$

where $\tilde{\varphi}_t^1$ and $\tilde{\varphi}_t^2$ are extensions of φ_t^1 and φ_t^2 to homomorphisms $\mathcal{U}\mathfrak{a}[[\hbar]] \longrightarrow \mathcal{U}\mathfrak{a}[[\hbar]]$ and $\mathcal{U}\mathfrak{a}_{r_t}^*[[\hbar]] \longrightarrow \mathcal{U}\mathfrak{a}_{r'_t}^*[[\hbar]]$, respectively. Putting $t = \hbar$ it is clear

that $(\varphi_{\hbar}^1, \varphi_{\hbar}^2)$ defines an isomorphism $\mathfrak{a} \oplus \mathfrak{a}_{r_{\hbar}}^* \longrightarrow \mathfrak{a} \oplus \mathfrak{a}_{r'_{\hbar}}^*$ and the elements $J_{r'_{\hbar}} = J_{r'_t} |_{t=\hbar}$, $J_{r_{\hbar}} = J_{r_t} |_{t=\hbar}$. We obtain the equality

$$J_{r'_{\hbar}} = (\tilde{\varphi}_{\hbar}^1; \tilde{\varphi}_{\hbar}^2)^{\otimes 2} J_{r_{\hbar}}.$$

Using an analogous proposition to proposition 3.9 in [6] with $\lambda = (\varphi_{\hbar}^1; \varphi_{\hbar}^2)$ and $u = 1$, we conclude that the map

$$\widetilde{(\varphi_{\hbar}^1; \varphi_{\hbar}^2)} = (\tilde{\varphi}_{\hbar}^1; \tilde{\varphi}_{\hbar}^2) : \hat{\mathcal{U}}(\mathfrak{a} \oplus \mathfrak{a}_{r_{\hbar}}^*) \longrightarrow \hat{\mathcal{U}}(\mathfrak{a} \oplus \mathfrak{a}_{r'_{\hbar}}^*)$$

is an isomorphism $A_{\mathfrak{a} \oplus \mathfrak{a}_{r_{\hbar}}^*, \Omega, J_{r_{\hbar}}^{-1}} \longrightarrow A_{\mathfrak{a} \oplus \mathfrak{a}_{r'_{\hbar}}^*, \Omega, J_{r'_{\hbar}}^{-1}}$. ■

About the converse of this proposition, we have [22] some examples of isomorphisms where $[\beta_{\hbar}] \neq [\beta'_{\hbar}]$.

References

- [1] F. Bayen, M. Flato, C. Fronsdal, A. Lichnerowicz, and D. Sternheimer, *Deformation theory and quantization*, Ann. Phys. **111** (1978), 61 – 151.
- [2] N. Bourbaki, *Groups et algèbres de Lie*, Hermann, Paris, 1960.
- [3] P. Cartier, *Hyperalgèbres et Groups de Lie formels*, Séminaire Sophus Lie, 2ème année, Faculté des Sciences de Paris, 19.
- [4] N. Bourbaki, *Algèbre chapitres 4-7*, Actualités Scientifiques et Industrielles, vol. 1293, Hermann, Paris, 1961.
- [5] V.G. Drinfeld, *On constant, quasiclassical solutions of the Yang-Baxter equations*, Sov. Math. Dokl. **28** (1983), 667 – 671.
- [6] V.G. Drinfeld, *Quasi-Hopf algebras*, Leningrad Math. J. **1** (6) (1990), 1419 – 1457.
- [7] V.G. Drinfeld, *On quasitriangular quasi-Hopf algebras and a group closely connected with $Gal(\overline{\mathbb{Q}}/\mathbb{Q})$* , Leningrad Math. J. **2** (4) (1991), 829 – 860.
- [8] D. Kazhdan and G. Lusztig, *Tensor structures arising from affine Lie algebras, III*, J. Amer. Math. Soc. **7** (1994), 335 – 381.
- [9] P. Etingof and D. Kazhdan, *Quantization of Lie Bialgebras, I*, Selecta Mathematica **2** (1) (1996), 1 – 41.
- [10] M. Gerstenhaber, *On the deformation of rings and algebras*, Ann. of Math. **79** (1964), 59 – 103.
- [11] S. Majid, *Foundations of Quantum Group Theory*, Cambridge University Press, 1995.
- [12] N.Yu. Reshetikhin and M.A. Semenov-Tian-Shansky, *Quantum R-matrices and factorization problems*, J. Geom. Phys. **5** (4) (1988), 533 – 550.
- [13] M. Kontsevich, *Deformation Quantization of Poisson Manifolds*, Lett. Math. Phys. **66** (2003), 157–216.
- [14] A. Lichnerowicz, *Les variétés de Poisson et leurs algèbres de Lie associées*, J. Diff. Geom. **18** (1983), 523.
- [15] Y. Kosmann-Schwarzbach, *Integrability of Nonlinear Systems*, Lectures Notes in Physics 495, Springer, 1997, pp. 104-170.
- [16] C. Moreno and L. Valero, in Y.Choquet-Bruhat and C. De-Witt-Morette, *Analysis, Manifolds and Physics. Part II*, North-Holland, Amsterdam, 2000.

- [17] C. Moreno, *Produits star sur certains G/K kähleriens. Équation de Yang-Baxter et produits star sur G* , Géométrie symplectique et mécanique (La Grande Motte 1988), Lect. Notes Math. **1416** (1990), 210-234.
- [18] C. Moreno and L. Valero, *Produits star invariants et équation de Yang-Baxter quantique constante*, Dans les Actes des Journées Relativistes (24-29 avril 1990, Aussois, France).
- [19] C. Moreno and L. Valero, *Star products and quantum groups*, in Physics and Manifolds, Proceedings of the International Colloquium in honour of Yvonne Choquet-Bruhat, Mathematical Physics Studies, vol. **15**, Kluwer Academic Publishers, 1994, pp. 203–234.
- [20] C. Moreno and L. Valero, *The set of equivalent classes of invariant star products on $(G; \beta_1)$* , J. Geom. Phys. **23** (4) (1997), 360 – 378.
- [21] C. Moreno and J. Teles, *Preferred quantizations of nondegenerate triangular Lie bialgebras and Drinfeld associators*, in Recent Advances in Lie Theory, Research and Exposition in Mathematics, vol. **25**, Heldermann Verlag, 2002, pp. 346-365.
- [22] J. Teles, *The formal Chevalley cohomology in the quantization by Etingof and Kazhdan of non-degenerate triangular Lie bialgebras and their classical doubles*, Ph.D. thesis, Departamento de Matemática, Universidade de Coimbra, 2003.

CARLOS MORENO

DEPARTAMENTO DE FÍSICA TEÓRICA II, UNIVERSIDAD COMPLUTENSE, E-28040 MADRID, SPAIN.

JOANA TELES

CMUC, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF COIMBRA, COIMBRA, PORTUGAL

E-mail address: `jteles@mat.uc.pt`