

ON THE ALEKSANDROV-BAKELMAN-PUCCI ESTIMATE FOR THE INFINITY LAPLACIAN

FERNANDO CHARRO, GUIDO DE PHILIPPIS, AGNESE DI CASTRO
AND DAVI MÁXIMO

ABSTRACT: We prove L^∞ bounds and estimates of the modulus of continuity of solutions to the Poisson problem for the normalized infinity and p -Laplacian, namely

$$-\Delta_p^N u = f \quad \text{for } n < p \leq \infty.$$

We are able to provide a stable family of results depending continuously on the parameter p . We also prove the failure of the classical Alexandrov-Bakelman-Pucci estimate for the normalized infinity Laplacian and propose alternate estimates.

KEYWORDS: Infinity Laplacian, a priori estimates, Maximum Principle.

AMS SUBJECT CLASSIFICATION (2010): 35B45, 35B50, 35J15, 35J60, 35J70.

1. Introduction

In this paper we investigate the validity of Alexandrov-Bakelman-Pucci estimates (ABP for short) for the infinity Laplacian,

$$\Delta_\infty u := \langle D^2 u \nabla u, \nabla u \rangle \quad (1)$$

and its normalized version

$$\Delta_\infty^N u := \left\langle D^2 u \frac{\nabla u}{|\nabla u|}, \frac{\nabla u}{|\nabla u|} \right\rangle \quad (2)$$

as well as L^∞ and $C^{0,\alpha}$ estimates, uniform in p , for the solutions of the Poisson problem for the normalized p -Laplacian

$$\Delta_p^N u := \frac{1}{p} |\nabla u|^{2-p} \operatorname{div}(|\nabla u|^{p-2} \nabla u). \quad (3)$$

These type of normalized operators have recently received great attention, mainly since they have an interpretation in terms of random Tug-of-War games, see [22, 23].

Received August 2, 2011.

FC partially supported by a MEC-Fulbright fellowship and Spanish project MTM2010-18128, MICINN.

ADC supported by CMUC/FCT and project UTAustin/MAT/0035/2008.

DM partially supported by NFS grant DMS-0545984.

As an application of these estimates, in Section 6 we prove convergence as $p \rightarrow \infty$ of solutions to

$$-\Delta_p^N u = f, \quad (4)$$

to solutions of

$$-\Delta_\infty^N u = f. \quad (5)$$

It is interesting to compare this result with the results in [4]. An important feature of this limit process is the lack of variational structure of problem (4) that yields complications in the proof of the uniform convergence of the solutions (for instance, Morrey's estimates, are not available). It is in this context that our ABP estimate provides a stable L^∞ bound that can be used in combination with our stable regularity results to prove convergence.

The Alexandrov-Bakelman-Pucci maximum principle is well-known since the decade of 1960 in the context of linear uniformly elliptic equations. It can be stated as follows,

$$\sup_\Omega u \leq \sup_{\partial\Omega} u + C(n, \lambda, \Lambda) \text{diam}(\Omega) \|f\|_{L^n(\Omega)}, \quad (6)$$

for f the right-hand side of the equation and $0 < \lambda \leq \Lambda$ the ellipticity constants.

These estimates have been generalized in several directions, including uniformly elliptic fully nonlinear equations where they are the central tool in the proof of the Krylov-Safonov Harnack inequality and regularity theory (see [6] and the references therein).

In Theorem 4.3 we recall the classical ABP estimate for solutions of (4) (see [1, 6, 10, 16] for different proofs of this result) in order to stress that the constant blows up as $p \rightarrow \infty$. This fact is not merely technical, actually we are able to prove that the classical ABP estimate fails to hold for the normalized infinity Laplacian (see Section 7). An open question is if an estimate of the form (6) holds for some other integral norm $\|f\|_{L^p(\Omega)}$ with $p > n$.

The failure of the classical ABP estimate reflects the very high degeneracy of the equation as the normalized infinity Laplacian only controls the second derivative in the direction of the gradient. Remarkably, this is still enough to get an estimate of $\sup u$ of the type,

$$\left(\sup_\Omega u - \sup_{\partial\Omega} u \right)^2 \leq C \text{diam}(\Omega)^2 \int_{\sup_{\partial\Omega} u}^{\sup_\Omega u} \|f\|_{L^\infty(\{u=r\})} dr,$$

since the second derivatives in the directions tangential to the level sets can be controlled in average using the Gauss-Bonnet theorem, see Section 3 for a more precise statement. See also Section 7 for an example in which this estimate is sharper than the estimate with a plain $\|f\|_{L^\infty(\Omega)}$ in the right-hand side.

With these ideas we are able to provide a stable family of estimates depending continuously on the parameter $p \in (1, \infty]$. Incidentally, our ABP-type estimates turn out to be independent of the dimension n . Maybe this could have some interest in the context of Tug-of-War games in infinity dimensional spaces.

Let us now briefly discuss the $\mathcal{C}^{0,\alpha}$ estimates. The well-known Krylov-Safonov estimates in [6] apply to the normalized p -Laplacian whenever $p < \infty$, but degenerate as $p \rightarrow \infty$ as they depend upon the ratio between the ellipticity constants (see also [14, Section 9.7 and 9.8]), which in this case is $p - 1$ and blows-up as $p \rightarrow \infty$.

We prove Hölder estimates for solutions of the normalized p -Laplacian as well as Lipschitz estimates for the normalized ∞ -Laplacian. The main interest of this estimates is that they are stable in p , so that the whole range $n < p \leq \infty$ can be treated in a coherent way with all the parameters involved varying continuously.

The ideas behind our proof are somehow reminiscent of the comparison with cones property in [8] for the homogeneous infinity Laplace equation and the comparison with polar quadratic polynomials introduced in [2, 19] to study the non-homogeneous normalized ∞ -Laplace equation. Actually, our Lipschitz estimates in the case of the infinity Laplacian are closely related to [2, Lemma 5.6].

Although we shall not address this issue here, let us mention that existence of solutions to (4) and (5) is a nontrivial question as the normalized infinity and p -Laplacian are non-variational (in contrast to the regular p -Laplacian) and discontinuous operators (see Section 2).

Existence of viscosity solutions to (5) with Dirichlet boundary conditions has been proved using game-theoretic arguments, finite difference methods, and PDE techniques, see [2, 19, 22]. Moreover, solutions to the Dirichlet problem can be characterized using comparison with quadratic functions.

Uniqueness of solutions to the Dirichlet problem is known if the right-hand side $f \in \mathcal{C}(\Omega) \cap L^\infty(\Omega)$ is either 0, or has constant sign (see [2, 17, 19]).

When these conditions are not met, interesting non-uniqueness phenomena can happen, see [2] for a detailed discussion.

About (4), we first notice that when the right-hand side is 0, both the normalized and regular p -Laplacian coincide. For a non-trivial right-hand side, existence of viscosity solutions can be addressed with a game-theoretic approach (see [23]). However, to the best of our knowledge, conditions that guarantee uniqueness are not well understood.

The paper is organized as follows. In Section 2 we introduce some notation and preliminaries. In Sections 3 and 4 we prove the ABP-type estimates. Then, in Section 5 we provide stable estimates of the modulus of continuity of the solutions in the case $n < p \leq \infty$. Next, in Section 6 we justify the limits as $p \rightarrow \infty$ and finally, in Section 7 we provide examples showing the failure of an estimate of the form (6) and analyze the sharpness of our ABP-type estimate.

2. Notation and preliminaries

In this section we are going to state notations and recall some facts about the operators we are going to use in the sequel.

In what follows, $\Omega \subset \mathbb{R}^n$, $n \geq 2$, will be an open and bounded set. $B_r(x)$ is the open ball of radius r and center x . We shall denote the diameter of a domain by d or $\text{diam}(\Omega)$. We write $1_A(x)$ for the indicator function of a set A . Given two vectors $\xi, \eta \in \mathbb{R}^n$ we denote their scalar product as $\langle \xi, \eta \rangle = \sum_{i=1}^n \xi_i \eta_i$ and their tensor product as the matrix $\xi \otimes \eta = (\xi_i \eta_j)_{1 \leq i, j \leq n}$. S^n is the space of symmetric $n \times n$ matrices.

\mathcal{H}^{n-1} stands for the $n - 1$ dimensional Hausdorff measure, which for \mathcal{C}^1 manifolds coincide with the classical surface measure. Wherever not specified otherwise, ‘‘a.e.’’ will refer to the Lebesgue measure.

Given a function u , we shall denote $u^+ = \max\{0, u\}$ and $u^- = \max\{0, -u\}$ so that $u = u^+ - u^-$. We shall say that a function is twice differentiable at a point $x_0 \in \Omega$ if

$$u(x) = u(x_0) + \langle \xi, (x - x_0) \rangle + \frac{1}{2} \langle A(x - x_0), (x - x_0) \rangle + o(|x - x_0|^2)$$

as $|x - x_0| \rightarrow 0$ for some $\xi \in \mathbb{R}^n$ and $A \in S^n$, which are easily seen to be unique and will be denoted by $\nabla u(x_0)$ and $D^2 u(x_0)$ respectively.

A function u is semiconvex if $u(x) + C|x|^2$ is convex for some constant $C \in \mathbb{R}$. Geometrically, this means that the graph of u can be touched from below by a paraboloid of the type $a + \langle b, x \rangle - C|x|^2$ at every point.

The infinity Laplacian and p -Laplacian for $1 < p < \infty$, as well as its normalized versions have already been defined in the introduction, see (1), (2), and (3). For the record, let us recall here different expressions for the variational p -Laplacian, $1 < p < \infty$,

$$\begin{aligned} \Delta_p u &:= \operatorname{div}(|\nabla u|^{p-2} \nabla u) \\ &= |\nabla u|^{p-2} \cdot \operatorname{trace} \left[\left(I + (p-2) \frac{\nabla u \otimes \nabla u}{|\nabla u|^2} \right) D^2 u \right] \end{aligned} \quad (7)$$

and its normalized version

$$\begin{aligned} \Delta_p^N u &= \frac{1}{p} \operatorname{trace} \left[\left(I + (p-2) \frac{\nabla u \otimes \nabla u}{|\nabla u|^2} \right) D^2 u \right] \\ &= \frac{1}{p} \Delta u + \frac{p-2}{p} \Delta_\infty^N u \\ &= \frac{1}{p} |\nabla u| \operatorname{div} \left(\frac{\nabla u}{|\nabla u|} \right) + \frac{p-1}{p} \Delta_\infty^N u \\ &= \frac{1}{p} \Delta_1^N u + \frac{p-1}{p} \Delta_\infty^N u. \end{aligned} \quad (8)$$

It is worth emphasizing the lack of variational structure of (8), a uniformly elliptic operator in trace form, in contrast to the regular p -Laplacian (7), a quasilinear operator in divergence form.

Both normalized operators (2) and (8) (except for $p = 2$) are undefined when $\nabla u = 0$, where they have a bounded discontinuity, even if u is regular. This can be remediated adapting the notion of viscosity solution (see [9, Section 9] and also [13, Chapter 2]) using the upper and lower semicontinuous envelopes (relaxations) of the operator.

Let us recall here the definition of viscosity solution to be used in the sequel. To this end, given a function h defined in a set L , we need to introduce its *upper semicontinuous envelope* h^* and *lower semicontinuous envelope* h_* defined by

$$\begin{aligned} h^*(x) &= \limsup_{r \downarrow 0} \{h(y) : y \in \overline{B}_r(x) \cap L\} \\ h_*(x) &= \liminf_{r \downarrow 0} \{h(y) : y \in \overline{B}_r(x) \cap L\} \end{aligned}$$

as functions defined in \overline{L} .

Definition 2.1. Let $\mathcal{F}(\xi, X)$ be defined in a dense subset of $\mathbb{R}^n \times S^n$ with values in \mathbb{R} and assume $\mathcal{F}_* < \infty$ and $\mathcal{F}^* > -\infty$ in $\mathbb{R}^n \times S^n$.

1. An upper semicontinuous function $u : \Omega \rightarrow \mathbb{R}$ is a *viscosity subsolution* of

$$\mathcal{F}(\nabla u, D^2 u) = f(x) \quad (9)$$

in Ω if for all $x_0 \in \Omega$ and $\varphi \in \mathcal{C}^2(\Omega)$ such that $u - \varphi$ attains a local maximum at x_0 , one has

$$\mathcal{F}_*(\nabla \varphi(x_0), D^2 \varphi(x_0)) \leq f(x_0).$$

2. A lower semicontinuous function $u : \Omega \rightarrow \mathbb{R}$ is a *viscosity supersolution* of (9) in Ω if for all $x_0 \in \Omega$ and $\varphi \in \mathcal{C}^2(\Omega)$ such that $u - \varphi$ attains a local minimum at x_0 , one has

$$\mathcal{F}^*(\nabla \varphi(x_0), D^2 \varphi(x_0)) \geq f(x_0).$$

3. We say that u is a *viscosity solution* of (9) in Ω if it is both a viscosity subsolution and supersolution.

Given a symmetric matrix $X \in S^n$, we shall denote by $M(X)$ and $m(X)$ its greatest and smallest eigenvalues respectively, that is,

$$M(X) = \max_{|\xi|=1} \langle X\xi, \xi \rangle, \quad m(X) = \min_{|\xi|=1} \langle X\xi, \xi \rangle.$$

In the case of operators (2) and (8), which are continuous in $\mathbb{R}^n \setminus \{0\} \times S^n$, Definition 2.1 can be particularized as follows.

Definition 2.2. Let Ω be a bounded domain and $1 < p < \infty$. An upper semicontinuous function $u : \Omega \rightarrow \mathbb{R}$ is a *viscosity subsolution* of

$$-\Delta_p^N u(x) = f(x) \quad \text{in } \Omega, \quad (10)$$

if for all $x_0 \in \Omega$ and $\varphi \in \mathcal{C}^2(\Omega)$ such that $u - \varphi$ attains a local maximum at x_0 , one has

$$\begin{cases} -\Delta_p^N \varphi(x_0) \leq f(x_0) & \text{if } \nabla \varphi(x_0) \neq 0, \\ -\frac{1}{p} \Delta \varphi(x_0) - \frac{p-2}{p} M(D^2 \varphi(x_0)) \leq f(x_0) & \text{if } \nabla \varphi(x_0) = 0 \text{ and } p \in [2, \infty), \\ -\frac{1}{p} \Delta \varphi(x_0) - \frac{p-2}{p} m(D^2 \varphi(x_0)) \leq f(x_0) & \text{if } \nabla \varphi(x_0) = 0 \text{ and } p \in (1, 2], \end{cases}$$

with $\Delta_p^N u$ given by (8). A lower semicontinuous function $u : \Omega \rightarrow \mathbb{R}$ is a *viscosity supersolution* of (10) in Ω if for all $x_0 \in \Omega$ and $\varphi \in \mathcal{C}^2(\Omega)$ such that $u - \varphi$ attains a local minimum at x_0 , one has

$$\begin{cases} -\Delta_p^N \varphi(x_0) \geq f(x_0) & \text{if } \nabla \varphi(x_0) \neq 0, \\ -\frac{1}{p} \Delta \varphi(x_0) - \frac{p-2}{p} m(D^2 \varphi(x_0)) \geq f(x_0) & \text{if } \nabla \varphi(x_0) = 0 \text{ and } p \in [2, \infty), \\ -\frac{1}{p} \Delta \varphi(x_0) - \frac{p-2}{p} M(D^2 \varphi(x_0)) \geq f(x_0) & \text{if } \nabla \varphi(x_0) = 0 \text{ and } p \in (1, 2]. \end{cases}$$

We say that u is a *viscosity solution* of (10) in Ω if it is both a viscosity subsolution and supersolution.

Remark 2.3. Geometrically, the condition “ $u - \varphi$ attains a local maximum (minimum) at x_0 ” means that up to a vertical translation, the graph of φ touches the graph of u from above (below).

Notice that although this definition is slightly different to the one in [23], it turns to be equivalent, see [18].

It is worth comparing this definition with the situation in the variational case in the range $1 < p < 2$. In that case the singularity in (7) when written in trace form is not bounded and it is not possible to use the theory mentioned above. Instead, one can adopt the definition proposed in a series of papers by Birindelli and Demengel, see [5] and the references therein. An alternative but equivalent definition in the case $f = 0$ can be found in [13, Section 2.1.3].

Analogously, we have the definition of solution for the normalized ∞ -Laplacian (2).

Definition 2.4. Let Ω be a bounded domain. An upper semicontinuous function $u : \Omega \rightarrow \mathbb{R}$ is a *viscosity subsolution* of

$$-\Delta_\infty^N u(x) = f(x) \quad \text{in } \Omega, \tag{11}$$

if for all $x_0 \in \Omega$ and $\varphi \in \mathcal{C}^2(\Omega)$ such that $u - \varphi$ attains a local maximum at x_0 , one has

$$\begin{cases} -\Delta_\infty^N \varphi(x_0) \leq f(x_0), & \text{whenever } \nabla \varphi(x_0) \neq 0, \\ -M(D^2 \varphi(x_0)) \leq f(x_0), & \text{otherwise.} \end{cases} \tag{12}$$

A lower semicontinuous function $u : \Omega \rightarrow \mathbb{R}$ is a *viscosity supersolution* of (11) in Ω if for all $x_0 \in \Omega$ and $\varphi \in \mathcal{C}^2(\Omega)$ such that $u - \varphi$ attains a local minimum at x_0 , one has

$$\begin{cases} -\Delta_{\infty}^N \varphi(x_0) \geq f(x_0), & \text{whenever } \nabla \varphi(x_0) \neq 0, \\ -m(D^2 \varphi(x_0)) \geq f(x_0), & \text{otherwise.} \end{cases} \quad (13)$$

We say that u is a *viscosity solution* of (11) in Ω if it is both a viscosity subsolution and supersolution.

The above definition agrees with the one in [19, 21] and is slightly different from the one in [22]. Anyway, it is easy to see that all this definitions are equivalent.

3. The ABP estimate for the infinity Laplacian

There are two parts in the proof of the classical ABP estimate. The first one is entirely geometric and allows to bound $\sup_{\Omega} u$ in terms of the integral of $\det D^2 u$ as follows,

$$\left(\frac{\sup_{\Omega} u - \sup_{\partial\Omega} u}{\text{diam}(\Omega)} \right)^n \leq C(n) \int_{\{u=\Gamma(u)\}} |\det D^2 u(x)| dx \quad (14)$$

where $\Gamma(u)$ is the concave envelope of u (see Definition 3.1 below). The argument leading to (14) is entirely geometric and does not involve any equation for u .

In the second part we take advantage of the fact that we are working in the contact set (so that the Hessian has a sign) and use the inequality between the arithmetic and geometric means to bound the determinant in terms of the trace. In the particular case of the Laplacian, the following estimate is obtained,

$$\int_{\{u=\Gamma(u)\}} |\det D^2 u(x)| dx \leq n^{-n} \int_{\{u=\Gamma(u)\}} (-\text{trace } D^2 u(x))^n dx. \quad (15)$$

Estimates (14) and (15) altogether yield the ABP estimate

$$\left(\frac{\sup_{\Omega} u - \sup_{\partial\Omega} u}{\text{diam}(\Omega)} \right)^n \leq C(n) \int_{\{u=\Gamma(u)\}} (-\Delta u(x))^n dx.$$

At this stage one can use the equation for u , namely $-\Delta u \leq f$ to control the right-hand side of the estimate.

It is apparent from the above discussion that if we are able to estimate $|\det D^2u|$ in terms of the infinity Laplacian we shall get an ABP-type estimate. However, a different kind of estimate is expected to hold since, as we have mentioned in the introduction, the classical ABP estimate is false for the infinity Laplacian (see Section 7).

We have to recall some necessary notions (see [6]).

Definition 3.1. Let u be a continuous function in an open convex set A . The concave envelope of u in A is defined by

$$\begin{aligned}\Gamma(u)(x) &= \inf_w \{w(x) : w \geq u \text{ in } A, w \text{ concave in } A\} \\ &= \inf_L \{L(x) : L \geq u \text{ in } A, L \text{ is affine}\}\end{aligned}$$

for $x \in A$. The upper contact set of u is defined as,

$$C^+(u) = \{x \in A : u(x) = \Gamma(u)(x)\}.$$

The following is one of the main results in this section.

Theorem 3.2. Let $f \in \mathcal{C}(\Omega)$ and consider $u \in \mathcal{C}(\overline{\Omega})$ that satisfies

$$-\Delta_\infty^N u \leq f(x) \quad \text{in } \Omega \tag{16}$$

in the viscosity sense. Then, we have

$$\left(\sup_\Omega u - \sup_{\partial\Omega} u^+\right)^2 \leq 2d^2 \int_{\sup_{\partial\Omega} u^+}^{\sup_\Omega u} \|f^+ \cdot 1_{C^+(u)}\|_{L^\infty(\{u^+=r\})} dr, \tag{17}$$

where $d = \text{diam}(\Omega)$. Analogously, whenever

$$-\Delta_\infty^N u \geq f(x) \quad \text{in } \Omega \tag{18}$$

in the viscosity sense, the following estimate holds,

$$\left(\sup_\Omega u^- - \sup_{\partial\Omega} u^-\right)^2 \leq 2d^2 \int_{\sup_{\partial\Omega} u^-}^{\sup_\Omega u^-} \|f^- \cdot 1_{C^+(-u)}\|_{L^\infty(\{u^-=r\})} dr. \tag{19}$$

Proof: We prove the first inequality, since the second one is similar. The proof is divided in two steps, in the first one we prove the result in the case of a semiconvex subsolution v of (16) and then we shall remove this assumption in the second step.

STEP 1. We can assume without loss of generality that $\sup_{\partial\Omega} v^+ = 0$ and $v(x_0) > 0$ for some $x_0 \in \Omega$. Let us consider the function v^+ extended by 0 to the whole \mathbb{R}^n . Notice that v^+ is also semiconvex and satisfies

$$-\Delta_\infty^N v^+ \leq f^+ \quad \text{in } \mathbb{R}^n.$$

Let $\Omega^* = \text{conv}(\Omega)$, the convex hull of Ω and for every $\sigma > 0$ let us consider $\Gamma_\sigma(v^+)$, the concave envelope of v^+ in Ω_σ^* , where

$$\Omega_\sigma^* = \{x \in \mathbb{R}^n \text{ such that } \text{dist}(x, \Omega^*) \leq \sigma\}.$$

Since we are assuming that v is semiconvex, Lemma A.1 implies that $\Gamma_\sigma(v^+)$ is $\mathcal{C}_{\text{loc}}^{1,1}(\Omega_\sigma^*)$.

Our aim is to estimate

$$\int_{\nabla\Gamma_\sigma(v^+)(\Omega_\sigma^*)} |\xi|^{2-n} d\xi, \quad (20)$$

both from above and from below. For the sake of brevity, let us denote in the sequel

$$M := \sup_{\Omega_\sigma^*} \Gamma_\sigma(v^+) = \sup_{\Omega} v^+ = \sup_{\Omega} v$$

and

$$A_\sigma = C_\sigma^+(v^+) \cap \{\text{Points of twice differentiability for } \Gamma_\sigma(v^+)\},$$

where $C_\sigma^+(v^+)$ is the set of points in Ω_σ^* where $v^+ = \Gamma_\sigma(v^+)$. First we estimate (20) from above. Applying the Area Formula for Lipschitz maps [12, Theorem 3.2.5], Lemma A.1, and then the Coarea Formula [12, Theorem 3.2.11], we get,

$$\begin{aligned} & \int_{\nabla\Gamma_\sigma(v^+)(\Omega_\sigma^*)} |\xi|^{2-n} d\xi \leq \int_{\Omega_\sigma^*} |\nabla\Gamma_\sigma(v^+)|^{2-n} \cdot \det(-D^2\Gamma_\sigma(v^+)) dx \\ & = \int_{\Omega_\sigma^*} 1_{A_\sigma}(x) \cdot |\nabla\Gamma_\sigma(v^+)|^{2-n} \cdot \det(-D^2\Gamma_\sigma(v^+)) dx \\ & = \int_0^M \int_{\{\Gamma_\sigma(v^+)=r\}} 1_{A_\sigma}(x) \cdot |\nabla\Gamma_\sigma(v^+)|^{1-n} \cdot \det(-D^2\Gamma_\sigma(v^+)) d\mathcal{H}^{n-1} dr. \end{aligned} \quad (21)$$

From Hadamard's formula [3, Section 2.10] for the determinant of a positive definite matrix and Proposition A.3, we get that for a.e. $r \in (0, M)$,

$$\det(-D^2\Gamma_\sigma(v^+)) \leq -\Delta_\infty^N \Gamma_\sigma(v^+) |\nabla\Gamma_\sigma(v^+)|^{n-1} \prod_{i=1}^{n-1} \kappa_i \quad (22)$$

holds \mathcal{H}^{n-1} -a.e. in $\{\Gamma_\sigma(v^+) = r\}$, for $\kappa_i(x)$ the principal curvatures of $\{\Gamma_\sigma(v^+) = r\}$ at x , as defined in Proposition A.3. Hence, from (21), (22) and Lemma

A.2 we have,

$$\begin{aligned}
& \int_{\nabla\Gamma_\sigma(v^+)(\Omega_\sigma^*)} |\xi|^{2-n} d\xi \\
& \leq \int_0^M \int_{\{\Gamma_\sigma(v^+)=r\}} 1_{A_\sigma}(x) \cdot f^+(x) \cdot \prod_{i=1}^{n-1} \kappa_i(x) d\mathcal{H}^{n-1} dr \\
& \leq \int_0^M \|f^+ \cdot 1_{C_\sigma^+(v^+)}\|_{L^\infty(\{v^+=r\})} \int_{\{\Gamma_\sigma(v^+)=r\}} \prod_{i=1}^{n-1} \kappa_i(x) d\mathcal{H}^{n-1} dr \\
& \leq \mathcal{H}^{n-1}(\partial B_1(0)) \int_0^M \|f^+ \cdot 1_{C_\sigma^+(v^+)}\|_{L^\infty(\{v^+=r\})} dr.
\end{aligned} \tag{23}$$

The last step follows from the Gauss-Bonnet theorem, Theorem A.4.

To estimate (20) from below, we observe that

$$\partial K_{x_0,\sigma}(\Omega_\sigma^*) \subset \nabla\Gamma_\sigma(v^+)(\Omega_\sigma^*)$$

where x_0 is a maximum point in Ω^* for $v^+(x)$, $K_{x_0,\sigma}(x)$ is the concave cone with vertex in $(x_0, v^+(x_0))$ and base Ω_σ^* and $\partial K_{x_0,\sigma}$ its super-differential (see [15, Section 1.4]). Collecting the previous information we get

$$\begin{aligned}
& \int_{\partial K_{x_0,\sigma}(\Omega_\sigma^*)} |\xi|^{2-n} d\xi \\
& \leq \mathcal{H}^{n-1}(\partial B_1(0)) \int_0^M \|f^+ \cdot 1_{C_\sigma^+(v^+)}\|_{L^\infty(\{v^+=r\})} dr.
\end{aligned} \tag{24}$$

We want now to pass to the limit as σ goes to 0. It is easy to see that $\partial K_{x_0,\sigma}(\Omega_\sigma^*) \uparrow \partial K_{x_0}(\Omega^*)$. We need now to estimate the limit of the right-hand side, to do this notice that $\Gamma_\sigma(v^+)$ is a decreasing sequence of concave functions which thus converges to a concave function w defined on Ω such that $w \geq \Gamma(v^+) \geq v^+$, so in particular

$$\limsup_{\sigma \rightarrow 0} C_\sigma^+(v^+) \subset \{w = v^+\} \subset C^+(v^+).$$

The last equation implies that for every $r \in (0, M)$

$$\limsup_{\sigma \rightarrow 0} \|f^+ \cdot 1_{C_\sigma^+(v^+)}\|_{L^\infty(\{v^+=r\})} \leq \|f^+ \cdot 1_{C^+(v^+)}\|_{L^\infty(\{v^+=r\})}.$$

Thus, (24) becomes

$$\begin{aligned} \int_{\partial K_{x_0}(\Omega^*)} |\xi|^{2-n} d\xi \\ \leq \mathcal{H}^{n-1}(\partial B_1(0)) \int_0^M \|f^+ \cdot 1_{C^+(v^+)}\|_{L^\infty(\{v^+=r\})} dr. \end{aligned} \quad (25)$$

It is now very easy to see that

$$B_{\frac{M}{d}}(0) \subset \partial K_{x_0}(\Omega^*).$$

where $d = \text{diam}(\Omega^*) = \text{diam}(\Omega)$ and $M = \sup_{\Omega} v = \sup_{\Omega^*} K_{x_0}(x)$. Consequently,

$$\int_{\partial K_{x_0}(\Omega^*)} |\xi|^{2-n} d\xi \geq \int_{B_{\frac{M}{d}}(0)} |\xi|^{2-n} d\xi = \frac{\mathcal{H}^{n-1}(\partial B_1(0))}{2} \left(\frac{M}{d}\right)^2. \quad (26)$$

Collecting (25) and (26) we arrive at

$$(\sup_{\Omega} v^+)^2 \leq 2d^2 \int_0^{\sup_{\Omega} v^+} \|f^+ \cdot 1_{C^+(v^+)}\|_{L^\infty(\{v^+=r\})} dr.$$

STEP 2. Now we consider the case of a general viscosity subsolution u of (16). Then, the sup-convolution of u , defined as

$$u^\varepsilon(x) = \sup_{y \in \bar{\Omega}} \left\{ u(y) - \frac{1}{2\varepsilon} |x - y|^2 \right\}, \quad (27)$$

is semiconvex and verifies

$$-\Delta_\infty^N u^\varepsilon \leq f_\varepsilon(x) := \sup \left\{ f(y) : y \in B_{2(\varepsilon\|u\|_{L^\infty(\Omega)})^{1/2}}(x) \right\} \quad \text{in } \Omega_\varepsilon$$

where

$$\Omega_\varepsilon := \{x \in \Omega : \text{dist}(x, \partial\Omega) > 2(\varepsilon\|u\|_{L^\infty(\Omega)})^{1/2}\}.$$

Moreover, $u \leq u^\varepsilon$ and $u^\varepsilon \rightarrow u$ uniformly as $\varepsilon \rightarrow 0$ in $\bar{\Omega}$, see [6, Chapter 5].

From the previous step we know that

$$\begin{aligned} & \left(\sup_{\Omega_\varepsilon} (u^\varepsilon)^+ - \sup_{\partial\Omega_\varepsilon} (u^\varepsilon)^+ \right)^2 \\ & \leq 2 \text{diam}(\Omega_\varepsilon)^2 \int_{\sup_{\partial\Omega_\varepsilon} (u^\varepsilon)^+}^{\sup_{\Omega_\varepsilon} (u^\varepsilon)^+} \|f_\varepsilon^+ \cdot 1_{C^+((u^\varepsilon)^+)}\|_{L^\infty(\{(u^\varepsilon)^+=r\})} dr. \end{aligned}$$

If we are able to pass to the limit in the previous inequality we shall obtain the desired estimate. It is easy to see that actually we can pass to the limit once we have proved that

$$\limsup_{\varepsilon \rightarrow 0} \|f_\varepsilon^+ \cdot 1_{C^+((u^\varepsilon)^+)}\|_{L^\infty(\{(u^\varepsilon)^+=r\})} \leq \|f^+ \cdot 1_{C^+(u^+)}\|_{L^\infty(\{u^+=r\})}.$$

Let us consider a sequence $\{\varepsilon_j\}$ such that the lim sup is realized and let $x_{\varepsilon_j} \in \{(u^{\varepsilon_j})^+ = r\}$ such that

$$\|f_{\varepsilon_j}^+ \cdot 1_{C^+((u^{\varepsilon_j})^+)}\|_{L^\infty(\{(u^{\varepsilon_j})^+=r\})} = f_{\varepsilon_j}^+(x_{\varepsilon_j}) \cdot 1_{C^+((u^{\varepsilon_j})^+)}(x_{\varepsilon_j}).$$

Up to a subsequence, still denoted with ε_j , we can assume that x_{ε_j} converges to $x_0 \in \overline{\Omega}$. Since x_{ε_j} belongs to $\{(u^{\varepsilon_j})^+ = r\}$ and u^{ε_j} converge uniformly to u we obtain $u^+(x_0) = r$, in particular $x_0 \notin \partial\Omega$ for any $r > \sup_{\partial\Omega} u$.

Now since f_{ε_j} converge uniformly to f on compact subsets of Ω and

$$\limsup_{\varepsilon_j \rightarrow 0} C^+((u^{\varepsilon_j})^+) \subset C^+(u^+)$$

thanks to [7, Lemma A.1], we get

$$\begin{aligned} \limsup_{\varepsilon_j \rightarrow 0} f_{\varepsilon_j}^+(x_{\varepsilon_j}) \cdot 1_{C^+((u^{\varepsilon_j})^+)}(x_{\varepsilon_j}) \\ \leq f^+(x_0) \cdot 1_{C^+(u^+)}(x_0) \leq \|f^+ \cdot 1_{C^+(u^+)}\|_{L^\infty(\{u^+=r\})} \end{aligned} \quad (28)$$

and hence the thesis follows. \blacksquare

The following is an easy consequence of Theorem 3.2.

Corollary 3.3. *Let $f \in \mathcal{C}(\Omega)$ and consider $u \in \mathcal{C}(\overline{\Omega})$ that satisfies*

$$-\Delta_\infty^N u \leq f(x) \quad \text{in } \Omega$$

in the viscosity sense. Then, the following estimate holds,

$$\sup_\Omega u \leq \sup_{\partial\Omega} u^+ + 2d^2 \|f^+\|_{L^\infty(C^+(u))}.$$

Analogously, whenever

$$-\Delta_\infty^N u \geq f(x) \quad \text{in } \Omega$$

in the viscosity sense,

$$\sup_\Omega u^- \leq \sup_{\partial\Omega} u^- + 2d^2 \|f^-\|_{L^\infty(C^+(-u))},$$

where $d = \text{diam}(\Omega)$.

Remark 3.4. Similar results can be derived from Theorems 3.5, 4.1, and 4.4 below.

If we replace (20) by

$$\int_{\nabla\Gamma_\sigma(v^+)(\Omega_\sigma^*)} |\xi|^{4-n} d\xi,$$

the same proof yields the following result.

Theorem 3.5. *Let $f \in \mathcal{C}(\Omega)$ and consider $u \in \mathcal{C}(\overline{\Omega})$ that satisfies*

$$-\Delta_\infty u \leq f(x) \quad \text{in } \Omega$$

in the viscosity sense. Then, we have

$$\left(\sup_\Omega u - \sup_{\partial\Omega} u^+ \right)^4 \leq 4 \cdot d^4 \int_{\sup_{\partial\Omega} u^+}^{\sup_\Omega u} \|f^+ \cdot 1_{C^+(u)}\|_{L^\infty(\{u^+=r\})} dr,$$

where $d = \text{diam}(\Omega)$. Analogously, whenever

$$-\Delta_\infty u \geq f(x) \quad \text{in } \Omega$$

in the viscosity sense, the following estimate holds,

$$\left(\sup_\Omega u^- - \sup_{\partial\Omega} u^- \right)^4 \leq 4 \cdot d^4 \int_{\sup_{\partial\Omega} u^-}^{\sup_\Omega u^-} \|f^- \cdot 1_{C^+(-u)}\|_{L^\infty(\{u^-=r\})} dr.$$

4. An stable ABP for the normalized p-Laplacian

The following is one of the main results in this section.

Theorem 4.1. *Let $1 < p < \infty$, $f \in \mathcal{C}(\Omega)$ and consider $u \in \mathcal{C}(\overline{\Omega})$ that satisfies*

$$-\Delta_p^N u \leq f(x) \quad \text{in } \Omega$$

in the viscosity sense. Then, we have

$$\left(\sup_\Omega u - \sup_{\partial\Omega} u^+ \right)^2 \leq \frac{2p d^2}{p-1} \int_{\sup_{\partial\Omega} u^+}^{\sup_\Omega u} \|f^+ \cdot 1_{C^+(u)}\|_{L^\infty(\{u^+=r\})} dr, \quad (29)$$

where $d = \text{diam}(\Omega)$. Analogously, whenever

$$-\Delta_p^N u \geq f(x) \quad \text{in } \Omega$$

in the viscosity sense, the following estimate holds,

$$\left(\sup_\Omega u^- - \sup_{\partial\Omega} u^- \right)^2 \leq \frac{2p d^2}{p-1} \int_{\sup_{\partial\Omega} u^-}^{\sup_\Omega u^-} \|f^- \cdot 1_{C^+(-u)}\|_{L^\infty(\{u^-=r\})} dr. \quad (30)$$

Proof: We only sketch the proof of the first inequality, since the second one is similar. The argument goes exactly in the same way of the one of Theorem 3.2 with the only observation that thanks to the concavity of $\Gamma_\sigma(u^+)$ and Lemmas A.1 and A.2 and Proposition A.3, at almost every contact point $x \in C_\sigma^+(u^+)$ one has,

$$\begin{aligned} & -\left(\frac{p-1}{p}\right) \Delta_\infty^N \Gamma_\sigma(u^+)(x) \\ & \leq -\left(\frac{p-1}{p}\right) \Delta_\infty^N \Gamma_\sigma(u^+)(x) + \frac{1}{p} |\nabla \Gamma_\sigma(u^+)(x)| \sum_{i=1}^{n-1} \kappa_i(x) \\ & = -\Delta_p^N \Gamma_\sigma(u^+)(x) \leq f^+(x), \end{aligned}$$

where $\kappa_i(x)$ $i = 1, \dots, n-1$ denote, as before, the principal curvatures at x of the level set of $\Gamma_\sigma(u^+)$ passing through x . \blacksquare

Remark 4.2. Estimates (29) and (30) are stable as $p \rightarrow \infty$ and, moreover, allow to recover estimates (17) and (19) in the limit.

It is interesting to compare estimates (29) and (30) with the usual ABP estimates that are not stable in the limit.

The following theorem can be easily proved with the techniques of [1, 6, 10, 16], the proof we sketch here actually allows a comparison with the previous one, see the remarks at the end of the proof.

Theorem 4.3. *Let $1 < p < \infty$, $f \in \mathcal{C}(\Omega)$ and consider $u \in \mathcal{C}(\overline{\Omega})$ that satisfies*

$$-\Delta_p^N u \leq f(x) \quad \text{in } \Omega,$$

in the viscosity sense. Then, the following estimate holds,

$$\sup_{\Omega} u \leq \sup_{\partial\Omega} u^+ + \frac{dp}{n|B_1(0)|^{\frac{1}{n}}(p-1)^{\frac{1}{n}}} \|f^+\|_{L^n(C^+(u))},$$

where $d = \text{diam}(\Omega)$.

Analogously, whenever $u \in \mathcal{C}(\overline{\Omega})$ satisfies

$$-\Delta_p^N u \geq f(x) \quad \text{in } \Omega$$

in the viscosity sense, the following estimate holds,

$$\sup_{\Omega} u^- \leq \sup_{\partial\Omega} u^- + \frac{dp}{n|B_1(0)|^{\frac{1}{n}}(p-1)^{\frac{1}{n}}} \|f^-\|_{L^n(C^+(-u))},$$

Proof: Again we don't take care of regularity issues which can be easily handled with the technique we used in the proof of Theorem 3.2.

The standard ABP argument yields,

$$\begin{aligned} & \left(\frac{\sup_{\Omega} u - \sup_{\partial\Omega} u^+}{d} \right)^n |B_1(0)| \\ & \leq |\nabla\Gamma(u^+)(C^+(u))| \leq \int_{C^+(u^+)} \det(-D^2u) dx. \end{aligned}$$

Proposition A.3 yields

$$\int_{C^+(u^+)} \det(-D^2u) dx \leq \int_{C^+(u^+)} -\Delta_{\infty}^N \Gamma(u^+) \cdot |\nabla\Gamma(u^+)|^{n-1} \cdot \prod_{i=1}^{n-1} \kappa_i dx.$$

Now, we multiply and divide by $p - 1$ the right-hand side of the previous inequality, and then apply the inequality between the arithmetic and geometric mean inequalities. We get,

$$\begin{aligned} & \int_{C^+(u^+)} -\Delta_{\infty}^N \Gamma(u^+) \cdot |\nabla\Gamma(u^+)|^{n-1} \cdot \prod_{i=1}^{n-1} \kappa_i dx \\ & \leq \frac{1}{(p-1)n^n} \int_{C^+(u^+)} \left(-(p-1)\Delta_{\infty}^N \Gamma(u^+) + |\nabla\Gamma(u^+)| \sum_{i=1}^{n-1} \kappa_i \right)^n dx \\ & = \frac{p^n}{(p-1)n^n} \int_{C^+(u^+)} \left(-\Delta_p^N \Gamma(u^+) \right)^n dx \\ & \leq \frac{p^n}{(p-1)n^n} \int_{C^+(u)} (f^+)^n dx. \end{aligned}$$

■

Some comments are in order. First, it is important to notice that the estimate in Theorem 4.3 is not stable in p , as the resulting constant blows up as $p \rightarrow \infty$. Hence we don't recover any estimate in the limit, in contrast to Theorem 4.1 that yields Theorem 3.2 as a limit case.

This fact can be understood by comparing the proofs of Theorems 4.1 and 4.3. In fact, they make apparent that the infinity Laplacian controls one single direction, the direction of steepest descent, totally neglecting the curvature of the level sets (whose average is in turn controlled by the Gauss-Bonnet Theorem). The p -Laplacian, instead, is a weighted mean of the curvatures

of the level sets and the normal direction. In particular, when $p = 2$, the classical Laplacian, all the directions are weighted the same way.

In this sense, it is completely natural that in the p -Laplacian case the ABP involves an integral norm, an average in all directions, while in the infinity Laplacian case the average is somehow unidimensional.

If we replace (20) by

$$\int_{\nabla\Gamma_\sigma(v^+)(\Omega_\sigma^*)} |\xi|^{p-n} d\xi,$$

the same proof of Theorem 4.1 yields similar estimates for the variational p -Laplacian (7).

Theorem 4.4. *Let $1 < p < \infty$, $f \in \mathcal{C}(\Omega)$ and consider $u \in \mathcal{C}(\bar{\Omega})$ that satisfies*

$$-\Delta_p u \leq f(x) \quad \text{in } \Omega$$

in the viscosity sense. Then, we have

$$\left(\sup_{\Omega} u - \sup_{\partial\Omega} u^+ \right)^p \leq \frac{p d^p}{p-1} \int_{\sup_{\partial\Omega} u^+}^{\sup_{\Omega} u} \|f^+ \cdot 1_{C^+(u)}\|_{L^\infty(\{u^+=r\})} dr,$$

where $d = \text{diam}(\Omega)$. Analogously, whenever

$$-\Delta_p u \geq f(x) \quad \text{in } \Omega$$

in the viscosity sense, the following estimate holds,

$$\left(\sup_{\Omega} u^- - \sup_{\partial\Omega} u^- \right)^p \leq \frac{p d^p}{p-1} \int_{\sup_{\partial\Omega} u^-}^{\sup_{\Omega} u^-} \|f^- \cdot 1_{C^+(-u)}\|_{L^\infty(\{u^-=r\})} dr.$$

5. Stable estimates of the modulus of continuity for $p \leq \infty$

In this section we obtain Hölder estimates for solutions of the normalized p -Laplacian as well as Lipschitz estimates for the normalized ∞ -Laplacian. The main interest of this estimates is that they are stable in p , and apply to the whole range $n < p \leq \infty$ with all the parameters involved varying continuously.

As mentioned in the introduction, the well-known estimates in [6] apply whenever $p < \infty$, but degenerate as $p \rightarrow \infty$ as they depend upon the ratio between the ellipticity constants (see also [14], Section 9.7 and 9.8), which in this case is $p - 1$ and blows-up as $p \rightarrow \infty$.

For the sake of a unified presentation, throughout this section we are going to denote

$$\alpha_p = \begin{cases} \frac{n-1}{p-1} & \text{for } n < p < \infty \\ 0 & \text{for } p = \infty \end{cases} \quad (31)$$

and

$$C_p = \begin{cases} \frac{p}{p+n-2} & \text{for } n < p < \infty \\ 1 & \text{for } p = \infty. \end{cases} \quad (32)$$

Theorem 5.1. *Let Ω be a bounded domain, $2 \leq n < p \leq \infty$, and u a viscosity solution of*

$$-\Delta_p^N u = f \quad \text{in } \Omega. \quad (33)$$

Then, for any $x \in \Omega$

$$\frac{|u(y) - u(x)|}{|y - x|^{1-\alpha_p}} \leq \sup_{z \in \partial\Omega} \frac{|u(z) - u(x)|}{|z - x|^{1-\alpha_p}} + \frac{C_p}{1 - \alpha_p} \text{diam}(\Omega)^{1+\alpha_p} \|f\|_{L^\infty(\Omega)}$$

for every $y \in \bar{\Omega}$, with α_p and C_p defined in (31) and (32) respectively.

Remark 5.2. Compare with [8, Lemma 2.5].

The following estimate is an easy consequence of Theorem 5.1.

Corollary 5.3. *Let Ω be a bounded domain, $2 \leq n < p \leq \infty$, and u a viscosity solution of (33). Let α_p and C_p defined by (31) and (32). Then, for any $x \in \Omega$ we have that*

$$\frac{|u(y) - u(x)|}{|y - x|^{1-\alpha_p}} \leq \frac{2\|u\|_{L^\infty(\Omega)}}{\text{dist}(x, \partial\Omega)^{1-\alpha_p}} + \frac{C_p}{1 - \alpha_p} \text{diam}(\Omega)^{1+\alpha_p} \|f\|_{L^\infty(\Omega)}$$

for every $y \in \bar{\Omega}$.

We also have global estimates for the Dirichlet problem.

Theorem 5.4. *Let Ω be a bounded domain, $2 \leq n < p \leq \infty$, and u a viscosity solution of*

$$\begin{cases} -\Delta_p^N u = f & \text{in } \Omega \\ u = g & \text{on } \partial\Omega \end{cases} \quad (34)$$

with $g \in C^{0,1-\alpha_p}(\partial\Omega)$ and α_p as in (31). Then, for every $x, y \in \bar{\Omega}$,

$$\frac{|u(x) - u(y)|}{|x - y|^{1-\alpha_p}} \leq C_p \|f\|_{L^\infty(\Omega)} \text{diam}(\Omega)^{1+\alpha_p} + [g]_{1-\alpha_p, \partial\Omega} \quad (35)$$

with C_p as in (32).

Corollary 5.5. *Let Ω be a bounded domain, $2 \leq n < p \leq \infty$, and u a viscosity solution of (34) with $f \in \mathcal{C}(\Omega)$, $g \in \mathcal{C}^{0,1-\alpha_p}(\partial\Omega)$ and α_p , C_p as in (31) and (32) respectively. Then,*

$$\begin{aligned} \|u\|_{\mathcal{C}^{0,1-\alpha_p}(\Omega)} &= \|u\|_{L^\infty(\Omega)} + \sup_{\substack{x,y \in \Omega \\ x \neq y}} \frac{|u(x) - u(y)|}{|x - y|^{1-\alpha_p}} \\ &\leq \|g\|_{\mathcal{C}^{0,1-\alpha_p}(\partial\Omega)} + \left(2 \tilde{C}_p \text{diam}(\Omega)^2 + C_p \text{diam}(\Omega)^{1+\alpha_p}\right) \|f\|_{L^\infty(\Omega)} \end{aligned}$$

with

$$\tilde{C}_p = \begin{cases} \frac{p}{p-1} & \text{for } n < p < \infty \\ 1 & \text{for } p = \infty. \end{cases} \quad (36)$$

Proof: From the ABP estimates (17) and (29) in Theorems 3.2 and 4.1, we have

$$\|u\|_{L^\infty(\Omega)} \leq \|g\|_{L^\infty(\partial\Omega)} + 2 \tilde{C}_p \text{diam}(\Omega)^2 \|f\|_{L^\infty(C^+(u))}.$$

for \tilde{C}_p as in (36). This estimate together with (35) yields the result. \blacksquare

We need some lemmas in the proof of Theorems 5.1 and 5.4.

Lemma 5.6. *Consider the function*

$$v(x) = h + \frac{A}{1-\alpha_p} |x - y|^{1-\alpha_p} - \frac{B}{2} |x - y|^2$$

with $A, B > 0$, $h \in \mathbb{R}$ and $y \in \mathbb{R}^n$. Let C_p as defined in (32). Then

$$-\Delta_p^N v(x) = C_p^{-1} B$$

for $x \neq y$ such that $\nabla v(x) \neq 0$.

Proof: For $x \neq y$ we have

$$\nabla v(x) = (A |x - y|^{-1-\alpha_p} - B) (x - y)$$

and

$$D^2 v(x) = (A |x - y|^{-1-\alpha_p} - B) \text{Id} - A(1 + \alpha_p) |x - y|^{-1-\alpha_p} \frac{(x - y) \otimes (x - y)}{|x - y|^2}.$$

Then,

$$\Delta_\infty^N v(x) = -A \alpha_p |x - y|^{-1-\alpha_p} - B$$

and the proof is complete in the case $p = \infty$. If $p < \infty$,

$$\Delta v(x) = \text{trace}(D^2 v(x)) = A(n-1-\alpha_p)|x-y|^{-1-\alpha_p} - nB$$

and hence

$$\Delta_p^N v(x) = \frac{1}{p}\Delta v(x) + \frac{p-2}{p}\Delta_\infty^N v(x) = -C_p^{-1}B. \quad \blacksquare$$

Remark 5.7. Notice that when $p = \infty$ these functions v are polar quadratic polynomials as defined in [2, 19], that replace cones as the functions to which compare in the context of the non-homogeneous normalized infinity Laplacian.

Remark 5.8. Both operators Δ_p^N and Δ_∞^N are linear when applied to radial functions. This fact motivates using cusps and cones (as they are p -harmonic away from the vertex) in Lemma 5.6 altogether with a quadratic perturbation.

Lemma 5.9. *Let Ω be a bounded domain, $x_0 \in \Omega$, $p > n$,*

$$A > B \text{diam}(\Omega)^{1+\alpha_p},$$

and $B > 0$. Then, the function

$$v(x) = h + \frac{A}{1-\alpha_p}|x-x_0|^{1-\alpha_p} - \frac{B}{2}|x-x_0|^2$$

satisfies $\nabla v(x) \neq 0$ for any $x \in \overline{\Omega} \setminus \{x_0\}$.

Proof: A direct computation shows that for any $x \in \overline{\Omega} \setminus \{x_0\}$ we have that

$$\begin{aligned} |\nabla v(x)| &= |x-x_0| \cdot |A|x-x_0|^{-1-\alpha_p} - B| \\ &\geq |x-x_0| \cdot (A \text{diam}(\Omega)^{-1-\alpha_p} - B) > 0 \end{aligned}$$

by our hypothesis on the size of A . \blacksquare

Lemma 5.10. *Let Ω be a bounded domain, $x_0 \in \Omega$, $p > n \geq 2$,*

$$A \geq (1-\alpha_p) \sup_{z \in \partial\Omega} \frac{|u(z) - u(x_0)|}{|z-x_0|^{1-\alpha_p}} + B \text{diam}(\Omega)^{1+\alpha_p}, \quad (37)$$

and $B = (1+\varepsilon)C_p\|f\|_{L^\infty(\Omega)}$ with $\varepsilon > 0$. Suppose that h is such that

$$v(x) = h + \frac{A}{1-\alpha_p}|x-x_0|^{1-\alpha_p} - \frac{B}{2}|x-x_0|^2$$

touches u from above at some point $\tilde{x} \in \overline{\Omega}$. Then, necessarily $\tilde{x} \equiv x_0$ and $h = u(x_0)$.

We arrive at the same conclusion if we suppose instead that h is such that

$$v(x) = h - \frac{A}{1 - \alpha_p} |x - x_0|^{1 - \alpha_p} + \frac{B}{2} |x - x_0|^2$$

touches u from below at some point $\tilde{x} \in \overline{\Omega}$.

Proof: We provide the proof in the first case as the second one follows in the same way.

1. First we are going to prove that the contact point $\tilde{x} \notin \partial\Omega$. Assume for contradiction that $\tilde{x} \in \partial\Omega$. From the contact condition, we get that

$$h = u(\tilde{x}) - \frac{A}{1 - \alpha_p} |\tilde{x} - x_0|^{1 - \alpha_p} + \frac{B}{2} |\tilde{x} - x_0|^2,$$

and $u(x_0) < v(x_0)$. This two facts together yield,

$$u(x_0) < u(\tilde{x}) - \frac{A}{1 - \alpha_p} |\tilde{x} - x_0|^{1 - \alpha_p} + \frac{B}{2} |\tilde{x} - x_0|^2.$$

Rearranging terms we get,

$$A < (1 - \alpha_p) \frac{(u(\tilde{x}) - u(x_0))}{|\tilde{x} - x_0|^{1 - \alpha_p}} + \frac{B}{2} (1 - \alpha_p) |\tilde{x} - x_0|^{1 + \alpha_p}.$$

Notice that,

$$\frac{B}{2} (1 - \alpha_p) |\tilde{x} - x_0|^{1 + \alpha_p} < B \cdot \text{diam}(\Omega)^{1 + \alpha_p},$$

so we arrive at a contradiction with the definition of A and hence $\tilde{x} \notin \partial\Omega$.

2. Now, assume for contradiction that $\tilde{x} \neq x_0$. We have that v touches u from above at \tilde{x} . From the previous step we know that \tilde{x} must be an interior point; hence from the hypothesis of contradiction we know that v is \mathcal{C}^2 in a neighborhood of \tilde{x} . As u solves (33) in the viscosity sense, we can use v as a test function in the definition of viscosity solution and get

$$-\Delta_p^N v(\tilde{x}) \leq f(\tilde{x}).$$

On the other hand, Lemma 5.9 implies $\nabla v \neq 0$ and then, from Lemma 5.6,

$$(1 + \varepsilon) \|f\|_{L^\infty(\Omega)} = C_p^{-1} B = -\Delta_p^N v(\tilde{x}),$$

with $\varepsilon > 0$, a contradiction. ■

We now complete the proof of Theorem 5.1.

Proof of Theorem 5.1: Consider a cusp pointing downwards centered at x ,

$$v(y) = h + \frac{A}{1 - \alpha_p} |y - x|^{1 - \alpha_p} - \frac{B}{2} |y - x|^2$$

with

$$A = (1 - \alpha_p) \sup_{z \in \partial\Omega} \frac{|u(z) - u(x)|}{|z - x|^{1 - \alpha_p}} + B \operatorname{diam}(\Omega)^{1 + \alpha_p}$$

and $B = (1 + \varepsilon) C_p \|f\|_{L^\infty(\Omega)}$ with $\varepsilon > 0$. Lemma 5.10 says that if v touches the graph of u from above, then the contact point is x and $h = u(x)$. We deduce,

$$\begin{aligned} u(y) &\leq v(y) = u(x) + \frac{A}{1 - \alpha_p} |y - x|^{1 - \alpha_p} - \frac{B}{2} |y - x|^2 \\ &\leq u(x) + \frac{A}{1 - \alpha_p} |y - x|^{1 - \alpha_p}. \end{aligned}$$

We get,

$$\begin{aligned} \frac{u(y) - u(x)}{|y - x|^{1 - \alpha_p}} &\leq \sup_{z \in \partial\Omega} \frac{|u(z) - u(x)|}{|z - x|^{1 - \alpha_p}} \\ &\quad + (1 + \varepsilon) \|f\|_{L^\infty(\Omega)} \frac{C_p}{1 - \alpha_p} \operatorname{diam}(\Omega)^{1 + \alpha_p}. \end{aligned}$$

As this estimate holds for any $\varepsilon > 0$, we can let $\varepsilon \rightarrow 0$.

On the other hand, we can do the same with a cusp pointing upwards and slide it until it touches from below. Namely, we get

$$\begin{aligned} u(y) &\geq v(y) = u(x) - \frac{A}{1 - \alpha_p} |y - x|^{1 - \alpha_p} + \frac{B}{2} |y - x|^2 \\ &\geq u(x) - \frac{A}{1 - \alpha_p} |y - x|^{1 - \alpha_p} \end{aligned}$$

and then argue as before. The two estimates together yield the result. \blacksquare

Now, we turn to the proof of the global estimates for the Dirichlet problem in Theorem 5.4. A key point in the proof is the following lemma, similar to the comparison with cones property in [8].

Lemma 5.11. *Consider $A > B \operatorname{diam}(\Omega)^{1 + \alpha_p}$, $B = (1 + \varepsilon) C_p \|f\|_{L^\infty(\Omega)}$ with $\varepsilon > 0$, and $x_0 \in \overline{\Omega}$. Let u be a viscosity solution of (33). Then if*

$$u(x) \leq u(x_0) + A |x - x_0|^{1 - \alpha_p} - \frac{B}{2} |x - x_0|^2$$

for all x in $\partial\Omega$ the same inequality holds in the interior, that is

$$u(x) \leq u(x_0) + A|x - x_0|^{1-\alpha_p} - \frac{B}{2}|x - x_0|^2$$

for all x in Ω . In the same way, if

$$u(x) \geq u(x_0) - A|x - x_0|^{1-\alpha_p} + \frac{B}{2}|x - x_0|^2$$

for all x in $\partial\Omega$ the same inequality holds in the interior, that is

$$u(x) \geq u(x_0) - A|x - x_0|^{1-\alpha_p} + \frac{B}{2}|x - x_0|^2$$

for all x in Ω .

Proof: We prove just the first claim since the other one is analogous. Denote as usual

$$v(x) = u(x_0) + A|x - x_0|^{1-\alpha_p} - \frac{B}{2}|x - x_0|^2.$$

As we intend to prove that $u(x) - v(x) \leq 0$ for all $x \in \Omega$, assume for the sake of contradiction that $u(\tilde{x}) - v(\tilde{x}) = \max_{\Omega}(u - v) > 0$ as we would be done otherwise. Since $u - v \leq 0$ on $\partial\Omega$ by hypothesis, \tilde{x} must be an interior point. Moreover, $u(\tilde{x}) - v(\tilde{x}) > 0$ so $\tilde{x} \neq x_0$ and v is \mathcal{C}^2 in a neighborhood of \tilde{x} . As u is a viscosity solution of (33), we have by definition that

$$-\Delta_p^N v(\tilde{x}) \leq f(\tilde{x})$$

On the other hand, as $\nabla v \neq 0$ by the hypothesis on the size of A (see Lemma 5.9) we have by Lemma 5.6 that

$$-\Delta_p^N v(\tilde{x}) = C_p^{-1}B = (1 + \varepsilon) \|f\|_{L^\infty(\Omega)}$$

a contradiction, as $\varepsilon > 0$. ■

Proof of Theorem 5.4: Pick $x_0 \in \partial\Omega$. Since $g \in \mathcal{C}^{0,1-\alpha_p}(\partial\Omega)$, we have that for any $x \in \partial\Omega$

$$u(x) = g(x) \leq u(x_0) + L|x - x_0|^{1-\alpha_p}$$

for $L = [g]_{1-\alpha_p, \partial\Omega}$. Let

$$A \geq L + B \operatorname{diam}(\Omega)^{1+\alpha_p} \tag{38}$$

and $B = (1 + \varepsilon) C_p \|f\|_{L^\infty(\Omega)}$ with $\varepsilon > 0$.

It is easy to see that (38) implies,

$$u(x) \leq u(x_0) + L|x - x_0|^{1-\alpha_p} \leq u(x_0) + A|x - x_0|^{1-\alpha_p} - \frac{B}{2}|x - x_0|^2$$

for any $x \in \partial\Omega$. Then, Lemma 5.11 implies that the same holds for any $x \in \Omega$.

With the same proof it also holds that

$$u(x) \geq u(x_0) - A|x - x_0|^{1-\alpha_p} + \frac{B}{2}|x - x_0|^2 \quad (39)$$

for any $x \in \Omega$ and $x_0 \in \partial\Omega$.

Choose now $y \in \Omega$, by (39) we know that for any $x \in \partial\Omega$ we have

$$u(x) \leq u(y) + A|x - y|^{1-\alpha_p} - \frac{B}{2}|x - y|^2$$

and thanks to Lemma 5.11 the same holds also for any $x \in \Omega$. Reversing the role of x and y we obtain

$$|u(x) - u(y)| \leq A|x - y|^{1-\alpha_p} - \frac{B}{2}|x - y|^2 \leq A|x - y|^{1-\alpha_p}$$

and the result follows. ■

6. Limit equation

Theorem 6.1. *Let Ω be a bounded domain, $2 \leq n < p \leq \infty$, and u_p a viscosity solution of*

$$\begin{cases} -\Delta_p^N u_p(x) = f_p(x) & \text{in } \Omega \\ u_p(x) = g_p(x) & \text{on } \partial\Omega, \end{cases} \quad (40)$$

with $f_p \in \mathcal{C}(\Omega)$ and $g_p \in \mathcal{C}^{0, \frac{p-n}{p-1}}(\partial\Omega)$. Suppose that f_p and g_p converge uniformly to some $f \in \mathcal{C}(\Omega)$ and g (which in turn is $\mathcal{C}^{0,1}(\partial\Omega)$). Then, there exists a subsequence $u_{p'}$ that converge uniformly to some u , a viscosity solution of

$$\begin{cases} -\Delta_\infty^N u(x) = f(x) & \text{in } \Omega \\ u(x) = g(x) & \text{on } \partial\Omega. \end{cases} \quad (41)$$

Moreover, $u \in \mathcal{C}^{0,1}(\Omega)$.

As it was mentioned in the introduction, this result is related to the results in [4]. However, a distinct feature of this limit process is the lack of variational structure of problem (40) that yields complications in the proof of the uniform convergence of the solutions. Our ABP estimate provides a stable L^∞ bound that can be used in combination with our stable regularity results to prove convergence.

It is also interesting to understand the limits of (40) to solutions of (41) as both equations can be interpreted in the framework of game theory, see [22, 23].

Once uniform estimates have been established, the proof of Theorem 6.1 is rather standard, but we include it for the sake of completeness.

Proof: Fix p_0 such that $n < p_0$. Then, for $p \geq p_0$, Corollary 5.5 yields the following estimate

$$\begin{aligned} \|u_p\|_{\mathcal{C}^{0, \frac{p_0-n}{p_0-1}}(\Omega)} &\leq \text{diam}(\Omega)^{\left(\frac{p-n}{p-1} - \frac{p_0-n}{p_0-1}\right)} \cdot \|u_p\|_{\mathcal{C}^{0, \frac{p-n}{p-1}}(\Omega)} \\ &\leq \text{diam}(\Omega)^{\left(\frac{p-n}{p-1} - \frac{p_0-n}{p_0-1}\right)} \cdot \|g_p\|_{\mathcal{C}^{0, \frac{p-n}{p-1}}(\partial\Omega)} \\ &\quad + \text{diam}(\Omega)^{2 - \frac{p_0-n}{p_0-1}} \cdot \left(\frac{2p}{p-1} \text{diam}(\Omega)^{\frac{p-n}{p-1}} + \frac{p}{p+n-2} \right) \|f_p\|_{L^\infty(\Omega)}. \end{aligned}$$

As the right hand side can be bounded independently of p , Arzela-Ascoli Theorem yields the existence of a subsequence converging uniformly to some limit $u \in \mathcal{C}(\Omega)$. We will still denote the subsequence by u_p .

Now, we turn to checking that the limit u is a viscosity solution of (41). Let $x_0 \in \Omega$ and a function $\varphi \in \mathcal{C}^2(\Omega)$ such that $u - \varphi$ attains a local minimum at x_0 . Up to replacing $\varphi(x)$ with $\varphi(x) - |x - x_0|^4$, we can assume without loss of generality that minimum to be strict.

As u is the uniform limit of the subsequence u_p and x_0 is a strict minimum point, there exists a sequence of points $x_p \rightarrow x_0$ as $p \rightarrow \infty$ such that $(u_p - \varphi)(x_p)$ is a local minimum for each p in the sequence.

Assume first that $|\nabla\varphi(x_0)| > 0$. Then, $|\nabla\varphi(x_p)| > 0$ for p large enough and, as u_p is a viscosity supersolution of (40), we have that,

$$\begin{aligned} -\frac{1}{p} \text{trace} \left[\left(I + (p-2) \frac{\nabla\varphi(x_p) \otimes \nabla\varphi(x_p)}{|\nabla\varphi(x_p)|^2} \right) D^2\varphi(x_p) \right] &= -\Delta_p^N \varphi(x_p) \\ &\geq f_p(x_p). \end{aligned} \quad (42)$$

Letting $p \rightarrow \infty$ we get

$$-\left\langle D^2\varphi(x_0) \frac{\nabla\varphi(x_0)}{|\nabla\varphi(x_0)|}, \frac{\nabla\varphi(x_0)}{|\nabla\varphi(x_0)|} \right\rangle = -\Delta_\infty^N \varphi(x_0) \geq f(x_0).$$

If we assume otherwise that $\nabla\varphi(x_0) = 0$, we have to consider two cases. Suppose first that there exists a subsequence still indexed by p such that

$|\nabla\varphi(x_p)| > 0$ for all p in the subsequence. Then, by Definition 2.4, we can let $p \rightarrow \infty$ to get

$$\begin{aligned} -\liminf_{p \rightarrow \infty} \left\langle D^2\varphi(x_p) \frac{\nabla\varphi(x_p)}{|\nabla\varphi(x_p)|}, \frac{\nabla\varphi(x_p)}{|\nabla\varphi(x_p)|} \right\rangle &= -m(D^2\varphi(x_0)) \\ &= -\Delta_\infty^N \varphi(x_0) \geq f(x_0). \end{aligned}$$

If such a subsequence does not exist, according to Definition 2.2, we have that

$$-\frac{1}{p} \Delta\varphi(x_p) - \frac{(p-2)}{p} m(D^2\varphi(x_p)) = -\Delta_p^N \varphi(x_p) \geq f_p(x_p)$$

for every p large enough. Letting $p \rightarrow \infty$, we have

$$-m(D^2\varphi(x_0)) = -\Delta_\infty^N \varphi(x_0) \geq f(x_0).$$

We have proved (13), so u is a viscosity supersolution of (41). The subsolution case is similar.

Finally, we conclude that $u \in \mathcal{C}^{0,1}(\Omega)$ either letting $p_0 \rightarrow \infty$ or using that u is a solution of (41), and hence the estimates in Corollary 5.5 apply. \blacksquare

7. Examples

In this section we provide some examples. The first one shows that the classical ABP estimate in the form

$$\sup_{\Omega} u \leq \sup_{\partial\Omega} u + C(n, \Omega) \|\Delta_\infty^N u\|_{L^n(\Omega)}$$

fails to hold. In order to prove this we shall construct a sequence of functions $\{u_\varepsilon\}$ defined on $B_1(0)$, vanishing on the boundary such that

$$\sup_{B_1(0)} u_\varepsilon(x) \approx 1 \quad \text{and} \quad \|\Delta_\infty^N u_\varepsilon\|_{L^n(B_1(0))} \rightarrow 0,$$

as $\varepsilon \rightarrow 0$. To do this we define

$$u_\varepsilon(x) = v_\varepsilon(|x|) = \frac{1}{1+\varepsilon} (1 - |x|^{1+\varepsilon}),$$

so that,

$$\sup_{B_1(0)} u_\varepsilon = \frac{1}{1+\varepsilon}$$

and

$$-\Delta_\infty^N u_\varepsilon(x) = -v_\varepsilon''(|x|) = \frac{\varepsilon}{|x|^{1-\varepsilon}}. \tag{43}$$

Now a straightforward computation gives

$$\|\Delta_\infty^N u_\varepsilon\|_{L^n(B_1(0))}^n = c(n)\varepsilon^n \int_0^1 \frac{d\rho}{\rho^{1-n\varepsilon}} = c(n)\varepsilon^{n-1}.$$

Notice however that equation (43) is a pointwise computation and one has to give it a meaning in the viscosity sense. However this difficulty can be easily overcome considering, for instance, the functions

$$u_{\varepsilon,\delta}(|x|) = \frac{(1 + \delta^2)^{\frac{1+\varepsilon}{2}}}{1 + \varepsilon} - \frac{(|x|^2 + \delta^2)^{\frac{1+\varepsilon}{2}}}{1 + \varepsilon}$$

which are \mathcal{C}^2 , details are left to the reader.

As we said in the introduction we are not able to provide a counterexample to the validity of an estimate of the form

$$\sup_\Omega u \leq \sup_{\partial\Omega} u + C(n, p, \Omega) \|\Delta_\infty^N u\|_{L^p(\Omega)} \quad (44)$$

for $p > n$. Our strategy, which is to approximate the “infinity harmonic” cone $1 - |x|$ with radial functions, cannot be used to obtain such a counterexample. In fact, it is easy to see that if $v_\varepsilon(|x|)$ uniformly converge to $1 - |x|$, then the sequence of unidimensional functions $v_\varepsilon''(\rho)$ has to converge in the sense of distributions to $-2\delta_0$, twice a Dirac mass in the origin. Hence to find a counterexample to (44) with our strategy one essentially needs to construct a sequence $v_\varepsilon(\rho)$ such that

$$\int_0^1 v_\varepsilon''(\rho) d\rho \rightarrow 1 \quad \text{and} \quad \int_0^1 (v_\varepsilon''(\rho))^p \rho^{n-1} d\rho \rightarrow 0.$$

An application of Hölder inequality gives that this is impossible for $p > n$.

The second example we provide shows that our estimate is, in some sense, sharper than the plain L^∞ bound

$$\sup_\Omega u \leq \sup_{\partial\Omega} u + c(n, \Omega) \|\Delta_\infty^N u\|_{L^\infty(\Omega)}. \quad (45)$$

Let us consider the functions

$$w_\varepsilon(x) = \begin{cases} 1 - |x| & \text{if } \varepsilon < |x| \leq 1 \\ 1 - \frac{\varepsilon}{2} - \frac{|x|^2}{2\varepsilon} & \text{if } |x| \leq \varepsilon. \end{cases}$$

One immediately sees that

$$\sup_{B_1(0)} w_\varepsilon \approx 1 \quad \text{and} \quad \|\Delta_\infty^N w_\varepsilon\|_{L^\infty(B_1(0))} = \frac{1}{\varepsilon},$$

while

$$\int_0^{\sup_{B_1(0)} w_\varepsilon} \|\Delta_\infty^N w_\varepsilon\|_{L^\infty(\{w_\varepsilon=r\})} dr \approx 1,$$

so our estimate can give more information than (45).

Appendix A

For the sake of completeness we provide here the proof of some technical details needed in the foregoing.

Lemma A.1. *Let Ω^* be a convex domain and u a continuous semiconvex function such that $u \leq 0$ in $\mathbb{R}^n \setminus \Omega^*$ and $u(x_0) > 0$ for some $x_0 \in \Omega^*$. Let $\Gamma_\sigma(u^+)$ be the concave envelope of u^+ (extended by 0 outside Ω^*) in*

$$\Omega_\sigma^* = \{x \in \mathbb{R}^n \text{ such that } \text{dist}(x, \Omega^*) \leq \sigma\}.$$

Then $\Gamma_\sigma(u)$ is $C_{\text{loc}}^{1,1}$ in Ω_σ^ (and hence second order differentiable a.e.). Moreover,*

$$\{x \in \Omega^* : \det D^2 \Gamma_\sigma(u) \neq 0\} \subset C_\sigma^+(u),$$

where $C_\sigma^+(u)$ is the set of points in Ω_σ^ where $u = \Gamma_\sigma(u)$.*

Proof: This lemma can be easily deduced from the arguments of [6, Lemma 3.5]. However, since our statement is slightly different, we shall briefly sketch the proof for the sake of completeness. Obviously the second derivative of $\Gamma_\sigma(u)$ is bounded from above by 0 and hence to prove the $C_{\text{loc}}^{1,1}$ regularity it is enough to show they are locally bounded from below. Since the graph of u can be touched from below by a paraboloid with fixed opening this is clearly true for every point in the contact set $C_\sigma^+(u)$.

Let now choose a compact set $\Omega^* \subset K \subset \Omega_\sigma^*$, pick $x \in K \setminus C_\sigma^+(u)$ and let L be a supporting hyperplane at x to the graph of $\Gamma_\sigma(u^+)$. Following the same argument of [6, Lemma 3.5] one can prove that

$$x \in \text{conv}\{y \in \Omega_\sigma^* \text{ such that } u^+(y) = L(y)\},$$

where conv denotes the closed convex hull of a set. Moreover, by Caratheodory's Theorem, we can write

$$x = \lambda_1 x_1 + \cdots + \lambda_{n+1} x_{n+1}$$

with $x_i \in \{u^+ = L\}$, $\lambda_i \geq 0$ and $\sum_{i=1}^{n+1} \lambda_i = 1$. We conclude that x belongs to the simplex $\text{conv}\{x_1, \dots, x_{n+1}\}$ and that $L \equiv \Gamma_\sigma(u^+)$ in this set. Moreover, all the $x_i \in \Omega^* \cap C_\sigma^+(u)$ with the exception of at most one, that may belong

to $\partial\Omega_\sigma^*$; otherwise by concavity one would have that $\Gamma_\sigma(u^+) = 0$ everywhere in Ω_σ^* since $u^+ = 0$ in $\Omega_\sigma^* \setminus \Omega^*$.

We now claim that there exists a constant $C = C(\sigma, n, \Omega^*, K)$ and an index i_0 such that $x_{i_0} \in \Omega^*$ and $\lambda_{i_0} \geq C$, this is true with $C = 1/(n+1)$ if all the x_i belong to Ω^* . Otherwise we can assume without loss of generality that $x_{n+1} \in \partial\Omega_\sigma^*$ and suppose that $\lambda_i < C$ for $i = 1, \dots, n$. Then,

$$0 < \text{dist}(K, \partial\Omega_\sigma^*) \leq |x - x_{n+1}| \leq \sum_{i=1}^n \lambda_i |x_i - x_{n+1}| < n C (\text{diam}(\Omega^*) + 2\sigma),$$

a contradiction if C is small enough. Assume without loss of generality that $i_0 = 1$, by the semiconvexity assumption we know that

$$\Gamma_\sigma(u^+)(x_1 + h) \geq u^+(x_1 + h) \geq L(x_1 + h) - M|h|^2$$

for $|h| \leq \text{dist}(\Omega^*, \partial\Omega_\sigma^*)$ and L an affine function whose graph is the supporting hyperplane to the graph of $\Gamma_\sigma(u^+)$ at x_1 . Writing

$$x + h = \lambda_1 \left(x_1 + \frac{h}{\lambda_1} \right) + \dots + \lambda_{n+1} x_{n+1},$$

thanks to the concavity of $\Gamma_\sigma(u^+)$ and the fact that $L \equiv \Gamma_\sigma(u^+)$ in the simplex $\text{conv}\{x_1, \dots, x_{n+1}\}$, we get

$$\begin{aligned} \Gamma_\sigma(u^+)(x + h) &\geq \lambda_1 \Gamma_\sigma(u^+) \left(x_1 + \frac{h}{\lambda_1} \right) \\ &\quad + \lambda_2 \Gamma_\sigma(u^+)(x_2) + \dots + \lambda_{n+1} \Gamma_\sigma(u^+)(x_{n+1}) \\ &\geq \lambda_1 L \left(x_1 + \frac{h}{\lambda_1} \right) - \frac{M}{\lambda_1} |h|^2 + \lambda_2 L(x_2) + \dots + \lambda_{n+1} L(x_{n+1}) \\ &= L(x + h) - \frac{M}{\lambda_1} |h|^2 \geq L(x + h) - \tilde{M}(n, \sigma, \Omega^*, K) |h|^2 \end{aligned}$$

for $|h| \leq \varepsilon(n, \sigma, \Omega^*, K)$ small enough, which is the claimed $\mathcal{C}_{\text{loc}}^{1,1}$ regularity. To prove the last assertion in the statement of the Lemma, just notice that for any $x \in \Omega^* \setminus C_\sigma^+(u)$ the function $\Gamma_\sigma(u^+)$ coincides with an affine function on a segment. \blacksquare

Lemma A.2. *Let u be a viscosity subsolution of*

$$\mathcal{F}(\nabla u, D^2 u) = f(x) \quad \text{in } \Omega$$

and let $x_0 \in \Omega$. If φ is a function such that $u - \varphi$ has a local maximum at x_0 and φ is twice differentiable at x_0 , then

$$\mathcal{F}_*(\nabla\varphi(x_0), D^2\varphi(x_0)) \leq f(x_0).$$

Proof: By assumption

$$\varphi(x) = \varphi(x_0) + \langle \nabla\varphi(x_0), (x - x_0) \rangle + \frac{1}{2} \langle D^2\varphi(x_0)(x - x_0), (x - x_0) \rangle + o(|x - x_0|^2).$$

Thus for every $\varepsilon > 0$ the \mathcal{C}^2 function

$$\varphi_\varepsilon(x) = \varphi(x_0) + \langle \nabla\varphi(x_0), (x - x_0) \rangle + \frac{1}{2} \langle D^2\varphi(x_0)(x - x_0), (x - x_0) \rangle + \frac{\varepsilon}{2} |x - x_0|^2.$$

touches u from above at x_0 , so by definition of viscosity subsolution we have

$$\mathcal{F}_*(\nabla\varphi(x_0), D^2\varphi(x_0) + \varepsilon\text{Id}) \leq f(x_0)$$

and passing to the limit in ε we obtain the thesis. ■

Before stating the next Proposition, we recall that a Lipschitz function $v : M \rightarrow N$ between two \mathcal{C}^1 manifolds is differentiable \mathcal{H}^{n-1} -a.e. and we shall denote with $\nabla^M v$ its tangential gradient, that is, the linear operator between $T_x M$ and $T_x N$ defined by

$$\nabla^M v \cdot \xi := \lim_{t \rightarrow 0} \frac{f(x + t\xi) - f(x)}{t},$$

see for instance [12, Section 3.1].

Proposition A.3. *Let w be a concave $\mathcal{C}^{1,1}$ function in a ball $B_R(0)$ with $w = 0$ on $\partial B_R(0)$. Then,*

- (1) *For every level $r \in (0, \sup_{B_R(0)} w)$, $\nabla w \neq 0$ and $M_r := \{w = r\}$ is a $\mathcal{C}^{1,1}$ manifold.*
- (2) *For a.e. $r \in [0, \sup_{B_R(0)} w]$, \mathcal{H}^{n-1} -a.e. $x \in M_r$ is a point of twice differentiability for w .*

(3) Let $r \in (0, \sup_{B_R(0)} w)$, if $x \in M_r$ is a point of twice differentiability, then,

$$\begin{aligned}
-D^2w(x) &= |\nabla w(x)| \sum_{i=1}^{n-1} \kappa_i(x) \tau_i(x) \otimes \tau_i(x) \\
&\quad - \sum_{i=1}^{n-1} \partial_{\nu \tau_i}^2 w(x) (\tau_i(x) \otimes \nu(x) + \nu(x) \otimes \tau_i(x)) \\
&\quad - \Delta_\infty^N w(x) \nu(x) \otimes \nu(x),
\end{aligned} \tag{46}$$

where:

(a) $\nu(x)$ is the exterior normal to M_r at x , i.e.

$$\nu(x) = -\frac{\nabla w(x)}{|\nabla w(x)|}.$$

(b) $\tau_i(x)$ for $i = 1, \dots, n-1$ is an orthonormal basis of $T_x M_r$ that diagonalizes the Weingarten operator $\nabla^M \nu$ at x .

(c) $\kappa_i(x)$ are the eigenvalues of $\nabla^M \nu$ at x , which are the principal curvatures of M_r at x .

Proof: (1) As w is concave we have that $\nabla w(x) = 0$ if and only if x is such that $w(x) = \sup_{B_R(0)} w$. Then, by the Implicit Function theorem, for every $r \in (0, \sup_{B_R(0)} w)$ the level set M_r can be locally represented as a graph of a $\mathcal{C}^{1,1}$ function. Hence, up to a change of coordinates, we can suppose that there exists $f : \mathcal{U} \subset \mathbb{R}^{n-1} \rightarrow \mathbb{R}$ such that

$$w(x', f(x')) = r, \quad \text{and} \quad \partial_n w(x', f(x')) \geq c > 0 \quad \forall x' \in \mathcal{U},$$

where we denote $x = (x', x_n) \in \mathbb{R}^{n-1} \times \mathbb{R}$. In particular,

$$\nabla' f(x') = -\frac{\nabla' w(x', f(x'))}{\partial_n w(x', f(x'))}$$

is Lipschitz.

(2) Now, let D be the set of points of twice differentiability for w . Then, we know that $|B_R(0) \setminus D| = 0$ (see [11]) and by the Coarea Formula,

$$0 = \int_{B_R(0) \setminus D} |\nabla w(x)| dx = \int_0^{\sup_{B_R(0)} w} \mathcal{H}^{n-1}(M_r \cap (B_R(0) \setminus D)) dr.$$

(3) Let now x_0 be a point of twice differentiability for w . By part (1), we know that $\nabla w(x_0) \neq 0$. Without loss of generality we can assume $x_0 = 0$.

By means of an axis rotation, we can also suppose that $\nu(0) = e_n$, that is,

$$\nabla w(0) = \partial_n w(0, 0) e_n$$

with $\partial_n w(0, 0) < 0$.

There exists $f : B'_\delta(0) \subset \mathbb{R}^{n-1} \rightarrow \mathbb{R}$ such that $w(x', f(x')) = r$, in particular, $f(0) = 0$ and $\nabla' f(0) = 0$. Moreover,

$$\nabla' f(x') = -\frac{\nabla' w(x', f(x'))}{\partial_n w(x', f(x'))}$$

is differentiable at 0.

Using another rotation of axis affecting only the first $n - 1$ variables, we can suppose that the matrix $D_{x'}^2 f(0)$ is diagonal in our coordinates, namely,

$$D_{x'}^2 f(0) = \text{diag}(-\kappa_1, \dots, -\kappa_{n-1}). \quad (47)$$

Moreover, the Gauss map coincides with

$$x' \mapsto \frac{1}{\sqrt{1 + |\nabla' f(x')|^2}} (-\nabla' f(x'), 1),$$

so differentiating, we obtain

$$\nabla^M \nu(0) = -D_{x'}^2 f(0).$$

In particular, from this equality and (47), we deduce that $\tau_1(0), \dots, \tau_{n-1}(0)$ and e_1, \dots, e_{n-1} coincide.

Differentiating twice the expression $w(x', f(x')) = r$ with respect to x' we finally obtain,

$$\begin{aligned} D_{x'}^2 w(0, 0) &= -\partial_n w(0, 0) D_{x'}^2 f(0) \\ &= |\nabla w(0)| D_{x'}^2 f(0) = -|\nabla w(0)| \nabla^M \nu(0). \end{aligned}$$

Moreover, we also have $\partial_{\nu, \nu}^2 w(0) = \langle D^2 w(0) \nu(0), \nu(0) \rangle = \Delta_\infty^N w(0)$ and the thesis is proved. \blacksquare

We recall the following version of the Gauss-Bonnet Theorem, the proof of which easily follows from the Area Formula between rectifiable sets (see [12, Corollary 3.2.20]).

Theorem A.4. *Let $K \subset \mathbb{R}^n$ a $\mathcal{C}^{1,1}$ convex set and ν_K its outer normal. Then,*

$$\int_{\partial K} \prod_{i=1}^{n-1} \kappa_i(x) d\mathcal{H}^{n-1} = \int_{\partial K} \det(\nabla^{\partial K} \nu_K) d\mathcal{H}^{n-1} = \mathcal{H}^{n-1}(\partial B_1(0)). \quad (48)$$

Proof: By the Area Formula (see [12, Corollary 3.2.20]) we have

$$\begin{aligned} \int_{\partial K} \det(\nabla^{\partial K} \nu_K) d\mathcal{H}^{n-1} &= \int_{\partial B_1(0)} \deg(\nu_K, \partial K, \partial B_1(0)) d\mathcal{H}^{n-1} \\ &= \mathcal{H}^{n-1}(\partial B_1(0)), \end{aligned}$$

where $\deg(\nu_K, \partial K, \partial B_1(0))$ is the Brouwer degree of the Gauss map $\nu_K : \partial K \rightarrow \partial B_1(0)$ which can be easily seen to be 1. \blacksquare

Acknowledgements: This paper was started while A. Di Castro and G. De Philippis were visiting the Mathematics Department of the University of Texas at Austin for whose hospitality the authors are very grateful.

References

- [1] R. Argiolas, F. Charro, I. Peral; *On the Aleksandrov-Bakelĭman-Pucci Estimate for Some Elliptic and Parabolic Nonlinear Operators*, Archive for Rational Mechanics and Analysis, to appear (Doi: 10.1007/s00205-011-0434-y).
- [2] S. Armstrong, C. Smart; *A finite difference approach to the infinity Laplace equation and tug-of-war games* (preprint).
- [3] E. Beckenbach, R. Bellman; *Inequalities*, Ergebnisse der Mathematik und ihrer Grenzgebiete, N. F., Bd. 30 Springer-Verlag, Berlin-Gĕttingen-Heidelberg 1961.
- [4] T. Bhattacharya, E. DiBenedetto, J. Manfredi; *Limit as $p \rightarrow \infty$ of $\Delta_p u_p = f$ and related extremal problems*, Rend. Sem. Mat. Univ. Politec. Torino (1989), 15–68.
- [5] I. Birindelli, F. Demengel; *The Dirichlet problem for singular fully nonlinear operators*, Discrete Contin. Dyn. Syst. (2007) Dynamical Systems and Differential Equations. Proceedings of the 6th AIMS International Conference, suppl., 110–121.
- [6] X. Cabré, L.A. Caffarelli; *Fully Nonlinear Elliptic Equations*, Amer. Math. Soc., Colloquium publications, Vol. 43 (1995).
- [7] L.A. Caffarelli, M.G. Crandall, M. Kocan, A. Świeĭch; *On viscosity solutions of fully nonlinear equations with measurable ingredients*, Comm. Pure Appl. Math., **49** (1996), 365–397.
- [8] M. G. Crandall, L. C. Evans, R. F. Gariepy; *Optimal Lipschitz extensions and the infinity Laplacian*, Calc. Var. Partial Differential Equations **13** (2001), no. 2, 123–139.
- [9] M. G. Crandall, H. Ishii, P. L. Lions; *User’s Guide to Viscosity Solutions of Second Order Partial Differential Equations*, Bull. Amer. Math. Soc. **27** (1992), no. 1, 1–67.
- [10] G. Dávila, P. Felmer and A. Quaas; *Alexandroff-Bakelman-Pucci estimate for singular or degenerate fully nonlinear elliptic equations*, C.R.Math. Acad. Sci. Paris **347** (2009), no.19-20, 1165–1168.
- [11] L.C. Evans, R. F. Gariepy; *Measure theory and fine properties of functions*, Studies in Advanced Mathematics. CRC Press, Boca Raton, FL, 1992. viii+268 pp.
- [12] H. Federer; *Geometric measure theory*, Die Grundlehren der mathematischen Wissenschaften, Band 153 Springer-Verlag New York Inc., New York 1969 xiv+676 pp.
- [13] Y. Giga; *Surface evolution equations. A level set approach*, Monographs in Mathematics, 99, Birkhäuser Verlag, Basel, 2006.
- [14] D. Gilbarg, N. S. Trudinger; *Elliptic Partial Differential Equations of Second Order*, Springer-Verlag, New York (1983).

- [15] C. Gutierrez; *The Monge-Ampère equation*, Birkhäuser, 2001.
- [16] C. Imbert; *Alexandroff-Bakelman-Pucci estimate and Harnack inequality for degenerate/singular fully non-linear elliptic equations*, J. Differential Equations **250** (2011), no. 3, 1553–1574.
- [17] R. Jensen; *Uniqueness of Lipschitz extensions: Minimizing the sup norm of the gradient*, Arch. Rational Mech. Anal. **123** (1993), 51–74.
- [18] B. Kawohl, Juan J. Manfredi, M. Parviainen; *Solutions of nonlinear PDEs in the sense of averages*, preprint.
- [19] G. Lu, P. Wang; *A PDE perspective of the normalized infinity Laplacian*, Comm. Partial Differential Equations **33** (2008), no. 10-12, 1788–1817.
- [20] G. Lu, P. Wang; *Inhomogeneous infinity Laplace equation* Advances in Mathematics, **217**(4) (2008), 1838–1868.
- [21] G. Lu, P. Wang; *Infinity Laplace equation with non-trivial right-hand side*, Electron. J. Differential Equations **77** (2010).
- [22] Y. Peres, O. Schramm, S. Sheffield, D. Wilson; *Tug-of-war and the infinity Laplacian*, J. Amer. Math. Soc. **22** (2009), no. 1, 167–210.
- [23] Y. Peres, S. Sheffield; *Tug-of-war with noise: a game theoretic view of the p -Laplacian*, Duke Math. J. Volume **145** (2008), 91–120.

FERNANDO CHARRO

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF TEXAS, 1 UNIVERSITY STATION C1200,
AUSTIN, TX 78712

E-mail address: fcharro@math.utexas.edu

GUIDO DE PHILIPPIS

SCUOLA NORMALE SUPERIORE, P.ZA DEI CAVALIERI 7, I-56126 PISA, ITALY

E-mail address: guido.dephilippis@sns.it

AGNESE DI CASTRO,

CMUC, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF COIMBRA, 3001-454 COIMBRA, PORTUGAL

E-mail address: dicastro@mat.uc.pt

DAVI MÁXIMO

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF TEXAS, 1 UNIVERSITY STATION C1200,
AUSTIN, TX 78712

E-mail address: maximo@math.utexas.edu