

THE CUBOID LEMMA AND MAL'TSEV CATEGORIES

MARINO GRAN AND DIANA RODELO

Dedicated to George Janelidze on the occasion of his sixtieth birthday

ABSTRACT: We prove that a regular category \mathbb{C} is a Mal'tsev category if and only if a strong form of the denormalised 3×3 Lemma holds true in \mathbb{C} . In this version of the 3×3 Lemma, the vertical exact forks are replaced by pullbacks of regular epimorphisms along arbitrary morphisms. The shape of the diagram it determines suggests to call it the Cuboid Lemma. This new characterisation of regular categories that are Mal'tsev categories (= 2-permutable) is similar to the one previously obtained for Goursat categories (= 3-permutable). We also analyse the "relative" version of the Cuboid Lemma and extend our results to that context.

KEYWORDS: Mal'tsev category, regular pushout, calculus of relations, homological diagram lemma, relative regular category.

AMS SUBJECT CLASSIFICATION (2000): 18C05, 08C05, 18B10, 18E10.

Introduction

One of the motivations to introduce the notion of a *Mal'tsev category*, already mentioned in the classical article by A. Carboni, J. Lambek and M.C. Pedicchio [5], was the possibility of establishing extended versions of homological diagram lemmas, such as the Snake Lemma, in a non-pointed context. This idea was further pursued in the work of D. Bourn [3], who established an extended version of the 3×3 Lemma in any regular Mal'tsev category, called the *denormalised 3×3 Lemma*. In general, in order to formulate this kind of results in a non-pointed context, the short exact sequences are simply replaced by *exact forks*, i.e. by diagrams of the form

$$R_f \begin{array}{c} \xrightarrow{f_1} \\ \xrightarrow{f_2} \end{array} A \xrightarrow{f} B$$

where f is a regular epimorphism, and (R_f, f_1, f_2) is the kernel pair of f .

Received September 17, 2012.

The first author was supported by the F.N.R.S. grant *Crédit aux chercheurs* 1.5.016.10F and by the Université catholique de Louvain.

The second author was supported by CMUC/FCT (Portugal) and the FCT Grant PTDC/MAT/120222/2010 through the European program COMPETE/FEDER.

The validity of the denormalised 3×3 Lemma is actually equivalent to the property of 3-permutability of the composition of equivalence relations [17, 10]: for all equivalence relations R and S on the same object one has that

$$RSR = SRS.$$

Those regular categories having 3-permutable equivalence relations are called *Goursat categories*, and they were introduced in [4]. However, the regular categories having the Mal'tsev property are precisely those that satisfy the stronger property of 2-permutability of the composition of equivalence relations, namely

$$RS = SR.$$

These observations naturally led to the problem of determining whether there existed a homological diagram lemma whose validity would characterise Mal'tsev categories among regular ones, in the same way as the denormalised 3×3 Lemma characterises Goursat categories. The present article answers this question: the *Cuboid Lemma* is a new diagram lemma, stronger than the denormalised 3×3 Lemma, allowing one to obtain the desired characterisation of Mal'tsev categories (Theorems 3.2 and 3.3). This result is obtained by using the “calculus of relations” available in any regular category, much in the spirit of the above mentioned article [5].

Our new observations can also be extended from the “absolute” context of regular categories to the wider context of “relative” regular categories [13, 14, 7]. Here, the role of regular epimorphisms (for regular categories) is played by morphisms that belong to a distinguished class \mathcal{E} of regular epimorphisms. Relative regular categories give a suitable setting to develop a “calculus of \mathcal{E} -relations”: this makes it possible to extend the main results obtained in a regular category to the “relative” context, by using exactly the same proofs.

Structure of the paper

In Section 1 we give the main background concerning regular categories and the calculus of relations. We revise some known properties of regular Mal'tsev categories and their characterisation through the so-called regular pushouts in Section 2. We also include a new characterisation of Mal'tsev categories through a suitable stability property of regular epimorphisms. In Section 3 we state the Cuboid Lemma and prove that it characterises regular categories that are Mal'tsev categories. We conclude the article by extending the main results of this paper to a relative context in Section 4.

1. Preliminaries

A finitely complete category \mathbb{C} is said to be a *regular* category when any kernel pair has a coequaliser and, moreover, regular epimorphisms are stable under pullbacks. In a regular category any morphism $f: A \rightarrow B$ has a factorisation $f = m \cdot p$, where p is a regular epimorphism and m is a monomorphism. The corresponding (regular epimorphism, monomorphism) factorisation system is then stable under pullbacks.

In this article \mathbb{C} will always denote a finitely complete regular category.

A relation R from A to B is a subobject $\langle r_1, r_2 \rangle: R \rightarrow A \times B$. The opposite relation, denoted R° , is the relation from B to A given by the subobject $\langle r_2, r_1 \rangle: R \rightarrow B \times A$. We identify a morphism $f: A \rightarrow B$ with the relation $\langle 1_A, f \rangle: A \rightarrow A \times B$ and write f° for the opposite relation. Given another relation S from B to C , the composite relation of R and S is a relation SR from A to C . With the above notation, we can write any relation $\langle r_1, r_2 \rangle: R \rightarrow A \times B$ as $R = r_2 r_1^\circ$. The following properties are well known (see [4], for instance); we collect them in a lemma for future references.

Lemma 1.1. *Let $f: A \rightarrow B$ be any morphism in a regular category \mathbb{C} . Then:*

- (1) $f f^\circ f = f$ and $f^\circ f f^\circ = f^\circ$;
- (2) $f f^\circ = 1_B$ if and only if f is a regular epimorphism.

A relation R from an object A to A is called a *relation on A* . Such a relation is *reflexive* if $1_A \leq R$, *symmetric* if $R^\circ \leq R$, and *transitive* when $RR \leq R$. As usual, a relation R on A is an *equivalence relation* when it is reflexive, symmetric and transitive. In particular, a kernel pair $\langle f_1, f_2 \rangle: R_f \rightarrow A \times A$ of a morphism $f: A \rightarrow B$ is an *effective equivalence relation*, which can be written either as $R_f = f^\circ f$, or as $R_f = f_2 f_1^\circ$, as mentioned above. When f is a regular epimorphism, then f is the coequaliser of f_1 and f_2 and the diagram

$$R_f \begin{array}{c} \xrightarrow{f_1} \\ \xrightarrow{f_2} \end{array} A \xrightarrow{f} B$$

is called an *exact fork*. Note that, if $f = m \cdot p$ is the (regular epimorphism, monomorphism) factorisation of an arbitrary morphism f , then $R_f = R_p$, so that an effective equivalence relation is always the kernel pair of a regular epimorphism.

2. Regular Mal'tsev categories

Recall that a finitely complete category \mathbb{C} is called a *Mal'tsev category* when any reflexive relation in \mathbb{C} is an equivalence relation [5, 4]. The following well known characterisation of regular Mal'tsev categories will be useful.

Proposition 2.1. *A regular category \mathbb{C} is a Mal'tsev category if and only if the composition of effective equivalence relations on any object in \mathbb{C} is commutative:*

$$R_f R_g = R_g R_f$$

for any pair of regular epimorphisms f and g in \mathbb{C} with the same domain.

Examples 2.2. A variety of universal algebras is a Mal'tsev category if and only if its theory has a ternary operation $p(x, y, z)$ satisfying the identities $p(x, y, y) = x$ and $p(x, x, y) = y$ [18]. Of course, the variety Grp of groups is a Mal'tsev category, as is any variety whose theory contains a binary operation satisfying the group identities. Also the variety of quasi-groups and the variety of Heyting algebras are Mal'tsev categories, as is the dual category of an elementary topos. The category of topological groups is a regular Mal'tsev category, as is the category of Hausdorff groups [4]. If \mathbb{C} is a finitely complete category, then the category $\text{Grp}(\mathbb{C})$ of internal groups in \mathbb{C} is a Mal'tsev category.

There are many known characterisations of Mal'tsev categories (see [2], for instance, and references therein). We shall now focus on the relationship with commutative diagrams of the form

$$\begin{array}{ccc} C & \xrightarrow{c} & A \\ g \downarrow \uparrow t & & f \downarrow \uparrow s \\ D & \xrightarrow{d} & B, \end{array} \tag{1}$$

where f and g are split epimorphisms ($f \cdot s = 1_B$, $g \cdot t = 1_D$), $f \cdot c = d \cdot g$, $s \cdot d = c \cdot t$, and c and d are regular epimorphisms. A diagram as in (1) is always a pushout; it is called a *regular pushout* [3] (alternatively, a *double extension* as in [11], that was inspired by [12]) when, moreover, the canonical morphism $\langle g, c \rangle: C \rightarrow D \times_B A$ to the pullback $D \times_B A$ of d and f is a regular epimorphism.

Remark 2.3. The condition of being a regular pushout can be expressed in terms of the calculus of relations: a commutative diagram of type (1) is a

regular pushout if and only if $cg^\circ = f^\circ d$ or, equivalently, $gc^\circ = d^\circ f$. Observe also that the vertical morphisms g and f are split epimorphisms, so that they induce a split epimorphism from R_c to R_d . Consequently, the image of R_c along g is R_d , which can be written as: $g\langle R_c \rangle = R_d$, i.e. $gc^\circ cg^\circ = d^\circ d$.

The regular categories that are Mal'tsev categories can be characterised through regular pushouts, as it follows from the results of D. Bourn in [3]. For the reader's convenience we give a new simple proof of this fact that uses the calculus of relations, and is also suitable to be extended to the "relative context" (see the last section).

Proposition 2.4. *A regular category \mathbb{C} is a Mal'tsev category if and only if any commutative diagram of the form (1) is a regular pushout; equivalently:*

$$f^\circ d = cg^\circ.$$

Proof: Suppose that \mathbb{C} is a regular Mal'tsev category. Then:

$$\begin{aligned} f^\circ d &= cc^\circ f^\circ d && (c \text{ is a regular epimorphism; Lemma 1.1(2)}) \\ &= cg^\circ d^\circ d && (f \cdot c = d \cdot g) \\ &= cg^\circ gc^\circ cg^\circ && (R_d = g\langle R_c \rangle; \text{ see Remark 2.3}) \\ &= cc^\circ cg^\circ gg^\circ && (R_g R_c = R_c R_g, \text{ by Proposition 2.1}) \\ &= cg^\circ. && (\text{Lemma 1.1(1)}) \end{aligned}$$

Conversely, let us consider regular epimorphisms $f: X \twoheadrightarrow Y$ and $g: X \twoheadrightarrow Z$. We want to prove that $R_f R_g = R_g R_f$. For this we build the following diagram

$$\begin{array}{ccc} R_f & \xrightarrow{\gamma} & g\langle R_f \rangle \\ f_1 \downarrow & & \downarrow r_1 \\ X & \xrightarrow{g} & Z \\ f \downarrow & & \\ Y & & \end{array}$$

that represents the regular image $g\langle R_f \rangle$ of R_f along g . It is easy to see that $g\langle R_f \rangle$ is a reflexive relation, so that we get two commutative diagrams of type (1). By assumption one then has the equalities: (A) $f_2 \gamma^\circ = g^\circ r_2$ and

(B) $\gamma f_1^\circ = r_1^\circ g$. Accordingly:

$$\begin{aligned}
R_f R_g &= f_2 f_1^\circ g^\circ g \\
&= f_2 \gamma^\circ r_1^\circ g \quad (g \cdot f_1 = r_1 \cdot \gamma) \\
&= g^\circ r_2 r_1^\circ g \quad (\text{A}) \\
&= g^\circ r_2 \gamma f_1^\circ \quad (\text{B}) \\
&= g^\circ g f_2 f_1^\circ \quad (r_2 \cdot \gamma = g \cdot f_2) \\
&= R_g R_f.
\end{aligned}$$

■

As first observed by D. Bourn [3], regular Mal'tsev categories have a strong stability property for regular epimorphisms. Again, we give a new simple proof here to make the paper self-contained.

Lemma 2.5. *Let \mathbb{C} be a regular Mal'tsev category. Consider a commutative diagram*

$$\begin{array}{ccc}
W \times_D C & \xrightarrow{\quad v \quad} & Y \times_B A \\
\begin{array}{c} \uparrow \\ k \\ \downarrow \\ j \end{array} & \searrow \gamma & \begin{array}{c} \uparrow \wedge \\ h \parallel i \\ \parallel \\ \parallel \\ \vee \\ \downarrow \end{array} & \xrightarrow{\quad c \quad} & A \\
W & \xrightarrow{\quad \delta \quad} & D & \xrightarrow{\quad d \quad} & B \\
\begin{array}{c} \uparrow \\ g \\ \downarrow \\ t \end{array} & \searrow w & Y & \xrightarrow{\quad \beta \quad} & B \\
\end{array} \quad (2)$$

where the front square is of the form (1), $\beta \cdot w = d \cdot \delta$, w is a regular epimorphism, $(W \times_D C, k, \gamma)$ and $(Y \times_B A, h, \alpha)$ are pullbacks. Then the comparison morphism $v: W \times_D C \rightarrow Y \times_B A$ is also a regular epimorphism.

Proof: The exterior rectangle in the commutative diagram

$$\begin{array}{ccccc}
W \times_D C & \xrightarrow{\quad \gamma \quad} & C & \xrightarrow{\quad c \quad} & A \\
\begin{array}{c} \uparrow \\ k \\ \downarrow \end{array} & & \begin{array}{c} \uparrow \\ g \\ \downarrow \end{array} & & \begin{array}{c} \uparrow \\ f \\ \downarrow \end{array} \\
W & \xrightarrow{\quad \delta \quad} & D & \xrightarrow{\quad d \quad} & B
\end{array}$$

is such that the comparison arrow $\langle k, c \cdot \gamma \rangle: W \times_D C \rightarrow W \times_B A$ to the pullback of $d \cdot \delta$ along f is a regular epimorphism, as a “composite” of a pullback with a regular pushout. Accordingly, the exterior rectangle in the

commutative diagram

$$\begin{array}{ccccc}
 W \times_D C & \xrightarrow{v} & Y \times_B A & \xrightarrow{\alpha} & A \\
 k \downarrow \uparrow & & h \downarrow \uparrow & & f \downarrow \uparrow \\
 W & \xrightarrow{w} & Y & \xrightarrow{\beta} & B.
 \end{array}$$

has the property that $\langle k, \alpha \cdot v \rangle: W \times_D C \rightarrow W \times_B A$ is a regular epimorphism. From the fact that the right hand square is a pullback and that w is a regular epimorphism it then easily follows that the arrow v is a regular epimorphism, as desired. ■

We now show that the property considered in Lemma 2.5 actually characterises the regular categories which are Mal'tsev categories.

Proposition 2.6. *Let \mathbb{C} be a regular category. The following conditions are equivalent:*

- (a) \mathbb{C} is a Mal'tsev category;
- (b) for any commutative cube (2), the comparison morphism $v: W \times_D C \rightarrow Y \times_B A$ is a regular epimorphism.

Proof: (a) \Rightarrow (b) by Lemma 2.5.

(b) \Rightarrow (a) Consider a commutative diagram of the type (1). It induces the commutative cube

$$\begin{array}{ccccc}
 C & \xrightarrow{\quad v \quad} & D \times_B A & & \\
 \uparrow \parallel & \searrow & \uparrow \parallel & \searrow \alpha & \\
 g \downarrow \uparrow & & h \parallel i & & A \\
 \parallel & & \parallel & & \parallel \\
 C & \xrightarrow{c} & D & & \\
 \parallel & & \parallel & & \parallel \\
 D & \xrightarrow{g} & D & \xrightarrow{d} & B, \\
 \parallel & & \parallel & & \parallel \\
 g \downarrow \uparrow & & t \downarrow \uparrow & & f \downarrow \uparrow \\
 D & \xrightarrow{d} & D & \xrightarrow{d} & B,
 \end{array}$$

where the right hand face is the pullback of d and f . By assumption, $v = \langle g, c \rangle$ is a regular epimorphism, and then \mathbb{C} is a Mal'tsev category by Proposition 2.4. ■

3. The Cuboid Lemma

In this section we use the characterisation of Mal'tsev categories given in Proposition 2.6 to explore a stronger form of the (denormalised) 3×3 Lemma [3, 17]. This follows the same line of research as in [10], where the so-called *Goursat pushouts* were used to show that a regular category is a Goursat category if and only if the 3×3 Lemma holds.

Recall that the 3×3 Lemma states the following: given a commutative diagram in \mathbb{C} of the form

$$\begin{array}{ccccc}
 R_{\bar{g}} & \xrightarrow{t_1} & R_g & \xrightarrow{v} & R_f \\
 \bar{g}_1 \downarrow & & g_1 \downarrow & & f_1 \downarrow \\
 R_c & \xrightarrow{c_1} & A & \xrightarrow{c} & C \\
 \bar{g} \downarrow & & g \downarrow & & f \downarrow \\
 S & \xrightarrow{s_1} & B & \xrightarrow{d} & D
 \end{array} \quad (3)$$

where the columns are exact (forks), as well as the middle row, then the upper row is exact if and only if the lower row is exact.

Following the analogous terminology introduced in [16], the 3×3 Lemma can be decomposed into two weaker formulations: the *Upper 3×3 Lemma*, stating that the exactness of the lower row implies the exactness of the upper row, and the *Lower 3×3 Lemma*, stating that the exactness of the upper row implies the exactness of the lower row. It was shown in [10] (Proposition 1), that these two (apparently) weaker formulations are equivalent to each other, and they are also equivalent to the 3×3 Lemma: all these formulations provide characterisations of regular Goursat categories (we refer the interested reader also to [8] for further developments).

By replacing the kernel pairs in the three exact columns of (3) with pull-backs of split epimorphisms along arbitrary morphisms, we get a “three dimensional version” of the 3×3 Lemma. Since these vertical exact forks are replaced by squares, horizontally we should now take 4 “forks” (instead of 3), two of which are then required to be exact forks. This gives rise to a diagram whose external part is a *cuboid*, and this explains the terminology adopted for the following property.

The Split Cuboid Lemma

Let \mathbb{C} be a regular category. Consider a commutative diagram in \mathbb{C}

$$\begin{array}{ccccc}
 & T & \xrightarrow{t_1} & V & \xrightarrow{v} & X \\
 \bar{k} \swarrow & & \xrightarrow{t_2} & & & \searrow h \\
 R_w & \xrightarrow{\bar{j}} & & W & \xrightarrow{w} & Y \\
 \delta \downarrow & & \xrightarrow{w_1} & & & \nearrow i \\
 & & \xrightarrow{w_2} & & & \\
 & R_c & \xrightarrow{c_1} & C & \xrightarrow{c} & A \\
 \bar{g} \swarrow & & \xrightarrow{c_2} & & & \searrow \beta \\
 S & \xrightarrow{\bar{t}} & & D & \xrightarrow{d} & B \\
 & & \xrightarrow{s_1} & & & \nearrow f \\
 & & \xrightarrow{s_2} & & & \\
 & & & & & \nearrow s
 \end{array}
 \tag{4}$$

where the three diamonds are pullbacks of split epimorphisms along arbitrary morphisms and the two middle rows are exact forks. Then *the upper row is exact if and only if the lower row is exact*.

Note that t_1, t_2, s_1 and s_2 are just parallel morphisms with no common splitting initially required. However, by the commutativity of the diagram, s_1 and s_2 do actually have a common splitting $\bar{g} \cdot \langle 1_C, 1_C \rangle \cdot t$. The commutativity of the diagram also implies that d is a regular epimorphism, that $d \cdot s_1 = d \cdot s_2$ since \bar{g} is an epimorphism and, moreover, that $v \cdot t_1 = v \cdot t_2$, because the pair of morphisms (h, α) is jointly monomorphic.

Remark 3.1. Observe that the existence of the split epimorphism \bar{g} implies that we always have $S = R_d$, i.e. the lower row is necessarily exact. Indeed, let $\phi: S \rightarrow R_d$ be the canonical arrow induced by the universal property of R_d thanks to the equality $d \cdot s_1 = d \cdot s_2$. This arrow ϕ is a monomorphism since S is a relation (the pair of arrows $(c_1 \cdot \bar{t}, c_2 \cdot \bar{t})$ is jointly monic by assumption). But ϕ is also a split epimorphism, since $\phi \cdot \bar{g}$ is a split epimorphism. Accordingly, the arrow $\phi: S \rightarrow R_d$ is an isomorphism. Consequently, the Split Cuboid Lemma is actually equivalent to the (apparently weaker) *Upper Split Cuboid Lemma*, asserting that *the upper row in diagram (4) is exact whenever the lower row is exact*.

We are now ready to state our main result:

Theorem 3.2. *Let \mathbb{C} be a regular category. The following conditions are equivalent:*

- (a) \mathbb{C} is a Mal'tsev category;
- (b) the Split Cuboid Lemma holds true in \mathbb{C} ;
- (c) the Upper Split Cuboid Lemma holds true in \mathbb{C} .

Proof: (a) \Rightarrow (b) From the exactness of the lower row in diagram (4), one clearly has that $T = R_v$, since $S = R_d$ and “kernel pairs commute with pullbacks”. Consequently, to prove the exactness of the upper row, it suffices to show that v is a regular epimorphism. The right cube of diagram (4) is of the form (2), and v is then a regular epimorphism by Proposition 2.6. This shows that the upper row is exact as well. Conversely, the fact that the lower row is always exact has been proved in Remark 3.1.

(b) \Rightarrow (c) Trivial.

(c) \Rightarrow (a) Consider a diagram of the form (2) and take the kernel pairs of c, d and w . We get an induced split epimorphism

$$\bar{g}: R_c \rightarrow R_d,$$

with splitting \bar{t} , and an induced morphism

$$\bar{\delta}: R_w \rightarrow R_d.$$

We obtain a diagram of the form (4) by defining $(T, \bar{k}, \bar{\gamma})$ as the pullback of \bar{g} and $\bar{\delta}$. Applying the Upper Split Cuboid Lemma to this diagram, we conclude that the upper row is exact and, consequently, v is a regular epimorphism. By Proposition 2.6, \mathbb{C} is a Mal'tsev category. ■

Since the condition of being a regular Mal'tsev category is known to be (strictly) stronger than the one of being a regular Goursat category, one would like to understand why the Split Cuboid Lemma is actually stronger than the (denormalised) 3×3 Lemma (see (3)). For this we now introduce a stronger form of the (Upper) Split Cuboid Lemma that will provide an immediate comparison with the 3×3 Lemma.

The Upper Cuboid Lemma

Let \mathbb{C} be a regular category. Consider a commutative diagram in \mathbb{C}

(b) \Rightarrow (a) This implication is obvious by Theorem 3.2 and the fact that the Upper Cuboid Lemma is stronger than the Upper Split Cuboid Lemma. ■

Remark 3.4. Of course, the validity of the Upper Cuboid Lemma is stronger than the one of the Upper 3×3 Lemma (which is itself equivalent to the 3×3 Lemma, as shown in [10]). Indeed, the Upper 3×3 Lemma is simply obtained as the special case of the Upper Cuboid Lemma where one takes twice the same regular epimorphism to build the pullbacks in diagram (5).

4. The relative context

In this section we briefly analyse the so-called “relative context” introduced by T. Janelidze [13, 14], and verify that our results easily extend from the “absolute context” considered in the previous sections to the relative one.

We begin by recalling the notion of a *relative regular category* as defined by J. Goedecke and T. Janelidze [7]. As in the previous sections, we shall always assume that the base category is finitely complete.

A *relative regular category* is a pair $(\mathbb{A}, \mathcal{E})$, where \mathbb{A} is a finitely complete category and \mathcal{E} is a class of regular epimorphisms in \mathbb{A} such that:

- (E1) \mathcal{E} contains all isomorphisms;
- (E2) the pullback of a morphism in \mathcal{E} belongs to \mathcal{E} ;
- (E3) \mathcal{E} is closed under composition;
- (E4) if f and $g \cdot f \in \mathcal{E}$, then $g \in \mathcal{E}$;
- (F) if a morphism factors as $f = e \cdot m$, with $e \in \mathcal{E}$ and m a monomorphism, then it also factors (essentially uniquely) as $f = m' \cdot e'$, where m' is a monomorphism and $e' \in \mathcal{E}$.

Given a finitely complete category \mathbb{A} that admits coequalisers of kernel pairs and \mathcal{E} the class of all regular epimorphisms in \mathbb{A} , then $(\mathbb{A}, \mathcal{E})$ is a relative regular category if and only if \mathbb{A} is a regular category: this context is usually referred to as the “absolute context”.

In the extension from the absolute to the relative context one replaces the regular epimorphisms with morphisms that belong to \mathcal{E} . As shown in [13] (see also [7]), relative regular categories provide the appropriate setting to develop a suitable calculus of \mathcal{E} -relations. Recall that an \mathcal{E} -relation R from A to B is a subobject $\langle r_1, r_2 \rangle: R \rightarrow A \times B$ such that $r_1, r_2 \in \mathcal{E}$. We can identify a morphism $f: A \rightarrow B$ in \mathcal{E} with the \mathcal{E} -relation $\langle 1_A, f \rangle: A \rightarrow A \times B$. The definition of the composition of \mathcal{E} -relations can be given exactly as in

the absolute case, and the properties gathered in Section 1 all extend to \mathcal{E} -relations for relative regular categories. However, for the relative version of Lemma 1.1 we require $f : A \rightarrow B$ to belong to \mathcal{E} : accordingly, the relative version of 1.1(2) should simply state that, for any $f \in \mathcal{E}$, one has the equality $ff^\circ = 1_B$ (since the arrows in \mathcal{E} are regular epimorphisms).

To extend our results to the relative case, we should consider:

- *equivalence \mathcal{E} -relations*: \mathcal{E} -relations which are equivalence relations;
- *\mathcal{E} -effective equivalence \mathcal{E} -relations*: \mathcal{E} -relations given by kernel pairs of morphisms in \mathcal{E} ;
- *\mathcal{E} -exact forks*: exact forks whose coequaliser belongs to \mathcal{E} .

The \mathcal{E} -image of an \mathcal{E} -relation R along a morphism $f \in \mathcal{E}$ is given by the monomorphism in the $(\mathcal{E}, \text{monomorphism})$ -factorisation of $(f \times f) \cdot \langle r_1, r_2 \rangle$

$$\begin{array}{ccc} R & \xrightarrow{\varphi} & S \\ \langle r_1, r_2 \rangle \downarrow & & \downarrow \langle s_1, s_2 \rangle \\ A \times A & \xrightarrow{f \times f} & B \times B, \end{array}$$

which exists by axiom (F) of the definition of a relative regular category; we write $f\langle R \rangle = S$. As in the absolute case, the relation $f\langle R \rangle$ is reflexive (or symmetric) whenever R is reflexive (or symmetric).

Let us then consider a diagram (of type (1))

$$\begin{array}{ccc} C & \xrightarrow{c} & A \\ g \updownarrow t & & f \updownarrow s \\ D & \xrightarrow{d} & B \end{array} \quad (6)$$

in a relative regular category, where f, c, g and d now belong to \mathcal{E} . It is called an \mathcal{E} -regular pushout when the induced morphism $\langle g, c \rangle : C \rightarrow D \times_B A$ also belongs to \mathcal{E} . In the context of a relative regular category $(\mathbb{A}, \mathcal{E})$ the property that any diagram (6) in \mathbb{A} is an \mathcal{E} -regular pushout will be called the \mathcal{E} -Mal'tsev axiom in what follows. This axiom has already been studied by T. Everaert, J. Goedecke and T. Van der Linden in [6], and is similar to the \mathcal{E} -Goursat axiom introduced in [7] by J. Goedecke and T. Janelidze.

Remark also that the notion of an \mathcal{E} -Mal'tsev category had essentially been introduced in T. Janelidze's PhD doctoral dissertation [13] (see Theorem 2.3.6 therein, for instance), even though the term \mathcal{E} -Mal'tsev had not been explicitly used there. As in the absolute case (Proposition 2.1), these

categories can be characterised by the commutativity of the composition of \mathcal{E} -effective equivalence \mathcal{E} -relations.

The statements from Remark 2.3 extend to the relative context by Theorem 2.10 of [14] and Lemmas 1.9 and 1.11 of [7]. We also obtain relative versions of Propositions 2.4 and 2.6 with the same proofs, which have been written on purpose in a way that can be now “understood” in the relative context:

Proposition 4.1. *A relative regular category $(\mathbb{A}, \mathcal{E})$ satisfies the \mathcal{E} -Mal'tsev axiom if and only if*

$$R_f R_g = R_g R_f$$

for any pair of arrows f and g in \mathcal{E} with the same domain.

Proposition 4.2. *Let $(\mathbb{A}, \mathcal{E})$ be a relative regular category. The following conditions are equivalent:*

- (a) *the \mathcal{E} -Mal'tsev axiom holds true in \mathbb{A} ;*
- (b) *$(\mathbb{A}, \mathcal{E})$ has the property that for any commutative cube (2), with $f, g, c, d, w \in \mathcal{E}$, the induced morphism $v: W \times_D C \rightarrow Y \times_B A$ belongs to \mathcal{E} .*

The Relative Cuboid Lemmas

Let $(\mathbb{A}, \mathcal{E})$ be a relative regular category. Consider a commutative diagram (4) (resp. (5)) where the three diamonds are pullbacks of split (resp. regular) epimorphisms which belong to \mathcal{E} along arbitrary morphisms and the two middle rows are \mathcal{E} -exact. Then *the upper row is \mathcal{E} -exact if and only if (resp. whenever) the lower row is \mathcal{E} -exact.*

With the same proof as in the absolute case, we obtain:

Theorem 4.3. *Let $(\mathbb{A}, \mathcal{E})$ be a relative regular category. The following conditions are equivalent:*

- (a) *the \mathcal{E} -Mal'tsev axiom holds true in \mathbb{A} ;*
- (b) *the Relative (Upper) Split Cuboid Lemma holds true in \mathbb{A} ;*
- (c) *the Relative Upper Cuboid Lemma holds true in \mathbb{A} .*

References

- [1] M. Barr, Exact Categories, in: *Lecture Notes in Math.* 236, Springer (1971) 1-120.
- [2] F. Borceux and D. Bourn, *Mal'cev, protomodular, homological and semi-abelian categories*, Kluwer, 2004.
- [3] D. Bourn, *The denormalized 3×3 lemma*, J. Pure Appl. Algebra 177 (2003) 113-129.
- [4] A. Carboni, G.M. Kelly, and M.C. Pedicchio, *Some remarks on Maltsev and Goursat categories*, Appl. Cat. Struct. 1 (1993) 385-421.

- [5] A. Carboni, J. Lambek, and M.C. Pedicchio, *Diagram chasing in Mal'tsev categories*, J. Pure Appl. Alg. 69 (1991) 271-284.
- [6] T. Everaert, J. Goedecke and T. Van der Linden, *Resolutions, higher extensions and the relative Mal'tsev axiom*, J. Algebra 371 (2012), 132-155.
- [7] J. Goedecke and T. Janelidze, *Relative Goursat categories*, J. Pure Appl. Algebra 216 (2012) 1726-1733.
- [8] M. Gran, Z. Janelidze and D. Rodelo, *3×3 -Lemma for star-exact sequences*, Homology, Homotopy Appl., Vol. 14 (2) (2012), 1-22.
- [9] M. Gran, Z. Janelidze and A. Ursini, *A good theory of ideal in regular multi-pointed categories*, J. Pure Appl. Algebra 216 (2012) 1905-1919.
- [10] M. Gran and D. Rodelo, *A new characterisation of Goursat categories*, Appl. Categ. Structures Vol. 20 (2012) 229-238.
- [11] M. Gran and V. Rossi, *Galois theory and double central extensions*, Homology, Homotopy Appl., Vol. 6 (1) (2004) 283-298.
- [12] G. Janelidze, *What is a double central extension?* Cah. Top. Géom. Diff. Catég. XXXII No. 3 (1991) 191-201.
- [13] T. Janelidze, *Foundation of relative non-abelian homological algebra*, Ph.D. Thesis, University of Cape Town (2009)
- [14] T. Janelidze, *Relative semi-abelian categories*, Appl. Categ. Struct. 17, (2009) 373-386.
- [15] Z. Janelidze, *Subtractive categories*, Appl. Categ. Struct. 13, (2005) 343-350.
- [16] Z. Janelidze, *The pointed subobject functor, 3×3 lemmas and subtractivity of spans*, Th. Appl. Categ. 23 (2010) 221-242.
- [17] S. Lack, *The 3-by-3 lemma for regular Goursat categories*, Homology, Homotopy Appl., Vol. 6 (1) (2004) 1-3.
- [18] J.D.H. Smith, *Mal'tsev Varieties*, Lecture Notes in Math. 554 (1976)

MARINO GRAN

INSITUT DE RECHERCHE EN MATHÉMATIQUE ET PHYSIQUE, UNIVERSITÉ CATHOLIQUE DE LOUVAIN,
CHEMIN DU CYCLOTRON 2, 1348 LOUVAIN-LA-NEUVE, BELGIUM

E-mail address: `marino.gran@uclouvain.be`

DIANA RODELO

DEPARTAMENTO DE MATEMÁTICA, FACULDADE DE CIÊNCIAS E TECNOLOGIA, UNIVERSIDADE DO AL-
GARVE, CAMPUS DE GAMBELAS, 8005-139 FARO, PORTUGAL
CMUC, UNIVERSITY OF COIMBRA, 3001-501 COIMBRA, PORTUGAL

E-mail address: `drodelo@ualg.pt`