

A NOTE ON PSEUDOVARITIES OF COMPLETELY REGULAR SEMIGROUPS

JORGE ALMEIDA AND ALFREDO COSTA

ABSTRACT: This note shows that, even though a paper of Almeida and Trotter on completely regular semigroups makes use of free profinite semigroupoids over profinite graphs with infinitely many vertices, which has since been shown to be a structure that in general needs to be handled with great care, the required properties do hold, and thus the results stand.

KEYWORDS: Completely regular semigroups, pseudovarieties, free profinite semigroups, free profinite semigroupoids, profinite graphs.

AMS SUBJECT CLASSIFICATION (2010): primary 20M07; secondary 20M17, 20M05, 22A15.

1. Introduction

In [6] one finds a systematic study of relatively free profinite categories and relatively free profinite semigroupoids generated by profinite graphs with a finite number of vertices. The case of profinite graphs with an infinite number of vertices is more delicate, as highlighted in [3]. The main problem is that in the infinite-vertex case, the topological closure of the subsemigroupoid generated by the graph may not be a subsemigroupoid of the corresponding free profinite semigroupoid. This problem was overlooked in [5] and [4], where infinite-vertex free profinite semigroupoids are seriously considered for the first time as a tool to be used in the study of relatively free profinite semigroups. While their role in [5] can be considered marginal, in [4] some specific relatively free profinite semigroupoids with an infinite number of vertices play a key role in the proofs of the main results remain valid by establishing that, for the infinite-vertex graphs which are used in the proofs, the semigroupoids generated by the graphs are dense in the corresponding relatively free profinite semigroupoids.

Received March 2, 2015.

This research was partially supported by Centro de Matemática da Universidade do Porto (CMUP) [J. Almeida] and Centro de Matemática da Universidade de Coimbra (CMUC) [A. Costa], both funded by the European Regional Development Fund, through the programme COMPETE, and by the Portuguese Government through Fundação para a Ciência e a Tecnologia (FCT), under, respectively, the projects PEst-C/MAT/UI0144/2013 and PEst-C/MAT/UI0324/2013.

2. The free profinite semigroupoid generated by $\partial_X P$

For general background on pseudovarieties of semigroups and of semigroupoids, and on their relatively free profinite structures, see [2, 1, 8]. We also adopt the notation of [4], which we recall here for the reader's benefit. Let A be a finite alphabet and let \mathbf{V} be a pseudovariety of semigroups. The free pro- \mathbf{V} semigroup generated by A is denoted by $\overline{\Omega}_A \mathbf{V}$, and the subsemigroup of $\overline{\Omega}_A \mathbf{V}$ generated by A is denoted by $\Omega_A \mathbf{V}$. Suppose that \mathbf{V} contains the pseudovariety \mathbf{Sl} of finite semilattices. This hypothesis enables us to consider the *content*, which is a continuous homomorphism c from $\overline{\Omega}_A \mathbf{V}$ to the union-semilattice of subsets of A that maps each letter $a \in A$ to the singleton set $\{a\}$. The content function extends to $(\overline{\Omega}_A \mathbf{V})^1$ by letting $c(1) = \emptyset$. Suppose also that $\overline{\Omega}_A \mathbf{V}$ has $0, \bar{0}, 1, \bar{1}$ functions, which means that, for every $w \in \overline{\Omega}_A \mathbf{V}$, there are factorizations $w = w_0 a w' = w'' b w_1$ in $(\overline{\Omega}_A \mathbf{V})^1$ such that $a, b \in A$ and

$$c(w) = c(w_0) \uplus \{a\} = \{b\} \uplus c(w_1),$$

where the factors w_0, a, b, w_1 are unique. We put $0(w) := w_0$, $1(w) := w_1$, $\bar{0}(w) := a$, and $\bar{1}(w) := b$. Note that the functions $0, \bar{0}, 1, \bar{1}$ are continuous.

Let X be a subset of A with at least two elements. Consider a subset P of $\overline{\Omega}_A \mathbf{V}$ such that $c(u) \subseteq X$ for every $u \in P$. We define a graph* $\partial_X P$ on the set of all factors $v \in (\overline{\Omega}_A \mathbf{V})^1$ of elements of P such that $|X \setminus c(v)| \in \{1, 2\}$. The vertices are those v such that $|X \setminus c(v)| = 2$ and the edges are the remaining elements of $\partial_X P$. The adjacency functions are $\alpha = 0$ and $\omega = 1$, where, in a graph, we respectively denote by $\alpha(u)$ and $\omega(u)$ the source and the target of an edge u . This definition of $\partial_X P$ is taken from [4, Subsection 3.1], where the necessary hypothesis that the content of all elements of P is contained in X is not made explicit, but is implicitly used.

Profinite graphs are defined in [4] as being the inverse limits of finite graphs, in the category of topological graphs, with finite graphs having the discrete topology. It is folklore that the profinite graphs are precisely the topological graphs whose topology is a Boolean space (a proof can be found in [9].) It follows immediately from this characterization that the graph $\partial_X P$ is profinite if P is a closed subset of $\overline{\Omega}_A \mathbf{V}$, which happens in particular when $P = \overline{\Omega}_X \mathbf{V}$, the case considered in the main results of [4].

Let \mathbf{W} be a pseudovariety of finite semigroupoids. We adopt the definition from [3] of a *pro- \mathbf{W}* semigroupoid as being a compact semigroupoid which

*Throughout this paper, by graph we mean a directed graph.

is *residually in W* in the sense that, for every pair u, v of distinct elements of S , there is a continuous semigroupoid homomorphism $\varphi: S \rightarrow F$ into a semigroupoid F of W satisfying $\varphi(u) \neq \varphi(v)$. For the case where W is the pseudovariety \mathbf{Sd} of all finite semigroupoids, one uses “*profinite*” as a synonym of “*pro- W* ”. We remark that there is an unpublished example due to G. Bergman (mentioned in [7]) of an infinite-vertex semigroupoid which is profinite according to this definition, but which is not an inverse limit of finite semigroupoids.

Let Γ be a profinite graph and let W be a pseudovariety of semigroupoids. The free pro- W semigroupoid generated by Γ , denoted $\overline{\Omega}_\Gamma W$, is a pro- W semigroupoid, together with a continuous graph homomorphism $\iota: \Gamma \rightarrow \overline{\Omega}_\Gamma W$, with the following property: for every continuous graph homomorphism $\varphi: \Gamma \rightarrow F$ into a semigroupoid of W , there is a unique continuous semigroupoid homomorphism $\hat{\varphi}: \overline{\Omega}_\Gamma W \rightarrow F$ such that $\hat{\varphi} \circ \iota = \varphi$. It is easy to show that this semigroupoid is indeed unique, up to isomorphism of compact semigroupoids, and a proof of its existence is made in [3] by a reduction to the finite-vertex case treated in [6]. Moreover, if W contains nontrivial semigroups, we may assume (as we do from hereon) that ι is the inclusion mapping [3]. The subsemigroupoid of $\overline{\Omega}_\Gamma W$ generated by $\iota(\Gamma)$ is denoted by $\Omega_\Gamma W$. If Γ has a finite number of vertices then $\Omega_\Gamma W$ is dense in $\overline{\Omega}_\Gamma W$ [6]. However, in general, that property fails, and it is only by iterating transfinitely algebraic and topological closures that one reaches the profinite semigroupoid $\overline{\Omega}_\Gamma W$ [3].

As remarked in [6], if Γ is a finite graph, then $\overline{\Omega}_\Gamma W$ is metrizable. The following is a generalization of this fact, which will be used later on.

Proposition 2.1. *Let Γ be a profinite graph and let W be a pseudovariety of semigroupoids. If Γ is metrizable, then so is $\overline{\Omega}_\Gamma W$.*

For the proof of Proposition 2.1 one uses the next folklore result, for which we do not have a direct reference. However, a proof is implicit in the proof of [10, Corollary 1.1.13] (there the reference is made to profinite second countable spaces, but by Urysohn’s metrization theorem [11, Theorem 23.1], a profinite space is second countable if and only if it is metrizable.) In the statement we use the notation in [10, Chapter I] for inverse systems.

Proposition 2.2. *Let the topological space X be the inverse limit of an inverse system $\{X_i, \varphi_{ij}, I\}$ of finite discrete spaces. Then X is metrizable if and only if for some countable totally ordered subset J of I (with order type*

equal to that of ω) the space X is the inverse limit of the restricted inverse system $\{X_i, \varphi_{ij}, J\}$.

Proof of Proposition 2.1: By Proposition 2.2, the profinite graph Γ is, in the category of profinite graphs, an inverse limit $\varprojlim_{n \geq 1} \Gamma_n$ for some inverse sequence of finite graphs Γ_n . By the construction of $\overline{\Omega}_\Gamma \mathbf{W}$ given in [3], the semigroupoid $\overline{\Omega}_\Gamma \mathbf{W}$ embeds as a closed subsemigroupoid of $\varprojlim_{n \geq 1} \overline{\Omega}_{\Gamma_n} \mathbf{W}$. Since Γ_n is finite, the profinite semigroupoid $\overline{\Omega}_{\Gamma_n} \mathbf{W}$ is metrizable. As the inverse limit of a sequence of metrizable spaces is metrizable (cf. [11, Theorem 22.3]), it follows that $\varprojlim_{n \geq 1} \overline{\Omega}_{\Gamma_n} \mathbf{W}$ and $\overline{\Omega}_\Gamma \mathbf{W}$ are metrizable. ■

Corollary 2.3. *The profinite semigroupoid $\overline{\Omega}_{\partial_X \overline{\Omega}_A \mathbf{V}} \mathbf{W}$ is metrizable, for every finite alphabet A and every subset X of A with at least two elements.*

Proof: Since A is finite, $\overline{\Omega}_A \mathbf{V}$ is metrizable, and therefore so is $\partial_X \overline{\Omega}_A \mathbf{V}$. ■

The relatively free profinite semigroupoids which intervene in the main results in [4] are of the form $\overline{\Omega}_{\partial_X \overline{\Omega}_A \mathbf{V}} \mathbf{W}$. We remark that in (the applications of) Theorem 2.5 of [4], the graph $\partial_X \Omega_A \mathbf{V}$ is identified with the graph $\partial_X(A^+)$ and the subsemigroupoid $\langle \partial_X \Omega_A \mathbf{V} \rangle$ of $\overline{\Omega}_{\partial_X \overline{\Omega}_A \mathbf{V}} \mathbf{W}$ generated by $\partial_X \Omega_A \mathbf{V}$ is identified with the free semigroupoid $(\partial_X(A^+))^+$. These identifications hold in many cases, namely when \mathbf{V} and \mathbf{W} contain all finite nilpotent semigroups: we then have $\Omega_A \mathbf{V} \cong A^+$ and $\Omega_\Gamma \mathbf{W} \cong \Gamma^+$ (cf. [3, Theorem 3.16].) But, for example, denoting by \mathbf{Ab} the pseudovariety of finite Abelian groups, we know that $\Omega_A \mathbf{Ab}$ is not isomorphic to A^+ . To fix this problem it suffices to replace $\partial_X(A^+)$ by $\partial_X \Omega_A \mathbf{V}$ and $(\partial_X(A^+))^+$ by $\langle \partial_X \Omega_A \mathbf{V} \rangle$ when appropriate in [4].

Another problem in [4] that needs to be fixed stems from the fact that in the proof of Theorem 2.5 in [4] it is assumed that the subsemigroupoid of $\overline{\Omega}_{\partial_X \overline{\Omega}_A \mathbf{V}} \mathbf{W}$ generated by $\partial_X \Omega_A \mathbf{V} \subseteq \partial_X \overline{\Omega}_A \mathbf{V}$ is dense in $\overline{\Omega}_{\partial_X \overline{\Omega}_A \mathbf{V}} \mathbf{W}$. As already mentioned, after the publication of [4], examples of profinite graphs Γ such that $\Omega_\Gamma \mathbf{W}$ is not dense in $\overline{\Omega}_\Gamma \mathbf{W}$ were given in [3]. Therefore, one needs to verify if the denseness assumption made in the proof of Theorem 2.5 in [4] really holds. In the next proposition, we show that it does hold when \mathbf{V} is a pseudovariety of semigroups such that $\mathbf{Sl} \subseteq \mathbf{V} \subseteq \mathbf{CR}$, where \mathbf{CR} denotes the pseudovariety of completely regular semigroups, and such that $\overline{\Omega}_A \mathbf{V}$ has 0 , $\bar{0}$, 1 , $\bar{1}$ functions for every finite set A . This suffices to guarantee the main results in Sections 3 to 6 of [4], all about pseudovarieties satisfying these conditions.

Theorem 2.4. *Let \mathbf{V} be a pseudovariety of semigroups such that $\mathbf{SI} \subseteq \mathbf{V} \subseteq \mathbf{CR}$ and let \mathbf{W} be any pseudovariety of semigroupoids containing some non-trivial semigroup. Let A be a finite alphabet, and let X be a subset of A with at least two elements. Suppose that $\overline{\Omega}_A \mathbf{V}$ has $0, \bar{0}, 1, \bar{1}$ functions. Then the subsemigroupoid $\langle \partial_X \Omega_A \mathbf{V} \rangle$ of $\overline{\Omega}_{\partial_X \overline{\Omega}_A \mathbf{V}} \mathbf{W}$ is dense. In particular, $\Omega_{\partial_X \overline{\Omega}_A \mathbf{V}} \mathbf{W}$ is dense in $\overline{\Omega}_{\partial_X \overline{\Omega}_A \mathbf{V}} \mathbf{W}$.*

Proof: To avoid confusion between the multiplication in $\overline{\Omega}_A \mathbf{V}$ and the edge multiplication in $\overline{\Omega}_{\partial_X \overline{\Omega}_A \mathbf{V}} \mathbf{W}$, we denote the multiplication of two composable elements e, f of $\overline{\Omega}_{\partial_X \overline{\Omega}_A \mathbf{V}} \mathbf{W}$ by $e \circ f$.

Note that $\partial_X \overline{\Omega}_A \mathbf{V} = \overline{\partial_X \Omega_A \mathbf{V}}$, so that, in particular, $\overline{\langle \partial_X \Omega_A \mathbf{V} \rangle}$ contains $\partial_X \overline{\Omega}_A \mathbf{V}$. Therefore, since $\overline{\Omega}_{\partial_X \overline{\Omega}_A \mathbf{V}} \mathbf{W}$ is the unique closed subsemigroupoid of $\overline{\Omega}_{\partial_X \overline{\Omega}_A \mathbf{V}} \mathbf{W}$ containing $\partial_X \overline{\Omega}_A \mathbf{V}$, to show that $\langle \partial_X \Omega_A \mathbf{V} \rangle$ is dense in $\overline{\Omega}_{\partial_X \overline{\Omega}_A \mathbf{V}} \mathbf{W}$ it suffices to show that $\overline{\langle \partial_X \Omega_A \mathbf{V} \rangle}$ is a subsemigroupoid of $\overline{\Omega}_{\partial_X \overline{\Omega}_A \mathbf{V}} \mathbf{W}$.

Thanks to Corollary 2.3, what we want to show translates into proving that if two edges e and f of $\overline{\Omega}_{\partial_X \overline{\Omega}_A \mathbf{V}} \mathbf{W}$ are limits of sequences $(e_n)_n$ and $(f_n)_n$ of elements of $\langle \partial_X \Omega_A \mathbf{V} \rangle$ such that $e \circ f$ is defined, then $e \circ f$ is also the limit of a sequence of elements of $\langle \partial_X \Omega_A \mathbf{V} \rangle$. This property is established if one shows that it is possible to replace each f_n by another element \tilde{f}_n of $\langle \partial_X \Omega_A \mathbf{V} \rangle$ such that $e_n \circ \tilde{f}_n$ is defined and $(\tilde{f}_n)_n$ converges to f . The claim follows by continuity of edge multiplication in $\overline{\Omega}_{\partial_X \overline{\Omega}_A \mathbf{V}} \mathbf{W}$. Actually, the replacement needs only to be done on a subsequence, so we may take subsequences of $(e_n)_n$ and $(f_n)_n$ as convenient.

The only relevant part of e is its end vertex u . We denote by u_n the end vertex of e_n and by u'_n the beginning vertex of f_n . We then have $\lim u_n = u = \lim u'_n$. Since the content function is continuous, by taking subsequences we may assume that $c(u_n) = c(u) = c(u'_n)$ for every n . Since $f_n \in \langle \partial_X \Omega_A \mathbf{V} \rangle$, there is a factorization $f_n = s_n \circ t_n$ such that $s_n \in \partial_X \Omega_A \mathbf{V}$ and $t_n \in \langle \partial_X \Omega_A \mathbf{V} \rangle \uplus \{I_n\}$, where I_n denotes an adjoined local identity. Again taking subsequences, we may assume that $(s_n)_n$ converges to s and $(t_n)_n$ converges to t (where t may be an adjoined local identity I .) Note that, by continuity of \circ , we have $f = s \circ t$. Consider factorizations of the form

$$s_n = 0(s_n)\bar{0}(s_n)x_n = y_n\bar{1}(s_n)1(s_n),$$

with $x_n, y_n \in (\overline{\Omega}_A \mathbf{V})^1$. We have $0(s_n) = \alpha(s_n) = u'_n$ and $\omega(s_n) = 1(s_n) = \alpha(t_n)$. Once again by taking subsequences, we may suppose that $(x_n)_n$ and $(y_n)_n$ respectively converge to some elements x and y of $(\overline{\Omega}_A \mathbf{V})^1$. Consider

the following element of $\overline{\Omega}_A \mathbf{V}$:

$$r_n = u_n \bar{0}(s_n) x_n \cdot s_n^{n!}.$$

Since $c(r_n) = c(s_n)$, we know that r_n is an edge of $\partial_X \Omega_A \mathbf{V}$ such that $1(r_n) = 1(s_n)$, and we also have $0(r_n) = u_n$ due to the equality $c(u_n) = c(0(s_n))$. Therefore, we may define the element $e_n \circ r_n \circ t_n$ of $\langle \partial_X \Omega_A \mathbf{V} \rangle$. By continuity of $\bar{0}$, we have

$$\lim u_n \bar{0}(s_n) x_n = u \bar{0}(s) x = \lim 0(s_n) \bar{0}(s_n) x_n = s.$$

On the other hand, the sequence $(s_n^{n!})_n$ converges to the idempotent s^ω . Therefore, we have $\lim r_n = s^{\omega+1} = s$, where the last equality holds because $\mathbf{V} \subseteq \mathbf{CR}$. Hence, we obtained a sequence $(e_n \circ r_n \circ t_n)_n$ of elements of $\langle \partial_X \Omega_A \mathbf{V} \rangle$ converging to $e \circ s \circ t = e \circ f$. ■

References

- [1] J. Almeida. *Finite Semigroups and Universal Algebra*. World Scientific, Singapore, 1995. English translation.
- [2] J. Almeida. Profinite semigroups and applications. In V. B. Kudryavtsev and I. G. Rosenberg, editors, *Structural Theory of Automata, Semigroups and Universal Algebra*, pages 1–45, New York, 2005. Springer.
- [3] J. Almeida and A. Costa. Infinite-vertex free profinite semigroupoids and symbolic dynamics. *J. Pure Appl. Algebra*, 213:605–631, 2009.
- [4] J. Almeida and P. G. Trotter. The pseudoidentity problem and reducibility for completely regular semigroups. *Bull. Austral. Math. Soc.*, 63:407–433, 2001.
- [5] J. Almeida and P. Weil. Profinite categories and semidirect products. *J. Pure Appl. Algebra*, 123:1–50, 1998.
- [6] P. R. Jones. Profinite categories, implicit operations and pseudovarieties of categories. *J. Pure Appl. Algebra*, 109:61–95, 1996.
- [7] J. Rhodes and B. Steinberg. Profinite semigroups, varieties, expansions and the structure of relatively free profinite semigroups. *Int. J. Algebra Comput.*, 11:627–672, 2002.
- [8] J. Rhodes and B. Steinberg. *The q-theory of finite semigroups*. Springer Monographs in Mathematics. Springer, 2009.
- [9] L. Ribes. Grupos profinitos y grafos topológicos. Number 4 in Publicacions de la Secció de Matemàtiques, pages 1–64. Universitat Autònoma de Barcelona, 1977.
- [10] L. Ribes and P. A. Zalesskiĭ. *Profinite Groups*. Number 40 in *Ergeb. Math. Grenzgebiete 3*. Springer, Berlin, 2000.
- [11] S. Willard. *General Topology*. Addison-Wesley, Reading, Mass., 1970.

JORGE ALMEIDA

CMUP, DEPARTAMENTO DE MATEMÁTICA, FACULDADE DE CIÊNCIAS, UNIVERSIDADE DO PORTO, RUA DO CAMPO ALEGRE 687, 4169-007 PORTO, PORTUGAL.

E-mail address: jalmeida@fc.up.pt

ALFREDO COSTA

CMUC, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF COIMBRA, 3001-501 COIMBRA, PORTUGAL.

E-mail address: amgc@mat.uc.pt