

HOMOGENIZATION OF OBSTACLE PROBLEMS IN ORLICZ-SOBOLEV SPACES

DIEGO MARCON, JOSÉ FRANCISCO RODRIGUES AND RAFAYEL TEYMURAZYAN

ABSTRACT: We study homogenization of obstacle problems in Orlicz-Sobolev spaces for a wide class of monotone operators (possibly degenerate or singular) of $p(\cdot)$ -Laplacian type. Our approach is based on Lewy-Stampacchia inequalities, which then give access to a compactness argument. We also prove convergence of the coincidence sets under non-degeneracy conditions.

KEYWORDS: Homogenization, obstacle problem, Orlicz-Sobolev spaces, convergence of coincidence sets.

AMS SUBJECT CLASSIFICATION (2010): 35J20, 35J62, 35J92, 35D30, 35A15.

1. Introduction

Let $\Omega \subset \mathbb{R}^n$ be a bounded domain and $p : \Omega \rightarrow \mathbb{R}$ be measurable and such that

$$1 < \alpha \leq p(x) \leq \beta < \infty \quad \text{a.e. in } \Omega, \quad (1.1)$$

where α and β are constants. The following variable exponent Lebesgue space is an Orlicz space:

$$L^{p(\cdot)}(\Omega) := \left\{ u : \Omega \rightarrow \mathbb{R} \text{ measurable } \rho(u) := \int_{\Omega} |u(x)|^{p(x)} dx < \infty \right\}.$$

This Orlicz space is a separable reflexive Banach space with the following (Luxembourg) norm:

$$\|u\|_{L^{p(\cdot)}(\Omega)} := \inf \left\{ \lambda > 0, \rho\left(\frac{|u|}{\lambda}\right) \leq 1 \right\}.$$

We define an Orlicz-Sobolev space by

$$W^{1,p(\cdot)}(\Omega) := \left\{ u \in L^{p(\cdot)}(\Omega), \nabla u \in \left(L^{p(\cdot)}(\Omega) \right)^n \right\},$$

Received June 22, 2018.

This work was partially supported by FCT-Portugal grant SFRH/BPD/92717/2013. RT thanks CMUC for providing excellent working conditions.

with the norm

$$\|u\|_{W^{1,p(\cdot)}(\Omega)} := \|u\|_{L^{p(\cdot)}(\Omega)} + \|\nabla u\|_{L^{p(\cdot)}(\Omega)}, \quad \|\nabla u\|_{L^{p(\cdot)}(\Omega)} = \sum_{i=1}^n \left\| \frac{\partial u}{\partial x_i} \right\|_{L^{p(\cdot)}(\Omega)}.$$

This Orlicz-Sobolev space is also a separable and reflexive Banach space. We also define

$$W_0^{1,p(\cdot)}(\Omega) := \left\{ u \in W_0^{1,1}(\Omega), \rho(|\nabla u|) < \infty \right\}.$$

The latter is a Banach space endowed with the norm

$$\|u\|_{W_0^{1,p(\cdot)}(\Omega)} := \|\nabla u\|_{L^{p(\cdot)}(\Omega)}.$$

In this paper we study the periodic homogenization of obstacle problems in Orlicz-Sobolev spaces. We consider

$$a(x, \xi) : \Omega \times \mathbb{R}^n \rightarrow \mathbb{R}^n$$

to be a Carathéodory vector function, that is, we assume it is continuous with respect to ξ , for almost every $x \in \mathbb{R}^n$, and that it is measurable with respect to x , for every ξ . Moreover, the functions $a(\cdot, \xi)$ and $p(\cdot)$ are assumed to be periodic with period 1 in each argument x_1, x_2, \dots, x_n . We denote the periodicity cell by Q , i.e. $Q := (0, 1]^n$. Additionally, we assume that the following structural conditions (monotonicity, coercitivity and boundedness) hold:

$$\begin{cases} (a(x, \xi) - a(x, \eta)) \cdot (\xi - \eta) > 0, & \text{for a.e. } x, \xi \neq \eta, \\ a(x, \xi) \cdot \xi \geq C_1 (|\xi|^{p(x)} - 1), \\ |a(x, \xi)| \leq C_2 (|\xi|^{p(x)-1} + 1), \end{cases} \quad (1.2)$$

where $C_1, C_2 > 0$ are constants. For $\varepsilon > 0$, we define

$$a_\varepsilon(x, \xi) := a\left(\frac{x}{\varepsilon}, \xi\right), \quad x \in \Omega, \xi \in \mathbb{R}^n \quad (1.3)$$

and $p_\varepsilon(x) = p(x/\varepsilon)$. The Orlicz-Sobolev spaces of periodic functions, denoted by $W_{\text{per}}^{1,p(\cdot)}(Q)$, is defined as the set of periodic functions u from $W_{\text{per}}^{1,1}(Q)$ with

$$\int_Q u \, dx = 0 \quad \text{and} \quad \int_Q |\nabla u|^{p(x)} \, dx < \infty.$$

For the homogenized functional defined by

$$h(\xi) := \min_{v \in W_{\text{per}}^{1,p(\cdot)}(Q)} \int_Q \frac{|\xi + \nabla v|^{p(x)}}{p(x)} \, dx, \quad (1.4)$$

we introduce also the Orlicz-Sobolev spaces

$$W^h(\Omega) := \{u \in W^{1,1}(\Omega), h(\nabla u) \in L^1(\Omega)\},$$

$$W_0^h(\Omega) := \{u \in W_0^{1,1}(\Omega), h(\nabla u) \in L^1(\Omega)\},$$

with the norm, $\|u\|_{W_0^h(\Omega)} := \|\nabla u\|_{L^h(\Omega)}$, and the vector Orlicz space

$$L^h(\Omega) := \{\xi \in [L^1(\Omega)]^n, h(\xi) \in L^1(\Omega)\},$$

normed by

$$\|\xi\|_{L^h(\Omega)} := \inf \left\{ \lambda > 0, \int_{\Omega} h\left(\frac{\xi}{\lambda}\right) \leq 1 \right\}.$$

By the properties of h , as it was observed in [24], we have the continuous embeddings

$$L^\beta(\Omega) \subset L^h(\Omega) \subset L^\alpha(\Omega),$$

Assuming that

$$f \text{ and } (A_\varepsilon \psi_\varepsilon - f)^+ \in L^s(\Omega), \quad (1.5)$$

$$\|(A_\varepsilon \psi_\varepsilon - f)^+\|_{L^s(\Omega)} \leq C, \quad (1.6)$$

where $C > 0$ is a constant independent of ε and

$$\psi_\varepsilon \in W^{1,p_\varepsilon(\cdot)}(\Omega), \psi_0 \in W^h(\Omega), \psi_\varepsilon^+ \in W_0^{1,p_\varepsilon(\cdot)}(\Omega), \psi_0^+ \in W_0^h(\Omega), \quad (1.7)$$

where $\alpha' = \alpha/(\alpha - 1)$, u^+ is the positive part of u and $s > \frac{n\alpha'}{n+\alpha'}$ if $\alpha < n$, $s > 1$, if $\alpha = n$ and $s = 1$ for $\alpha > n$, we show (Theorem 3.1) that the unique solution $u_\varepsilon \in K_\varepsilon$ of the obstacle problem

$$\int_{\Omega} a_\varepsilon(x, \nabla u_\varepsilon) \cdot \nabla(v - u_\varepsilon) dx \geq \int_{\Omega} f(v - u_\varepsilon) dx, \quad \forall v \in K_\varepsilon, \quad (1.8)$$

where

$$K_\varepsilon := \{v \in W_0^{1,p_\varepsilon(\cdot)}(\Omega), v \geq \psi_\varepsilon \text{ a.e. in } \Omega\},$$

converges to the unique solution $u_0 \in K_0$ of the following homogenized obstacle problem

$$\int_{\Omega} a_0(\nabla u_0) \cdot \nabla(v - u_0) dx \geq \int_{\Omega} f(v - u_0) dx, \quad \forall v \in K_0, \quad (1.9)$$

where

$$K_0 := \{v \in W_0^h(\Omega), v \geq \psi_0 \text{ a.e. in } \Omega\}.$$

The homogenized operator $a_0 : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is given in terms of the weighted average of a as in [24], that is,

$$a_0(\xi) := \int_Q a(x, \xi + \nabla v) dx, \quad (1.10)$$

with $v \in W_{\text{per}}^{1,p(\cdot)}(Q)$, such that,

$$\int_Q a(x, \xi + \nabla v) \cdot \nabla \varphi dx = 0, \quad \forall \varphi \in W_{\text{per}}^{1,p(\cdot)}(Q),$$

where Q is the periodicity cell.

Note that, due to the Lavrent'ev effect, if instead of $W_{\text{per}}^{1,p(\cdot)}(Q)$, we take $\varphi \in C_{\text{per}}^\infty(Q)$, we may end up with a different homogenized operator, since in general the space $C_{\text{per}}^\infty(Q)$ is not dense in $W_{\text{per}}^{1,p(\cdot)}(\Omega)$. These homogenized operators, referred to as W and H solutions in [24], respectively, in general may be different, but our results hold for both solutions, with minor modifications for the space framework of the H solutions. Although we prefer to work with W solutions, that is due to the fact that [24, Theorem 3.1] (see Theorem 2.1 below) is true for both types of solutions. Observe that we do not impose any regularity assumption on $p(\cdot)$. However, in the particular case when p is log-Lipschitz continuous, i.e., when for a constant $L > 0$

$$-|p(x) - p(y)| \log |x - y| \leq L, \quad \forall x, y \in \overline{\Omega}, \quad |x - y| < 1/2,$$

the notion of W and H solutions coincide (see [10, 14]), since then the smooth functions are dense in the Orlicz-Sobolev space.

Our approach is a development of the classical methods [6, 11] (see also [20, 21, 24]) combined with the Lewy-Stampacchia inequalities in the Orlicz-Sobolev framework, in accordance with [19], which then allows the use of a Rellich-Kondrachov compactness argument.

The result generalizes, in part, that of [5], which covers the case when p is constant (and hence the homogenization is in usual Sobolev spaces). The latter, in turn, implies the case of $p = 2$ obtained in [4]. Nonetheless, we observe that the structural assumptions (1.2) allow us to consider a wider range of monotone operators, which cover these cases and include other interesting quasilinear operators, some of which we list below.

1. If $a(x, \xi) = |\xi|^{p(x)-2}\xi$, we deal with the obstacle problem for the $p(x)$ -Laplace operator.

2. We can also consider perturbations of the p -Laplace (p constant) and of the $p(x)$ -Laplace operators, taking

$$a(x, \xi) = \gamma(x)|\xi|^{p-2}\xi \quad \text{and} \quad a(x, \xi) = \gamma(x)|\xi|^{p(x)-2}\xi$$

for any non-negative bounded periodic function $\gamma(x)$.

3. It is possible to consider functions which are essentially different from these previous “power like” functions. One general example can be

$$a(x, \xi) = \gamma_1(x)|\xi|^{p(x)-1}\xi \log(\gamma_2(x)|\xi| + \gamma_3(x)),$$

where $\gamma_3(x)$, $p(x) > 1$ and $\gamma_1(x)$, $\gamma_2(x) > 0$ a.e. in Ω are bounded periodic functions.

The paper is organized as follows: in Section 2, we state some preliminaries facts, which then serve to prove our main result in Section 3 (Theorem 3.1). In Section 4, we prove the convergence of the coincidence sets (Theorems 4.1 and 4.2).

2. Preliminaries

In this section we give some preliminaries. In particular, we provide the concept of G -convergence of operators in our framework, as well as convergence of sets in Mosco sense. We also recall some results from [22] and [24] for future reference. We start by setting some notations, which will be used throughout the paper: $p_\varepsilon(x) = p(x/\varepsilon)$; $\alpha' = \frac{\alpha}{\alpha-1}$; \rightharpoonup denotes the weak convergence;

$$A_\varepsilon u := -\operatorname{div}(a_\varepsilon(x, \nabla u)) \quad \text{and} \quad A_0 u := -\operatorname{div}(a_0(\nabla u)),$$

where a_ε is defined by (1.3), and a_0 is defined by (1.10). Next, we define the notion of G -convergence of a_ε to a_0 . Observe, that most definitions of G -convergence that can be found in the literature (see, for example, [2, 3, 7, 17]), allow a_0 to depend on x as well, just as a_ε depends. However, in some particular cases, more information can be said about the limiting operator. One example is that of operators with rapidly oscillating “coefficients”. Since our assumptions ensure that $a(x, \xi)$ and $p(\cdot)$ are periodic with respect to x in each of the arguments x_1, x_2, \dots, x_n , there is no loss in generality to impose a_0 to be independent of x in the definition of G -convergence, which is more relevant for our purposes.

Definition 2.1. Consider $a_\varepsilon : \Omega \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ and $a_0 : \mathbb{R}^n \rightarrow \mathbb{R}^n$ as above. We say that a_ε G -converges to a_0 when, considering the unique solution $u_\varepsilon \in W_0^{1,p_\varepsilon(\cdot)}(\Omega)$ of

$$-\operatorname{div}(a_\varepsilon(x, \nabla u_\varepsilon)) = f, \quad f \in W_0^{-1,\alpha'}(\Omega) \text{ in } \mathcal{D}'(\Omega)$$

and $u_0 \in W_0^h(\Omega)$ the unique solution of

$$-\operatorname{div}(a_0(\nabla u_0)) = f \text{ in } \mathcal{D}'(\Omega),$$

there holds:

- (1) $u_\varepsilon \rightharpoonup u_0$ in $W_0^{1,\alpha}(\Omega)$, as $\varepsilon \rightarrow 0$;
- (2) $a_\varepsilon(x, \nabla u_\varepsilon) \rightharpoonup a_0(\nabla u_0)$ in $(L^{\beta'}(\Omega))^n$, as $\varepsilon \rightarrow 0$.

Note that the choice of s in (1.5) guarantees, in particular, $f \in W^{-1,\alpha'}(\Omega)$. Additionally, $a(x, \xi)$ is assumed to be continuous with respect to ξ , for almost every $x \in \mathbb{R}^n$.

Next, we state a theorem from [24, Theorem 3.1] that insures the G -convergence of a_ε to a function a_0 , as $\varepsilon \rightarrow 0$, given explicitly in terms of a . Its proof is based on a compensated compactness argument from [23, 24], which, in the case of $p(\cdot) = \text{constant}$, resembles the well known result of Tartar-Murat (see [16]).

Theorem 2.1. Let $a(x, \xi)$ be a Carathéodory vector function, which is periodic with respect to x in each argument and satisfy (1.2). Let also p be periodic, measurable and satisfy (1.1). If structural conditions (1.2) hold, then a_ε G -converges to a_0 , where a_0 is defined by (1.10). Moreover,

$$\int_{\Omega} a_\varepsilon(x, \nabla u_\varepsilon) \cdot \nabla u_\varepsilon \, dx \rightarrow \int_{\Omega} a_0(\nabla u_0) \cdot \nabla u_0 \, dx,$$

as $\varepsilon \rightarrow 0$.

As it is shown in [24], the vector function $a_0(\xi)$ is strictly monotone, i.e.,

$$(a_0(\xi) - a_0(\eta)) \cdot (\xi - \eta) > 0, \quad \xi \neq \eta,$$

and coercive, that is,

$$a_0(\xi) \cdot \xi > c_0(h(\xi) - 1),$$

where $c_0 > 0$ is a constant, and the homogenized functional $h(\xi)$ is defined by (1.4). Moreover, h satisfies the so-called Δ_2 condition, [24, Proposition 2.1], which implies that the Orlicz space $L^h(\Omega)$ is reflexive. As it is observed in

[24], $h(\xi)$ being defined by (1.4), is convex on \mathbb{R}^n and satisfies the following two-sided estimate:

$$c_1|\xi|^\alpha - 1 \leq h(\xi) \leq c_2|\xi|^\beta + 1,$$

for a $c_1 > 0$ constant. As a consequence, we have

$$W_0^{1,\beta}(\Omega) \subset W_0^h(\Omega) \subset W_0^{1,\alpha}(\Omega),$$

which implies that

$$K_0 \subset W_0^{1,\alpha}(\Omega).$$

The following result is from [22], and it provides more information on the homogenized functional.

Lemma 2.1. *If u_ε is a sequence uniformly bounded in $W_0^{1,p_\varepsilon(\cdot)}(\Omega)$, such that, $u_\varepsilon \rightharpoonup u_0$ in $W_0^{1,\alpha}(\Omega)$ as $\varepsilon \rightarrow 0$, then $h(\nabla u_0) \in L^1(\Omega)$.*

Observe that Lemma 2.1 guarantees that, within G -convergence, the weak limits of u_ε in $W_0^{1,\alpha}(\Omega)$ belong to $W_0^h(\Omega)$, and therefore, if also $u_\varepsilon \in K_\varepsilon$ then $u_0 \in K_0$.

In order to state our main result, we will also need to redefine the Mosco convergence of sets.

Definition 2.2. *The sequence of closed convex sets $K_\varepsilon \subset W_0^{1,p_\varepsilon(\cdot)}(\Omega)$, is said to converge to the set $K_0 \subset W_0^h(\Omega)$ in the Mosco sense, if*

- for any $v_0 \in K_0$ there exists a sequence $v_\varepsilon \in K_\varepsilon$, such that, $v_\varepsilon \rightarrow v_0$ in $W_0^{1,\alpha}(\Omega)$;
- weak limits in $W_0^{1,\alpha}(\Omega)$ of any sequence of elements in K_ε , that is uniformly bounded in $W_0^{1,p_\varepsilon(\cdot)}(\Omega)$, belong to K_0 .

Remark 2.1. *Since $W_0^{1,p_\varepsilon(\cdot)}(\Omega)$ is continuously embedded into $W_0^{1,\alpha}(\Omega)$ (see, for example, [10]), then $\psi_\varepsilon \rightarrow \psi_0$ in $W^{1,\beta}(\Omega)$ provides $K_\varepsilon \rightarrow K_0$ in the Mosco sense, where K_ε and K_0 are as in (1.8) and (1.9) respectively.*

3. Homogenization of the obstacle problem

We are now ready to prove our main result, which states as follows.

Theorem 3.1. *Let $a(x, \xi)$ be a Carathéodory vector function satisfying (1.2) and periodic with respect to x in each argument. Let $p(\cdot)$ be periodic, measurable and satisfying (1.1). Assume further that (1.5)-(1.7) hold. If $K_\varepsilon \rightarrow K_0$ in the Mosco sense, then the unique solution of (1.8) converges weakly in*

$W_0^{1,\alpha}(\Omega)$, as $\varepsilon \rightarrow 0$, to the unique solution of (1.9), where a_0 is given by (1.10).

Proof: We divide the proof into five steps.

Step 1 (*A priori estimates*). Existence and uniqueness of the solution of (1.8) (and (1.9)) is a classical result (see, for instance, [9, 18, 19]). As in the proof of [5, Theorem 2.3] (see also [18, page 145]), the coercitivity and boundedness assumptions from (1.2) imply that u_ε is bounded in $W_0^{1,p_\varepsilon(\cdot)}(\Omega)$ by a constant depending only from C_1, C_2 but independent of ε . For the details we refer the reader to [12]. As a consequence we obtain that u_ε is bounded also in $W_0^{1,\alpha}(\Omega)$, since $W_0^{1,p_\varepsilon(\cdot)}(\Omega) \subset W_0^{1,\alpha}(\Omega)$. Set

$$\sigma_\varepsilon := a_\varepsilon(x, \nabla u_\varepsilon), \quad \mu_\varepsilon := -\operatorname{div}(a_\varepsilon(x, \nabla u_\varepsilon)) - f. \quad (3.1)$$

The boundedness condition from (1.2) implies that σ_ε and μ_ε are bounded (see [5, 24]), therefore we can extract weakly convergent subsequence (still denoted by ε) from each one of them. Thus, there exist u^*, σ^*, μ^* such that

$$u_\varepsilon \rightharpoonup u^* \quad \text{in } W_0^{1,\alpha}(\Omega) \quad \text{and} \quad u_\varepsilon \rightarrow u^* \quad \text{in } L^\alpha(\Omega), \quad (3.2)$$

$$\sigma_\varepsilon \rightharpoonup \sigma^* \quad \text{in } \left(L^{\beta'}(\Omega)\right)^n, \quad (3.3)$$

$$\mu_\varepsilon \rightharpoonup \mu^* \quad \text{in } W^{-1,\beta'}(\Omega). \quad (3.4)$$

Note that

$$\mu^* = -\operatorname{div}\sigma^* - f. \quad (3.5)$$

Moreover, using Lemma 2.1 and since $K_\varepsilon \rightarrow K_0$ in the Mosco sense, then

$$u^* \in K_0. \quad (3.6)$$

Step 2 (*Compactness*). Note that our assumptions provide the Lewy-Stampacchia inequalities (see [19]), that is, we have

$$f \leq f + \mu_\varepsilon \leq (A_\varepsilon \psi_\varepsilon - f)^+ + f,$$

which implies, by a Rellich-Kondrachov compactness argument,

$$\mu_\varepsilon \rightarrow \mu^* \quad \text{in } W^{-1,\alpha'}(\Omega). \quad (3.7)$$

Step 3. In this step we prove that $\sigma^* = a_0(\nabla u^*)$, where a_0 is defined by (1.10). To see this, let $w_0 \in \mathcal{D}(\Omega)$ and $w_\varepsilon \in W_0^{1,p_\varepsilon(\cdot)}(\Omega)$ be the unique solution of

$$\operatorname{div}(a_\varepsilon(x, \nabla w_\varepsilon)) = \operatorname{div}(a_0(\nabla w_0)) \quad \text{in } \mathcal{D}'(\Omega). \quad (3.8)$$

From Theorem 2.1, we have that a_ε G -converges to a_0 , as $\varepsilon \rightarrow 0$, where $a_0(\xi)$ is defined by (1.10). In particular,

$$\begin{cases} w_\varepsilon \rightharpoonup w_0 & \text{in } W^{1,\alpha}(\Omega) \\ a_\varepsilon(x, \nabla w_\varepsilon) \rightharpoonup a_0(\nabla w_0) & \text{in } (L^{\beta'}(\Omega))^n. \end{cases} \quad (3.9)$$

Fix now φ such that

$$\varphi \in \mathcal{D}(\Omega), \quad 0 \leq \varphi \leq 1. \quad (3.10)$$

From the monotonicity of a_ε one has

$$\int_{\Omega} \varphi (a_\varepsilon(x, \nabla u_\varepsilon) - a_\varepsilon(x, \nabla w_\varepsilon)) \cdot (\nabla u_\varepsilon - \nabla w_\varepsilon) dx \geq 0. \quad (3.11)$$

Since $u^* \in K_0$, and $K_\varepsilon \rightarrow K_0$ in the Mosco sense, there exists a sequence \bar{u}_ε , such that,

$$\bar{u}_\varepsilon \in K_\varepsilon \quad \text{and} \quad \bar{u}_\varepsilon \rightarrow u^* \quad \text{in } W_0^{1,\alpha}(\Omega). \quad (3.12)$$

Next, we write (3.11) as

$$\begin{aligned} \int_{\Omega} \varphi \sigma_\varepsilon \cdot (\nabla u_\varepsilon - \nabla \bar{u}_\varepsilon) &+ \int_{\Omega} \varphi \sigma_\varepsilon \cdot \nabla \bar{u}_\varepsilon - \int_{\Omega} \varphi \sigma_\varepsilon \cdot \nabla w_\varepsilon \\ &- \int_{\Omega} \varphi a_\varepsilon(x, \nabla w_\varepsilon) \cdot \nabla (u_\varepsilon - w_\varepsilon) \\ &:= I_1 + I_2 + I_3 + I_4. \end{aligned} \quad (3.13)$$

Since $0 \leq \varphi \leq 1$ on Ω , and K_ε is convex, then the function $v = \varphi \bar{u}_\varepsilon + (1-\varphi)u_\varepsilon$ can be used as a test function in (1.8), which gives

$$\int_{\Omega} \sigma_\varepsilon \cdot \nabla (\varphi(\bar{u}_\varepsilon - u_\varepsilon)) \geq \int_{\Omega} f \varphi (\bar{u}_\varepsilon - u_\varepsilon) \quad (3.14)$$

and so

$$\begin{aligned} I_1 &= \int_{\Omega} \sigma_\varepsilon \cdot \nabla (\varphi(u_\varepsilon - \bar{u}_\varepsilon)) - \int_{\Omega} (u_\varepsilon - \bar{u}_\varepsilon) \sigma_\varepsilon \cdot \nabla \varphi \\ &\leq \int_{\Omega} f \varphi (u_\varepsilon - \bar{u}_\varepsilon) - \int_{\Omega} (u_\varepsilon - \bar{u}_\varepsilon) \sigma_\varepsilon \cdot \nabla \varphi. \end{aligned}$$

Since u_ε and \bar{u}_ε converge to u^* weakly in $W_0^{1,\alpha}(\Omega)$ (and strongly in $L^\alpha(\Omega)$), we obtain

$$\limsup_{\varepsilon \rightarrow 0} I_1 \leq 0. \quad (3.15)$$

As we know from (3.12), $\bar{u}_\varepsilon \rightarrow u^*$ in $W_0^{1,\alpha}(\Omega)$, which gives

$$\lim_{\varepsilon \rightarrow 0} I_2 = \int_{\Omega} \varphi \sigma^* \cdot \nabla u^*. \quad (3.16)$$

Note that

$$I_3 = - \int_{\Omega} \sigma_\varepsilon \cdot \nabla (\varphi w_\varepsilon) + \int_{\Omega} w_\varepsilon \sigma_\varepsilon \cdot \nabla \varphi.$$

From (3.7) and (3.12), we pass to the limit in the first term of I_3 . Using (3.3) and (3.12), we pass to the limit also in the second term of I_3 , arriving at

$$\lim_{\varepsilon \rightarrow 0} I_3 = - \int_{\Omega} \varphi \sigma^* \nabla w_0. \quad (3.17)$$

Observe that

$$I_4 = - \int_{\Omega} a_\varepsilon(x, \nabla w_\varepsilon) \cdot \nabla (\varphi(u_\varepsilon - w_\varepsilon)) + \int_{\Omega} (u_\varepsilon - w_\varepsilon) a_\varepsilon(x, \nabla w_\varepsilon) \cdot \nabla \varphi,$$

and recalling (3.2) and (3.9) and passing to the limit we obtain

$$\lim_{\varepsilon \rightarrow 0} I_4 = - \int_{\Omega} \varphi a_0(\nabla w_0) \cdot \nabla (u^* - w_0). \quad (3.18)$$

Combining (3.13), (3.15)-(3.18), one has

$$\int_{\Omega} \varphi (\sigma^* - a_0(\nabla w_0)) \cdot \nabla (u^* - w_0) \geq 0 \quad \text{for } w_0 \in \mathcal{D}(\Omega). \quad (3.19)$$

By density, (3.19) is true also for any w_0 in $W_0^{1,\alpha}(\Omega)$. Consider $w_0 = u^* + t\varphi$, with $t \geq 0$ and $\varphi \in W_0^{1,\alpha}(\Omega)$. Letting $t \rightarrow 0$ and using Minty's trick as in [5, page 94] (see also [15]), we conclude

$$\sigma^* = a_0(\nabla u^*). \quad (3.20)$$

Step 4 (*Lower semicontinuity of the energy*). From (3.11) and (3.13) one has

$$\begin{aligned} \int_{\Omega} \varphi \sigma_\varepsilon \cdot \nabla u_\varepsilon &\geq \int_{\Omega} \varphi \sigma_\varepsilon \cdot \nabla w_\varepsilon + \int_{\Omega} \varphi a_\varepsilon(x, \nabla w_\varepsilon) \cdot \nabla (u_\varepsilon - w_\varepsilon) \\ &= -I_3 - I_4. \end{aligned}$$

From (3.17), (3.18) and (3.20) for any $w_0 \in \mathcal{D}(\Omega)$ we have

$$\begin{aligned} & \liminf_{\varepsilon \rightarrow 0} \int_{\Omega} \varphi \sigma_{\varepsilon} \cdot \nabla u_{\varepsilon} \\ & \geq \int_{\Omega} \varphi a_0(\nabla u^*) \cdot \nabla w_0 + \int_{\Omega} \varphi a_0(\nabla w_0) \cdot \nabla(u^* - w_0). \end{aligned} \quad (3.21)$$

Letting w_0 go to u^* in $W_0^{1,\alpha}(\Omega)$, one gets from (3.21)

$$\liminf_{\varepsilon \rightarrow 0} \int_{\Omega} \varphi \sigma_{\varepsilon} \cdot \nabla u_{\varepsilon} \geq \int_{\Omega} \varphi a_0(\nabla u^*) \cdot \nabla u^*, \quad (3.22)$$

$\forall \varphi \in \mathcal{D}(\Omega)$ such that $0 \leq \varphi \leq 1$.

Step 5. Finally, we claim that u^* is the unique solution u_0 of (1.9).

Let $v_0 \in K_0$ and since $K_{\varepsilon} \rightarrow K_0$ in the Mosco sense, then there is a sequence $\bar{v}_{\varepsilon} \in K_{\varepsilon}$ such that $\bar{v}_{\varepsilon} \rightarrow v_0$ in $W_0^{1,\alpha}(\Omega)$. Using \bar{v}_{ε} as a test function in (1.8) for $\varphi \in \mathcal{D}(\Omega)$, $0 \leq \varphi \leq 1$, one gets

$$\int_{\Omega} \sigma_{\varepsilon} \cdot \nabla \bar{v}_{\varepsilon} - \int_{\Omega} f(\bar{v}_{\varepsilon} - u_{\varepsilon}) \geq \int_{\Omega} \sigma_{\varepsilon} \cdot \nabla u_{\varepsilon} \geq \int_{\Omega} \varphi (\sigma_{\varepsilon} \cdot \nabla u_{\varepsilon}). \quad (3.23)$$

Recalling (3.22) and passing to the limit in ε in (3.23), we obtain

$$\int_{\Omega} a_0(\nabla u^*) \cdot \nabla v_0 - \int_{\Omega} f(v_0 - u^*) \geq \int_{\Omega} \varphi a_0(\nabla u^*) \cdot \nabla u^*.$$

Letting $\varphi \rightarrow 1$ in the last inequality, one gets

$$\int_{\Omega} a_0(\nabla u^*) \cdot \nabla(v_0 - u^*) - \int_{\Omega} f(v_0 - u^*) \geq 0, \quad \forall v_0 \in K_0.$$

The latter, combined with (3.6), allow us to conclude that u^* coincides with the unique solution u_0 of (1.9) and the whole sequence $u_{\varepsilon} \rightharpoonup u_0$ in $W_0^{1,\alpha}(\Omega)$. \blacksquare

Remark 3.1. *One can also show the convergence of the energies. More precisely,*

$$\int_{\Omega} a_{\varepsilon}(x, \nabla u_{\varepsilon}) \cdot \nabla u_{\varepsilon} dx \rightarrow \int_{\Omega} a_0(\nabla u_0) \cdot \nabla u_0 dx. \quad (3.24)$$

Proof: For any $\varphi \in \mathcal{D}(\Omega)$ such that $0 \leq \varphi \leq 1$ from (3.14) we have

$$\int_{\Omega} \varphi \sigma_{\varepsilon} \cdot \nabla u_{\varepsilon} \leq \int_{\Omega} \sigma_{\varepsilon} \cdot \nabla(\varphi \bar{u}_{\varepsilon}) - \int_{\Omega} u_{\varepsilon} \sigma_{\varepsilon} \cdot \nabla \varphi - \int_{\Omega} f \varphi (\bar{u}_{\varepsilon} - u_{\varepsilon}),$$

which gives

$$\limsup_{\varepsilon \rightarrow 0} \int_{\Omega} \varphi \sigma_{\varepsilon} \cdot \nabla u_{\varepsilon} \leq \int_{\Omega} a_0(\nabla u_0) \cdot \nabla u_0. \quad (3.25)$$

The latter, combined with (3.22), implies

$$\sigma_{\varepsilon} \cdot \nabla u_{\varepsilon} \rightarrow a_0(\nabla u_0) \cdot \nabla u_0 \quad \text{in } \mathcal{D}'(\Omega).$$

Since $K_{\varepsilon} \rightarrow K_0$ in the Mosco sense, then taking $v_0 = u_0$ in (3.23), we get

$$\begin{aligned} \int_{\Omega} a_0(\nabla u_0) \cdot \nabla u_0 &\geq \limsup_{\varepsilon \rightarrow 0} \int_{\Omega} \sigma_{\varepsilon} \cdot \nabla u_{\varepsilon} \\ &\geq \liminf_{\varepsilon \rightarrow 0} \int_{\Omega} \sigma_{\varepsilon} \cdot \nabla u_{\varepsilon} \\ &\geq \int_{\Omega} \varphi a_0(\nabla u_0) \cdot \nabla u_0, \end{aligned}$$

and letting $\varphi \rightarrow 1$, we obtain (3.24). ■

Remark 3.2. *If in (1.8) we have f_{ε} instead of f and $f_{\varepsilon} \rightharpoonup f$ in $L^s(\Omega)$, then the conclusion of the Theorem 3.1 still holds.*

Remark 3.3. *Since there are Lewy-Stampacchia inequalities also for the two obstacles problem (see [19]), the Theorem 3.1 can be extended for two obstacles problems with similar assumptions.*

4. Convergence of the coincidence sets

In this section, using the Lewy-Stampacchia inequalities, we prove a stability result for the coincidence sets as it was done, for example, in Theorem 6:6.1 in [18].

Theorem 4.1. *Let the conditions of Theorem 3.1 hold. If, as $\varepsilon \rightarrow 0$,*

$$u_{\varepsilon} - \psi_{\varepsilon} \rightarrow u_0 - \psi_0 \quad \text{in } L^1(\Omega), \quad (4.1)$$

$$(A_{\varepsilon} \psi_{\varepsilon} - f)^+ \rightarrow (A_0 \psi_0 - f)^+ \quad \text{in } L^1(\Omega), \quad (4.2)$$

$$A_{\varepsilon} u_{\varepsilon} \rightarrow A_0 u_0 \quad \text{in } \mathcal{D}'(\Omega), \quad (4.3)$$

$$\int_S d(A_0 \psi_0 - f) \neq 0, \quad \forall S \subset \Omega \quad \text{such that } |S| > 0, \quad (4.4)$$

and

$$A_0 u_0 - f = (A_0 \psi_0 - f) \chi_0 \quad \text{a.e. in } \Omega, \quad (4.5)$$

where χ_0 is the characteristic function of the set $I_0 := \{u_0 = \psi_0\}$, then the coincidence sets $I_\varepsilon := \{u_\varepsilon = \psi_\varepsilon\}$ converge in measure, i.e.,

$$\chi_\varepsilon \rightarrow \chi_0 \quad \text{in } L^p(\Omega), \forall p \in [1, \infty),$$

where χ_ε is the characteristic function of I_ε .

Proof: From the Lewy-Stampacchia inequalities we have

$$f \leq A_\varepsilon u_\varepsilon \leq f + (A_\varepsilon \psi_\varepsilon - f)^+ \quad \text{a.e. in } \Omega.$$

Hence, there exists a function $q_\varepsilon \in L^\infty(\Omega)$, such that,

$$A_\varepsilon u_\varepsilon - f = q_\varepsilon (A_\varepsilon \psi_\varepsilon - f)^+ \quad \text{a.e. in } \Omega, \quad (4.6)$$

and

$$0 \leq q_\varepsilon \leq \chi_\varepsilon \leq 1 \quad \text{a.e. in } \Omega. \quad (4.7)$$

Then for a subsequence (still denoted by ε), one has

$$q_\varepsilon \rightarrow q \quad \text{and} \quad \chi_\varepsilon \rightharpoonup \chi_* \quad \text{in } L^\infty(\Omega) - \text{weak}^* \quad (4.8)$$

for functions $q, \chi_* \in L^\infty(\Omega)$. The inequalities (4.7) imply

$$0 \leq q \leq \chi_* \leq 1 \quad \text{a.e. in } \Omega. \quad (4.9)$$

Using (4.2), (4.3) and (4.8), we pass to the limit, as $\varepsilon \rightarrow 0$, in (4.6) and obtain

$$A_0 u_0 - f = q(A_0 \psi_0 - f)^+ \quad \text{a.e. in } \Omega.$$

The latter, combined with (4.5) provides

$$q(A_0 \psi_0 - f)^+ = (A_0 \psi_0 - f) \chi_0 \quad \text{a.e. in } \Omega. \quad (4.10)$$

Note that in the region $\{A_0 \psi_0 > f\}$, (4.10) and (4.4) imply that $q = \chi_0$, while in $\{A_0 \psi_0 \leq f\}$, $\chi_0 = 0$. Therefore, $q \geq \chi_0$ a.e. in Ω . Consequently, from (4.9) we get

$$\chi_0 \leq \chi_* \quad \text{a.e. in } \Omega.$$

On the other hand, from (4.1) and (4.8) one has

$$0 = \int_{\Omega} \chi_\varepsilon (u_\varepsilon - \psi_\varepsilon) \rightarrow \int_{\Omega} \chi_* (u_0 - \psi_0) = 0,$$

thus $\chi_*(u_0 - \psi_0) = 0$ a.e. in Ω . Consequently, if $u_0 > \psi_0$, then $\chi_* = 0$, and since $0 \leq \chi_* \leq 1$, one obtains

$$\chi_0 \geq \chi_* \quad \text{a.e. in } \Omega.$$

Therefore, $\chi_0 = \chi_*$, and the whole sequence χ_ε converges to χ_0 as $\varepsilon \rightarrow 0$, first weakly, and since they are characteristic functions, also strongly in any $L^p(\Omega)$, for any $p \in [1, \infty)$. ■

Remark 4.1. *If $\psi_0 = 0$ and the right hand side is regular enough, the condition (4.5) holds automatically, since in this particular case one has porosity of the free boundary from [9] (hence, the free boundary has Lebesgue measure zero), which provides (4.5).*

Remark 4.2. *The assumption (4.4) is a weaker version of the condition*

$$A_0\psi_0 - f \neq 0 \quad \text{a.e. in } \Omega, \quad \text{when } A_0\psi_0 \in L^1(\Omega).$$

Theorem 4.2. *Let the conditions of Theorem 3.1 and also $s > n/2$. If $\psi_\varepsilon \rightarrow \psi_0$, uniformly, $\psi_0|_{\partial\Omega} < 0$ and*

$$\overline{\text{int}\{u_0 = \psi_0\}} = \{u_0 = \psi_0\} = I_0,$$

then the coincidence sets $I_\varepsilon := \{u_\varepsilon = \psi_\varepsilon\}$ converge in the Hausdorff distance to I_0 .

Proof: Using [13, Theorem 3.2], we obtain the uniform Hölder continuity of solutions. The uniform Hölder continuity of the obstacles then implies, as $\varepsilon \rightarrow 0$, the convergence $u_\varepsilon \rightarrow u_0$, uniformly in compact subsets of Ω . This, in turn, provides the convergence of the coincidence sets in Hausdorff distance as in [8] and [18, Theorem 6:6.5]. ■

References

- [1] H. Attouch and C. Picard, *Variational inequalities with varying obstacles. The general form of the limit problem*, J. Funct. Anal. 50 (1983), 329-386.
- [2] L. Boccardo, *Lewy-Stampacchia inequality in quasilinear unilateral problems and applications to the G-convergence*, Boll. Unione Mat. Ital. (9) 4 (2011), 275-282.
- [3] L. Boccardo and I.C. Dolcetta, *G-convergenza e problema di Dirichlet unilaterale*, Boll. Unione Mat. Ital. 4 (1975), 115-123.
- [4] L. Boccardo and F. Murat, *Nouveaux résultats de convergence dans des problèmes unilatéraux*, Research Notes in Math. 60 (1982), 64-85.
- [5] L. Boccardo and F. Murat, *Homogenization of nonlinear unilateral problems*, Composite media and homogenization theory; an international center for theoretical physics workshop, Trieste, Italy, January 1990 (1991), 81-105.
- [6] D. Cioranescu and P. Donato, *An introduction to homogenization*, Oxford lecture series in Mathematics and its applications 17, Oxford University Press Inc., New York, 1999.
- [7] V. Chiado Piat, G. Dal Maso and A. Defranceschi, *G-convergence of monotone operators*, Ann. Inst. Henri Poincaré 7 (1990), 123-160.

- [8] M. Codegone and J.F. Rodrigues, *Convergence of the coincidence set in the homogenization of the obstacle problem*, Ann. Fac. Sci. Toulouse Math. (5) 3 (1981), 275-285.
- [9] S. Challal, A. Lyaghfour, J.F. Rodrigues and R. Teymurazyan, *On the regularity of the free boundary for quasilinear obstacle problems*, Interfaces Free Bound. 16 (2014), 359-394.
- [10] L. Diening, P. Harjulehto, P. Hästö and M. Ruzicka, *Lebesgue and Sobolev spaces with variable exponents*, Lecture Notes in Mathematics 2017 (2011).
- [11] L.C. Evans, *Weak convergence methods for nonlinear partial differential equations*, American Mathematical Society, 1990.
- [12] X. Fan and Q. Zhang, *Existence of solutions for $p(x)$ -Laplacian Dirichlet problem*, Nonlinear Anal., Theory Methods Appl. 52 (2003), 1843-1852.
- [13] X. Fan and D. Zhao, *The quasi-minimizer of integral functionals with $m(x)$ growth condition*, Nonlinear Anal. 39 (2000), 807-816.
- [14] X. Fan and D. Zhao, *On the spaces $L^{p(x)}(\Omega)$ and $W^{m,p(x)}(\Omega)$* , J. Math. Anal. Appl. 263 (2001), 424-446.
- [15] D. Kinderlehrer and G. Stampacchia, *An introduction to variational inequalities and their applications*, SIAM Classics in Applied Mathematics 31, 2000, xxii+306p.
- [16] F. Murat, *Compacité par compensation*, Ann. Scuola Norm. Sup. Pisa. Cl. Sci (4) 5 (1978), 489-507.
- [17] A.A. Pankov, *G-convergence and homogenization of nonlinear partial differential operators*, Kluwer Academic Publishers, Dordrecht, 1997.
- [18] J.F. Rodrigues, *Obstacle problems in mathematical physics*, North-Holland Mathematics Studies 134 (Notas de Matematica 114), Elsevier Science Publishers B.V., 1991.
- [19] J.F. Rodrigues and R. Teymurazyan, *On the two obstacles problem in Orlicz-Sobolev spaces and applications*, Complex Var. Elliptic Equ. 56 (2011), 769-787.
- [20] G.V. Sandrakov, *Homogenization of variational inequalities for obstacle problems*, Sbornik, Mathematics 196 (2005), 541-560.
- [21] G.V. Sandrakov, *Homogenization of nonlinear equations and variational inequalities with obstacles*, Doklady Mathematics 73 (2006), 178-181.
- [22] V.V. Zhikov, *Laurent'ev effect and the homogenization of nonlinear variational problems*, Differ. Equations 27 (1991), 32-39.
- [23] V.V. Zhikov and S.E. Pastukhova, *On the compensated compactness principle*, Doklady Mathematics 82 (2010), 590-595.
- [24] V.V. Zhikov and S.E. Pastukhova, *Homogenization of monotone operators under conditions of coercivity and growth of variable order*, Mathematical Notes 90 (2011), 48-63.

DIEGO MARCON

DEPARTMENT OF MATHEMATICS, FEDERAL UNIVERSITY OF RIO GRANDE DO SUL, AV. BENTO GONCALVES 9500, 91509-900 PORTO ALEGRE, RS, BRAZIL.

E-mail address: diego.marcon@ufrgs.br

JOSÉ FRANCISCO RODRIGUES

CMAFcIO, DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCES, UNIVERSITY OF LISBON, PORTUGAL.

E-mail address: jfrodrigues@ciencias.ulisboa.pt

RAFAYEL TEYMURAZYAN

CMUC, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF COIMBRA, 3001-501 COIMBRA, PORTUGAL

E-mail address: rafayel@utexas.edu