

EXTREMAL CURVES ON STIEFEL AND GRASSMANN MANIFOLDS

VELIMIR JURDJEVIC, IRINA MARKINA AND FÁTIMA SILVA LEITE

ABSTRACT: This paper uncovers a large class of left-invariant sub-Riemannian systems on Lie groups that admit explicit solutions with certain properties, and provides geometric origins for a class of important curves on Stiefel manifolds, called quasi-geodesics, that project on Grassmann manifolds as Riemannian geodesics. We show that quasi-geodesics are the projections of sub-Riemannian geodesics generated by certain left-invariant distributions on Lie groups that act transitively on each Stiefel manifold $\text{St}_k^n(V)$. This result is valid not only for the real Stiefel manifolds in $V = \mathbb{R}^n$, but also for the Stiefel in the Hermitian space $V = \mathbb{C}^n$ and the quaternion space $V = \mathbb{H}^n$.

KEYWORDS: Sub-Riemannian geometry, Riemannian and sub-Riemannian geodesics, quasi-geodesic curves, horizontal distributions, Grassmann manifolds, Stiefel manifolds, Lie groups actions on manifolds, Pontryagin Maximum Principle, extremal curves.

AMS SUBJECT CLASSIFICATION (1991): Primary 53C17, 53C22, 53B21, 53C25, 30C80; Secondary 49J15, 58E40.

1. Introduction

This paper provides geometric origins for a class of curves on Stiefel manifolds, called quasi-geodesic, that have proved to be particularly important in solving interpolation problems arising in real applications [12]. We show that quasi-geodesic curves are the projections of sub-Riemannian geodesics generated by certain left-invariant distributions on Lie groups G that act on Stiefel manifolds. This quest for the geometric characterization of quasi-geodesic curves uncovered a large class of left-invariant sub-Riemannian systems on Lie groups that admit explicit solutions, in the form that will be made clear below. As a result, the paper is as much about sub-Riemannian structures on Lie groups as it is about quasi-geodesic curves.

The first part of the paper deals with sub-Riemannian structures associated with homogeneous spaces $M = G/K$ induced by a transitive left action of a semi-simple Lie group G on a smooth manifold M , where K denotes the isotropy subgroup relative to a fixed point $m_0 \in M$. The sub-Riemannian structures will be defined by a left-invariant distribution \mathcal{H} generated by a

vector space $\mathfrak{p} \subset \mathfrak{g}$ that is transversal to the Lie algebra \mathfrak{k} of the isotropy group K , and satisfies additional Lie algebraic relations:

$$\mathfrak{g} = \mathfrak{p} \oplus \mathfrak{k}, \quad [\mathfrak{p}, \mathfrak{k}] \subseteq \mathfrak{p}, \quad \mathfrak{k} \subseteq [\mathfrak{p}, \mathfrak{p}].$$

The sub-Riemannian metrics are defined by bilinear, symmetric, non-degenerate, Ad_G invariant forms $\langle \cdot, \cdot \rangle$ that are positive definite on \mathfrak{p} . Under additional assumption that \mathfrak{p}^\perp , the orthogonal complement relative to the Killing form, is a Lie subalgebra of \mathfrak{g} , we show that the sub-Riemannian geodesics are of the form

$$g(t) = g_0 e^{t(P+Q)} e^{-tQ},$$

where $P \in \mathfrak{p}$ and $Q \in \mathfrak{p}^\perp$, see Theorem 1.

In these situations, a Riemannian metric on M is induced by the push forward of the sub-Riemannian metric on G . We show that the corresponding Riemannian geodesics on M are the projections of curves in (1) where P and Q satisfy the additional relation that $P + Q$ is in \mathfrak{k}^\perp . The above findings coincide with the analogous results on semi-simple Lie groups with an involutive automorphism, as in the theory of symmetric Riemannian spaces, where $\mathfrak{p}^\perp = \mathfrak{k}$, see, for instance, [10].

The second part of the paper deals with the application of these results to the actions of Lie groups on Stiefel and Grassmann manifolds and their relevance to the quasi-geodesic curves. Rather than dealing exclusively with the real case, we also include Stiefel and Grassmann manifolds over complex and quaternion algebras.

We realize each Stiefel manifold in two ways as homogeneous manifolds G/K . On groups G that define the homogeneous structure we consider three distinct sub-Riemannian structures whose sub-Riemannian geodesics are described by Theorem 1. We then single out the sub-classes of sub-Riemannian geodesics that project either to Riemannian geodesics or to quasi-geodesic curves on the Stiefel manifolds. Additionally, we show that quasi-geodesics are curves of constant curvature relative to the induced Riemannian metric. In the process we discovered an interesting fact that two distinct sub-Riemannian structures on G can induce isometric Riemannian structures on the homogeneous manifold G/K , see Theorem 2.

We end the paper with a brief discussion of Lagrangian manifolds. First, we show the relation of Theorem 1 to the canonical (symmetric) Riemannian

metric, and secondly, we show that the projections of curves on Stiefel manifolds, descending from the sub-Riemannian curves on G , have constant geodesic curvature on the Grassmann manifold relative to its canonical metric. In particular, we show that the quasi-geodesic curves on the Stiefel manifolds project onto the Riemannian geodesics on the Grassmannian manifold.

This article constitutes a full version of the results announced in a short paper [11] on Stiefel manifolds embedded in the Euclidean space \mathbb{R}^n .

2. Group actions and sub-Riemannian problems

2.1. Notations and background material

We begin with the notations and basic concepts that will be necessary throughout the paper. Details about this background material may be found, for instance, in [9, 13, 16, 17].

In particular, M will denote a smooth (or real analytic) manifold, TM and T^*M denote the tangent and the cotangent bundle of M respectively, and T_mM and T_m^*M denote the tangent and cotangent spaces at $m \in M$. If F is a smooth map between smooth manifolds, then F_* will denote the tangent map and F^* the dual map.

The set of smooth vector fields on M is denoted by $\Gamma(TM)$. A smooth curve $m(t)$ defined on an open interval $(-\epsilon, \epsilon)$ in M is said to be an integral curve, or a solution curve, of X if $\frac{dm}{dt} = X(m(t))$ for all $t \in (-\epsilon, \epsilon)$. A vector field X is said to be complete if each integral curve of X can be extended to the interval $(-\infty, \infty)$. We denote by $\phi_t^X(m)$ the flow on M generated by a complete vector field $X \in \Gamma(TM)$. Sometimes we will regard a flow as a one parameter group of diffeomorphisms $\{\phi_t^X : t \in \mathbb{R}\}$. We will make use of the fact that any one parameter group of diffeomorphisms ϕ_t on M is generated by the flow of a complete vector field X , in the sense that $\phi_t = \phi_t^X$. In this context, X is called the infinitesimal generator of the group of diffeomorphisms.

Throughout the paper G will denote a Lie group, and \mathfrak{g} will denote its Lie algebra. We think of \mathfrak{g} as the tangent space T_eG at the identity $e \in G$, with the Lie bracket induced by the left-invariant vector fields in G : $[A, B] = [X, Y](e) = (YX - XY)(e)$, where $X(g) = (L_g)_*(A)$ and $Y(g) = (L_g)_*(B)$, $A, B \in \mathfrak{g}$, and L_g is the left translation by g in G . We will adopt a short-hand notation and write gA (Ag) for the left (right) invariant vector fields.

If $X(g) = gA$ is a left (Ag right) invariant vector field defined by $A \in \mathfrak{g}$, then its flow $\phi_t^X(g)$ is the left (right) translate $L_g\phi_t^X(e)$ ($R_g\phi_t^X(e)$) of the

flow through the group identity. We will write $\exp(tA)$, as well as e^{tA} when convenient, for the curve $\phi_t^X(e)$ with $X(g) = gA$.

A Lie group G is said to act on a manifold M through the left action $\phi: G \times M \rightarrow M$ if ϕ satisfies

$$\phi(g, \phi(h, m)) = \phi(L_g(h), m) = \phi(gh, m) \quad \text{and} \quad \phi(e, m) = m,$$

for all g, h in G , and all m in M . We use ϕ_g to denote the diffeomorphism $m \mapsto \phi_g(m)$ on M . If m_0 is a point in M , then $K_{m_0} = \{g \in G : \phi(g, m_0) = m_0\}$ denotes the isotropy group of m_0 . An isotropy group is a closed subgroup of G and any two isotropy groups are conjugate. If a group G acts transitively on M , then M can be regarded as the quotient G/K_{m_0} with $m \in M$ identified with gK_{m_0} whenever $\phi_g(m_0) = m$. Then the natural projection $\pi: G \rightarrow G/K_{m_0}$ is given by the map $g \rightarrow \phi_g(m_0)$.

Each element $A \in \mathfrak{g}$ induces a one-parameter group of diffeomorphisms $\{\phi_{e^{tA}} : t \in \mathbb{R}\}$ on M . We will use X_A to denote its infinitesimal generator, so that $\phi_{e^{tA}}(m) = \phi_t^{X_A}(m)$. We will refer to $\mathcal{F} = \{X_A : A \in \mathfrak{g}\}$ as the family of vector fields on M subordinated to the group action. Since G acts by the left action on M , \mathcal{F} is a homomorphic image of the family of right-invariant vector fields on G . Therefore, \mathcal{F} is a finite dimensional Lie algebra of complete vector fields on M .

If G is connected, then any $g \in G$ can be written as $g = e^{t_k A_k} \dots e^{t_1 A_1}$ for some elements A_1, \dots, A_k in \mathfrak{g} . Then,

$$\phi_g(m) = \phi_{e^{t_k A_k} \dots e^{t_1 A_1}}(m) = \phi_{t_k}^{X_{A_k}} \dots \phi_{t_1}^{X_{A_1}}(m). \quad (1)$$

This implies that M is equal to the orbit of \mathcal{F} through any point $m \in M$ whenever G is connected, and the action of G on M is transitive. In such a case any absolutely continuous curve $m(t)$ in M is a solution of

$$\frac{dm}{dt} = \sum_{i=1}^k u_i(t) X_{A_i}(m)$$

for some choice of elements A_1, \dots, A_k in \mathfrak{g} and some choice of measurable and bounded functions $u_1(t), \dots, u_k(t)$.

2.2. Sub-Riemannian problems on Lie groups

In what follows we will always suppose that G is a connected Lie group that acts transitively on M from the left, and K will always denote the isotropy subgroup of a fixed point m_0 in M . Then \mathfrak{k} will denote the Lie algebra of K .

We will now assume that \mathfrak{p} is a linear subspace in \mathfrak{g} such that

$$\mathfrak{g} = \mathfrak{p} \oplus \mathfrak{k}, \quad [\mathfrak{p}, \mathfrak{k}] \subseteq \mathfrak{p}, \quad \mathfrak{k} \subseteq [\mathfrak{p}, \mathfrak{p}]. \quad (2)$$

The space \mathfrak{p} induces a family $\mathcal{H}_{\mathfrak{p}}$ of left-invariant vector fields $X_A(g) = gA, g \in G, A \in \mathfrak{p}$. The associated distribution $\mathcal{H}_{\mathfrak{p}}(g), g \in G$, will be called horizontal, or the Ehresmann \mathfrak{p} connection, see for instance [13]. Left-invariant vector fields $gB, B \in \mathfrak{k}$ will be called vertical. Then \mathcal{V} will denote the family of vertical vector fields.

Absolutely continuous curves $g(t) \in G$ that satisfy $\frac{dg}{dt} = \dot{g}(t) \in \mathcal{H}_{\mathfrak{p}}(g(t))$ for almost all t in some interval $[0, T]$ are called *horizontal*. Alternatively, horizontal curves can be described as the solution curves of a control system

$$\frac{dg}{dt} = \sum_{i=1}^k u_i(t)gA_i,$$

where A_1, \dots, A_k is a basis for \mathfrak{p} , and $u_1(t), \dots, u_k(t)$ are bounded and measurable functions on $[0, T]$.

Conditions (2) imply that $\mathfrak{p} + [\mathfrak{p}, \mathfrak{p}] = \mathfrak{g}$, which in turn implies that $\mathcal{H}_{\mathfrak{p}}$ is a two step bracket generating distribution, in the sense that

$$\mathcal{H}_{\mathfrak{p}}(g) + [\mathcal{H}_{\mathfrak{p}}, \mathcal{H}_{\mathfrak{p}}](g) = T_g G, \quad g \in G.$$

Therefore, any two points in G can be joined by a horizontal curve whenever G is connected [2, 5, 15].

A curve $g(t)$ in G is called a *horizontal lift of a curve* $m(t) \in M = G/K$ if $g(t)$ is a horizontal curve that projects onto $m(t)$, that is, $\pi(g(t)) = \phi_{g(t)}(m_0) = m(t)$.

Proposition 1. *Every absolutely continuous curve $m(t)$ in M is the projection of a horizontal curve. Moreover, if $g_1(t)$ and $g_2(t)$ are horizontal lifts of a curve $m(t)$ in M , then $g_1(t) = g_2(t)h$ for some constant element $h \in K$.*

Proof: Equation (1) implies that every absolutely continuous curve $m(t)$ in M is the projection of an absolutely continuous curve $g(t) \in G$. Then $\dot{g}(t) = g(t)U(t)$ for some curve $U(t) \in \mathfrak{g}$, and $U(t) = P(t) + Q(t)$, with $P(t) \in \mathfrak{p}$ and $Q(t) \in \mathfrak{k}$.

Now define a new curve $\tilde{g}(t) = g(t)h(t)$, where $h(t)$ is a solution in K of $\dot{h}(t) = -Q(t)h(t)$. The curve $\tilde{g}(t)$ also projects on $m(t)$ and furthermore,

$$\dot{\tilde{g}}(t) = g(t)(P(t) + Q(t))h(t) + g(t)\dot{h}(t) = \tilde{g}(t)h^{-1}(t)P(t)h(t).$$

The condition $[\mathfrak{p}, \mathfrak{k}] \subseteq \mathfrak{p}$ implies that $h^{-1}(t)P(t)h(t)$ is in \mathfrak{p} for all t , hence $\tilde{g}(t)$ is a horizontal curve that projects onto $m(t)$.

If $g_1(t)$ and $g_2(t)$ are horizontal lifts of a curve $m(t)$ in M , then $\dot{g}_1 = gU_1(t)$ and $\dot{g}_2 = g_2U_2(t)$ for some curves $U_1(t)$ and $U_2(t)$ in \mathfrak{p} . Since $g_1(t)$ and $g_2(t)$ project onto the same curve $m(t)$, $g_1(t) = g_2(t)h(t)$ for some curve $h(t) \in K$. But then,

$$g_1U_1(t) = \dot{g}_1 = \dot{g}_2h + g_2\dot{h} = g_1(h^{-1}U_2h + h^{-1}\dot{h}).$$

Hence, $U_1 = h^{-1}U_2h + h^{-1}\dot{h}$. Since $h^{-1}U_2h$ is in \mathfrak{p} , $h^{-1}\dot{h} = 0$. Therefore h is constant. \blacksquare

Assume now that $\langle \cdot, \cdot \rangle_{\mathfrak{p}}$ is any positive definite Ad_K -invariant symmetric bilinear form on \mathfrak{p} . This bilinear form induces a left-invariant inner product $\langle gA, gB \rangle_g = \langle A, B \rangle_{\mathfrak{p}}$, on $\mathcal{H}_{\mathfrak{p}}(g)$. We define $\|\frac{dg}{dt}\|_{\mathfrak{p}} = \sqrt{\langle g^{-1}(t)\frac{dg}{dt}, g^{-1}(t)\frac{dg}{dt} \rangle_{\mathfrak{p}}}$ for any horizontal curve $g(t)$. This metric is called a (left-invariant) *sub-Riemannian metric* relative to the distribution $\mathcal{H}_{\mathfrak{p}}$. We denote it by the same symbol $\langle \cdot, \cdot \rangle_{\mathfrak{p}}$. The pair $(\mathcal{H}_{\mathfrak{p}}, \langle \cdot, \cdot \rangle_{\mathfrak{p}})$ is called a (left-invariant) *sub-Riemannian structure* on G and the triplet $(G, \mathcal{H}_{\mathfrak{p}}, \langle \cdot, \cdot \rangle_{\mathfrak{p}})$ is called a *sub-Riemannian manifold*, see for instance [3, 4]. The sub-Riemannian metric $\langle \cdot, \cdot \rangle_{\mathfrak{p}}$ induces a length $l(g, T)$ of a horizontal curve $g: [0, T] \rightarrow G$, by

$$l(g, T) = \int_0^T \langle \dot{g}(t), \dot{g}(t) \rangle_{\mathfrak{p}}^{1/2} dt = \int_0^T \langle g^{-1}(t)\dot{g}(t), g^{-1}(t)\dot{g}(t) \rangle_{\mathfrak{p}}^{1/2} dt.$$

The sub-Riemannian distance function d_{sR} on G is defined by

$$d_{sR}(g_1, g_2) = \inf \left\{ l(g, T) : \dot{g} \in \mathcal{H}_{\mathfrak{p}}(g), \quad g(0) = g_1, \quad g(T) = g_2 \right\}.$$

Definition 1. A horizontal curve $g: [0, T] \rightarrow G$ is called *geodesic* if for any $t \in (0, T)$ there exists $\varepsilon > 0$ such that

$$d_{sR}(g(t_1), g(t_2)) = \int_{t_1}^{t_2} \langle \dot{g}(t), \dot{g}(t) \rangle_{\mathfrak{p}}^{1/2} dt.$$

for any t_1, t_2 with $t - \varepsilon < t_1 < t_2 < t + \varepsilon$.

Any Ad_K left-invariant sub-Riemannian metric $\langle \cdot, \cdot \rangle_{\mathfrak{p}}$ induces a Riemannian metric on $M = G/K$, whereby the length of a curve $m(t)$ on an interval $[0, T]$ is equal to the length of a horizontal lift $g(t)$ of $m(t)$ in G . That is, the length of $m(t)$ is given by $l(m, T) = \int_0^T \sqrt{\langle U(t), U(t) \rangle_{\mathfrak{p}}} dt$, where $U(t) = g^{-1}(t)\dot{g}(t)$. If \tilde{g} is another horizontal lift of $m(t)$ then, according to the

previous proposition, $\tilde{g} = gh$ for some constant $h \in K$, and $\tilde{U}(t) = h^{-1}U(t)h$. But then $\int_0^T \sqrt{\langle U(t), U(t) \rangle_{\mathfrak{p}}} dt = \int_0^T \sqrt{\langle \tilde{U}(t), \tilde{U}(t) \rangle_{\mathfrak{p}}} dt$, by Ad_K -invariance of the metric. Hence the length of $m(t)$ is well defined.

Definition 2. *A Riemannian metric on $M = G/K$ that is the push forward of a sub-Riemannian left-invariant metric $\langle \cdot, \cdot \rangle_{\mathfrak{p}}$ on a horizontal distribution $\mathcal{H}_{\mathfrak{p}}$ in G will be called homogeneous.*

In the present paper we are essentially interested in the structure of sub-Riemannian geodesics on G and their relation to the Riemannian geodesics on $M = G/K$ relative to the homogeneous metric. Our fundamental results will be extracted through the length minimizing property of the geodesics and the following auxiliary optimal problem on G :

given any two points g_1 and g_2 in G , find a horizontal curve of shortest length that connects g_1 to g_2 .

The solutions to this auxiliary problem are intimately related to the sub-Riemannian geodesics, because any horizontal curve of shortest length that connects g_1 to g_2 is a sub-Riemannian geodesic. Conversely, every sub-Riemannian geodesic $g(t)$ is a curve of shortest length relative to the points $g_1 = g(0)$ and $g_2 = g(t_1)$ for sufficiently small t_1 .

This formulation permits easy comparison between the sub-Riemannian geodesics in G and the Riemannian geodesics in $M = G/K$ via the following proposition.

Proposition 2. *Suppose that m_1 and m_2 are given points in M . Let $S_1 = \pi^{-1}(m_1)$ and $S_2 = \pi^{-1}(m_2)$ be the fibers above these points. Then, the projection of a horizontal curve $g(t)$ is a curve of minimal length in M that connects m_1 to m_2 if and only if $g(t)$ is the curve of minimal length that connects S_1 to S_2 .*

The proof is simple and will be omitted.

2.3. The associated optimal control problem

The optimal problem of finding curves of shortest length that connect two given points can be easily formulated as a time-optimal control problem, but since it is more convenient to work with the energy functional $E = \frac{1}{2} \int_0^T \|U(t)\|_{\mathfrak{p}}^2 dt$ rather than the length functional $\int_0^T \|U(t)\|_{\mathfrak{p}} dt$, we will, instead, pass to the following energy-optimal control problem on G :

if $T > 0$ is a fixed number, and if g_1 and g_2 are fixed points in G , find a horizontal curve $g(t)$ that satisfies $g(0) = g_1$, $g(T) = g_2$, along which the total energy $E = \frac{1}{2} \int_0^T \|U(t)\|_{\mathfrak{p}}^2 dt$, $U(t) = g^{-1}(t) \frac{dg}{dt}(t)$, is minimal among all the horizontal curves that satisfy the boundary conditions $g(0) = g_1$ and $g(T) = g_2$.

The following proposition clarifies the relation between these two optimal control problems.

Proposition 3. *Every horizontal curve of shortest length is a solution of the optimal control problem for a suitable $T > 0$.*

Proof: If $g(t)$ is any horizontal curve that is a solution of $\frac{dg}{dt} = gU(t)$ that connects $g_1 = g(0)$ to $g_2 = g(T)$, then by the Cauchy inequality

$$\int_0^T \|U(t)\|_{\mathfrak{p}} dt \leq T^{\frac{1}{2}} \left(\int_0^T \|U(t)\|_{\mathfrak{p}}^2 dt \right)^{\frac{1}{2}}.$$

The equality occurs only when $\|U(t)\|_{\mathfrak{p}}$ is constant, that is, when T is proportional to the length of $g(t)$ on $[0, T]$.

In particular, if $g(t)$ is a curve of minimal length, and if $\|U(t)\|_{\mathfrak{p}} = 1$, then T is the length of $g(t)$ on $[0, T]$, and $E = \frac{1}{2}T$ is the minimal value of the energy functional relative to the boundary values g_1 and g_2 . Conversely, suppose that $\hat{g}(t)$ is a horizontal curve parametrized by arc length: $\|\hat{U}(t)\|_{\mathfrak{p}} = 1$, such that $\hat{g}(0) = g_1$ and $\hat{g}(T) = g_2$. Then, by the above inequality, $E = \frac{1}{2}T$ is the minimal value of the energy functional over all horizontal curves that satisfy the boundary conditions $g(0) = g_0$ and $g(T) = g_1$, and $\hat{g}(t)$ attains this minimal value.

Suppose now that $g(s)$ is a horizontal curve of minimal length L such that $g(0) = g_1$ and $g(S) = g_2$, for some $S > 0$. Then $L = \int_0^S \|U(\tau)\|_{\mathfrak{p}} d\tau$. It is not difficult to show that $g(s)$ is a regular curve on the interval $[0, S]$, in the sense that $\frac{dg}{ds}(s) \neq 0$, $s \in [0, S]$. Therefore, $g(s)$ can be reparametrized by a parameter $s(t)$, $t \in [0, T]$ so that the reparametrized curve $\tilde{g}(t) = g(s(t))$ has constant speed $\|\tilde{g}^{-1} \frac{d\tilde{g}}{dt}\|_{\mathfrak{p}} = \lambda$. In fact, $s(t)$ is the inverse function of $t(s) = \frac{1}{\lambda} \int_0^s \|U(\tau)\|_{\mathfrak{p}} d\tau$. Since the property to be horizontal and the length functional are invariant under reparametrizations, $\tilde{g}(t)$ is a horizontal curve of minimal length that reaches g_2 from g_1 in T units of time, by a control of constant magnitude $\lambda = \frac{L}{T}$. Therefore, $\tilde{g}(t)$ is a solution of the optimal control problem on the interval $[0, T]$ relative to the boundary conditions g_1

and g_2 . Hence it attains the optimal value $E = \frac{1}{2}T$. It then follows that

$$L = \lambda T = \int_0^T \|\tilde{U}(t)\|_{\mathfrak{p}} dt = T^{\frac{1}{2}}(2E)^{\frac{1}{2}} = T,$$

and, therefore, $\lambda = 1$ and $L = T$. ■

Proposition 4. *Given any pair of points g_1 and g_2 in G , there exists a horizontal curve $g(t)$ such that $g(0) = g_1$, and $g(T) = g_2$, along which the total energy $E = \frac{1}{2} \int_0^T \|U(t)\|^2 dt$ is minimal.*

The proof is essentially the same as that given in [10, Proposition 9.5, p. 151] and will be omitted.

Corollary 1. *For any two points g_1 and g_2 in G there exists a sub-Riemannian geodesic that connects g_1 to g_2 .*

2.4. The Maximum Principle and the extremal curves

We will now turn to the Maximum Principle of Pontryagin to obtain the necessary conditions of optimality for the above optimal control problem under the additional assumptions that G is semi-simple group, and that $\langle \cdot, \cdot \rangle_{\mathfrak{p}}$ is a positive definite bilinear form on \mathfrak{p} that is the restriction of a symmetric, non-degenerate Ad_G -invariant bilinear form $\langle \cdot, \cdot \rangle$ on \mathfrak{g} . The Ad_G -invariance implies that

$$\langle A, [B, C] \rangle = \langle B, [C, A] \rangle, \quad (3)$$

for all A, B, C in \mathfrak{g} . It also implies that $\langle \cdot, \cdot \rangle_{\mathfrak{p}}$ is Ad_K -invariant. Typically, $\langle \cdot, \cdot \rangle$ could be any scalar multiple of the Killing form $\text{Kl}(A, B) = \text{Tr}(\text{ad}A \circ \text{ad}B)$ that is positive definite on \mathfrak{p} .

To make an easier transition to the literature on control theory, we will represent curves $U(t)$ in \mathfrak{p} in terms of an orthonormal basis A_1, \dots, A_k as $U(t) = \sum_{i=1}^k u_i(t)A_i$. In this representation, horizontal curves are the solutions of $\frac{dg}{dt} = \sum_{i=1}^k u_i(t)gA_i$, and their energy is given by

$$\frac{1}{2} \int_0^T \langle U(t), U(t) \rangle_{\mathfrak{p}} dt = \frac{1}{2} \int_0^T \sum_{i=1}^k u_i^2(t) dt.$$

To take advantage of the left-invariant symmetries, the cotangent bundle T^*G will be represented by $G \times \mathfrak{g}^*$, where \mathfrak{g}^* stands for the dual of \mathfrak{g} . In this representation, points $\xi \in T^*G$ are viewed as the pairs (g, ℓ) , $\ell \in \mathfrak{g}^*$ defined

by $\xi(gA) = \ell(A)$ for any $A \in \mathfrak{g}$. Then the Hamiltonian $h_A(\xi)$ of any left-invariant vector field $X_A(g) = gA$ is given by $h_A(\ell) = \ell(A)$. In particular, the control system $\frac{dg}{dt} = \sum_{i=1}^k u_i(t)gA_i$, together with the associated energy functional, lifts to the extended Hamiltonian

$$h_{U(t)}(\ell) = -\frac{\lambda}{2} \sum_{i=1}^k u_i^2(t) + \sum_{i=1}^k u_i(t)h_i(\ell), \quad \lambda = 1, 0,$$

where $h_i(\ell) = \ell(A_i)$, $i = 1, \dots, k$. The corresponding Hamiltonian vector field $\vec{h}_{U(t)}$ is called the Hamiltonian lift of the energy-extended control system. Its integral curves $(g(t), \ell(t))$ are the solutions of

$$\frac{dg}{dt} = gU(t), \quad \frac{d\ell}{dt} = -\text{ad}^*U(t)(\ell(t)), \quad U(t) = \sum_{i=1}^k u_i(t)A_i,$$

where $(\text{ad}^*U(t)(\ell))(A) = \ell([U(t), A])$, $A \in \mathfrak{g}$, see [10].

According to the Maximum Principle, every optimal solution $g(t)$ generated by a control $U(t)$ is the projection of an integral curve $(g(t), \ell(t))$ of the Hamiltonian vector field $\vec{h}_{U(t)}$ such that

$$h_{U(t)}(\ell(t)) \geq -\frac{\lambda}{2} \sum_{i=1}^k v_i^2 + \sum_{i=1}^k v_i h_i(\ell(t)), \quad (4)$$

for any $(v_1, \dots, v_k) \in \mathbb{R}^k$ and all t . In addition, the Maximum Principle requires that $\ell(t) \neq 0$ when $\lambda = 0$.

Integral curves $(g(t), \ell(t))$ of $\vec{h}_{U(t)}$ that satisfy the conditions of the Maximum Principle (4) are called *extremal*; *abnormal extremal* when $\lambda = 0$, and *normal extremal* when $\lambda = 1$. In the abnormal case, inequality (4) yields constraints

$$h_i(\ell(t)) = 0, \quad i = 1, \dots, k,$$

while in the normal case, the inequality shows that the extremal control $U(t)$ is a critical point of the Hamiltonian $h_{U(t)}(\ell) = -\frac{1}{2} \sum_{i=1}^k u_i^2(t) + \sum_{i=1}^k u_i(t)h_i(\ell)$, that is, the extremal control is of the form $U(t) = \sum_{i=1}^k h_i(\ell(t))A_i$.

The above shows that the normal extremals are the solution curves of a single Hamiltonian vector field corresponding to the Hamiltonian

$$H(\ell) = \frac{1}{2} \sum_{i=1}^k h_i^2(\ell).$$

We will not pursue the abnormal extremals since it is known that their projections on G cannot be optimal when $\mathcal{H}_{\mathfrak{p}}$ is a two step bracket generating distribution (see [2]).

It follows that the sub-Riemannian geodesics are the projections of the integral curves of the Hamiltonian vector field \vec{H} on energy level $H = \frac{1}{2}$, since the energy functional is equal to the length functional only over the horizontal curves parametrized by arc length by Proposition 3. That is, the sub-Riemannian geodesics are the projections of curves $(g(t), \ell(t))$ that are the solutions of

$$\frac{dg}{dt} = g(dH), \quad \frac{d\ell}{dt} = -\text{ad}^*dH(\ell)(\ell(t)), \quad (5)$$

on the energy level set $H = \frac{1}{2}$, where $dH = \sum_{i=1}^k h_i(\ell)A_i$ is the differential of H .

For our purposes, however, it will be more convenient to express equation (5) on the tangent bundle $G \times \mathfrak{g}$ rather than the cotangent bundle $G \times \mathfrak{g}^*$. For that reason, \mathfrak{g}^* will be identified with \mathfrak{g} via the bilinear form $\langle \cdot, \cdot \rangle$, i.e.,

$$\ell \in \mathfrak{g}^* \iff L \in \mathfrak{g} \quad \text{if and only if} \quad \ell(A) = \langle L, A \rangle, \text{ for all } A \in \mathfrak{g}.$$

Then, relying on (3),

$$\begin{aligned} \langle \frac{dL}{dt}, A \rangle &= \frac{d\ell}{dt}(A) = -\text{ad}^*dH(\ell)(\ell(t))(A) = -\ell([dH, A]) \\ &= \langle L, [dH, A] \rangle = \langle [dH, L], A \rangle. \end{aligned}$$

Since A is an arbitrary element of \mathfrak{g} , $\frac{dL}{dt} = [dH, L]$. Hence the extremal equations (5) are equivalent to

$$\frac{dg}{dt} = g(t)dH(\ell(t)), \quad \frac{dL}{dt} = [dH(\ell), L]. \quad (6)$$

Let now \mathfrak{p}^\perp denote the orthogonal complement of \mathfrak{p} in \mathfrak{g} relative to $\langle \cdot, \cdot \rangle$. Since $\langle \cdot, \cdot \rangle$ is a symmetric and non-degenerate quadratic form, $\mathfrak{g} = \mathfrak{p} \oplus \mathfrak{p}^\perp$. Then each $L \in \mathfrak{g}$ can be written as $L = L_{\mathfrak{p}} + L_{\mathfrak{p}^\perp}$ with $L_{\mathfrak{p}} \in \mathfrak{p}$ and $L_{\mathfrak{p}^\perp} \in \mathfrak{p}^\perp$. Relative to the orthonormal basis A_1, \dots, A_k , $L_{\mathfrak{p}} = \sum_{i=1}^k \langle L, A_i \rangle A_i$. But, $\langle L, A_i \rangle = \ell(A_i) = h_i(\ell)$, hence $dH = L_{\mathfrak{p}}$.

The above shows that the Hamiltonian H can be written as $H = \frac{1}{2}\langle L_{\mathfrak{p}}, L_{\mathfrak{p}} \rangle$ and the associated equations (6) can be written as

$$\frac{dg}{dt} = g L_{\mathfrak{p}}, \quad \frac{dL}{dt} = [L_{\mathfrak{p}}, L].$$

Under the previous assumptions, together with the condition that \mathfrak{p}^{\perp} is a Lie subalgebra of \mathfrak{g} , we now come to the main theorem of the paper.

Theorem 1. *Assume that \mathfrak{p}^{\perp} is a Lie subalgebra of \mathfrak{g} . Then, the sub-Riemannian geodesics are given by*

$$g(t) = g(0) \exp(t(P_{\mathfrak{p}} + P_{\mathfrak{p}^{\perp}})) \exp(-tP_{\mathfrak{p}^{\perp}}), \quad (7)$$

for some constant elements $P_{\mathfrak{p}} \in \mathfrak{p}$ and $P_{\mathfrak{p}^{\perp}} \in \mathfrak{p}^{\perp}$ with $\|P_{\mathfrak{p}}\| = 1$.

The Riemannian geodesics on $M = G/K$ are the projections of the sub-Riemannian geodesics for which $P_{\mathfrak{p}} + P_{\mathfrak{p}^{\perp}}$ is orthogonal to \mathfrak{k} .

Proof: Sub-Riemannian geodesics are the projections of

$$\frac{dg}{dt} = g(t)L_{\mathfrak{p}}(t), \quad \frac{dL}{dt} = [L_{\mathfrak{p}}(t), L(t)],$$

on the energy level set $H = \frac{1}{2}\|L_{\mathfrak{p}}\|^2 = \frac{1}{2}$.

We now address the solutions of

$$\frac{dL}{dt} = [L_{\mathfrak{p}}(t), L(t)],$$

or, equivalently, the solutions of

$$\frac{dL_{\mathfrak{p}}}{dt} + \frac{dL_{\mathfrak{p}^{\perp}}}{dt} = [L_{\mathfrak{p}}(t), L(t)] = [L_{\mathfrak{p}}(t), L_{\mathfrak{p}^{\perp}}(t)]. \quad (8)$$

Due to the $\text{ad}_{\mathfrak{k}}$ -invariance property of $\langle \cdot, \cdot \rangle$, and the assumption that \mathfrak{p}^{\perp} is a Lie subalgebra of \mathfrak{g} , we have

$$\langle [L_{\mathfrak{p}}, L_{\mathfrak{p}^{\perp}}], \mathfrak{p}^{\perp} \rangle = \langle L_{\mathfrak{p}}, [L_{\mathfrak{p}^{\perp}}, \mathfrak{p}^{\perp}] \rangle = 0,$$

from which we conclude that $[L_{\mathfrak{p}}(t), L_{\mathfrak{p}^{\perp}}(t)] \in \mathfrak{p}$. Consequently, $\frac{dL_{\mathfrak{p}^{\perp}}}{dt} = 0$, hence $L_{\mathfrak{p}^{\perp}}(t) = P_{\mathfrak{p}^{\perp}}$, for some constant element $P_{\mathfrak{p}^{\perp}} \in \mathfrak{p}^{\perp}$.

So, equation (8) reduces to

$$\frac{dL_{\mathfrak{p}^{\perp}}}{dt} = 0, \quad \frac{dL_{\mathfrak{p}}}{dt} = [L_{\mathfrak{p}}(t), P_{\mathfrak{p}^{\perp}}].$$

Therefore,

$$U(t) = L_{\mathfrak{p}}(t) = \exp(tP_{\mathfrak{p}^{\perp}})P_{\mathfrak{p}} \exp(-tP_{\mathfrak{p}^{\perp}}), \quad (9)$$

where $P_{\mathfrak{p}} = L_{\mathfrak{p}}(0)$.

The corresponding sub-Riemannian geodesics are the solutions of $\frac{dg}{dt} = g(t)U(t)$. In order to show that the solution of this differential equation has the required form, we define $\tilde{g}(t) = g(t) \exp(tP_{\mathfrak{p}^\perp})$ and use (9) to obtain

$$\begin{aligned} \frac{d\tilde{g}}{dt} &= g(t)U(t) \exp(tP_{\mathfrak{p}^\perp}) + \tilde{g}(t)P_{\mathfrak{p}^\perp} \\ &= \tilde{g}(t) \left(\exp(-tP_{\mathfrak{p}^\perp})U(t) \exp(tP_{\mathfrak{p}^\perp}) + P_{\mathfrak{p}^\perp} \right) \\ &= \tilde{g}(t)(P_{\mathfrak{p}} + P_{\mathfrak{p}^\perp}). \end{aligned}$$

Hence, $g(t)$ is given by (7).

To complete the proof, we will use the fact that the geodesics in the quotient space $M = G/K$ are the projections of the extremal curves in G that satisfy the transversality conditions, implied by Proposition 2, which, after the identification $\ell \rightarrow L$, means that $L(t)$, the extremal curve that projects onto a geodesic in G/K , is orthogonal to \mathfrak{k} at $t = 0$ and $t = T$. Since the horizontal distribution $\mathcal{H}_{\mathfrak{p}}$ is K -invariant, the Hamiltonian lift $h_A(L) = \langle L, A \rangle$ of any left-invariant vector field gA , $A \in \mathfrak{k}$ is constant along the solutions of (8). Indeed, $\frac{d}{dt} \langle L(t), A \rangle = \langle [L_{\mathfrak{p}(t)^\perp}, L_{\mathfrak{p}}(t)], A \rangle = \langle L_{\mathfrak{p}(t)^\perp}, [L_{\mathfrak{p}}(t), A] \rangle = 0$, because of $[L_{\mathfrak{p}}(t), A] \in \mathfrak{p}$.

So, since $\langle L(t), A \rangle$ is constant, $\langle L(0), A \rangle = 0$ if and only if $\langle L(t), A \rangle = 0$ for all t . It follows that the orthogonality conditions reduce to $L(0) = P_{\mathfrak{p}} + P_{\mathfrak{p}^\perp} \perp \mathfrak{k}$. \blacksquare

As a corollary we have the following result, [10, Proposition 8.30].

Corollary 2. *If $\mathfrak{p}^\perp = \mathfrak{k}$, the sub-Riemannian geodesics are given by*

$$g(t) = g(0) \exp(t(P_{\mathfrak{p}} + P_{\mathfrak{k}})) \exp(-tP_{\mathfrak{k}}), \quad (10)$$

for some constant elements $P_{\mathfrak{p}} \in \mathfrak{p}$ and $P_{\mathfrak{k}} \in \mathfrak{k}$ with $\|P_{\mathfrak{p}}\| = 1$.

The Riemannian geodesics on $M = G/K$ are the projections of the sub-Riemannian geodesics (10) for which $P_{\mathfrak{k}} = 0$.

Proposition 5. *If $\mathfrak{p}^\perp = \mathfrak{k}$, then the projection of a sub-Riemannian geodesic on the quotient space $M = G/K$ is a curve of constant geodesic curvature relative to the Riemannian metric.*

Proof: Recall that the geodesic curvature of a curve $m(t)$ parametrized by its arc length is equal to the length of the covariant derivative of $\frac{dm}{dt}$ along

$m(t)$. In this context it is most convenient to express the covariant derivative in terms of its horizontal lift as follows.

Let $Y(t)$ denote a vector field defined along the curve $m(t)$ in M . Then $Y(t)$ is the projection of a horizontal curve $g(t)W(t)$, for some curve $W(t) \in \mathfrak{p}$, where $g(t)$ denotes the horizontal curve that projects onto $m(t)$. It follows that $\frac{dm}{dt}$ is the projection of a curve $g(t)U(t)$ for some $U(t) \in \mathfrak{p}$. Then the covariant derivative $\frac{DY}{dt}$ of $Y(t)$ along $m(t)$ is the projection of

$$g(t) \left(\frac{dW}{dt} + \frac{1}{2}[U(t), W(t)]_{\mathfrak{p}} \right),$$

on M , where $[U, W]_{\mathfrak{p}}$ denotes the projection of $[U, W]$ on \mathfrak{p} .

In this situation $m(t) = \pi(g(t))$, where $g(t)$ is given by (10). Then, $\frac{dg}{dt} = g(t)U(t) = g(t)(\exp(tP_{\mathfrak{k}})P_{\mathfrak{p}}\exp(-tP_{\mathfrak{k}}))$ and $\|U(t)\| = \|P_{\mathfrak{p}}\| = 1$. Hence $m(t)$ is parametrized by arc length. It follows that

$$\frac{D}{dt} \left(\frac{dm}{dt} \right) = \pi_* \left(g(t) \frac{dU}{dt} \right) = \pi_* \left(g(t) \exp(tP_{\mathfrak{k}})[P_{\mathfrak{p}}, P_{\mathfrak{k}}] \exp(-tP_{\mathfrak{k}}) \right).$$

But then,

$$\left\| \frac{D}{dt} \left(\frac{dm}{dt} \right) \right\| = \left\| \exp(tP_{\mathfrak{k}})[P_{\mathfrak{p}}, P_{\mathfrak{k}}] \exp(-tP_{\mathfrak{k}}) \right\| = \left\| [P_{\mathfrak{p}}, P_{\mathfrak{k}}] \right\|.$$

■

3. Homogeneous metrics on general Stiefel manifolds $\text{St}_k^n(V)$

3.1. Stiefel manifolds

A Stiefel manifold $\text{St}_k^n(V)$ consists of ordered k orthonormal vectors v_1, \dots, v_k in an n -dimensional Euclidean vector space V . We will focus on the cases when the vector space V is an Euclidean space $V = \mathbb{R}^n$, an Hermitian complex vector space $V = \mathbb{C}^n$, or a quaternionic space $V = \mathbb{H}^n$ (with the right multiplication by scalars) equipped with its inner product $(u, v) = \sum_{l=1}^n \bar{u}_l v_l$. We will identify points $m = (v_1, \dots, v_k) \in \text{St}_k^n(V)$ with matrices M_{nk} whose columns consist of the coordinate vectors v_1, \dots, v_k with respect to a chosen orthonormal basis for V . Each such matrix M_{nk} satisfies $M_{nk}^* M_{nk} = I_k$, where $*$ stands for the transpose, the complex conjugate, or the quaternion conjugate, depending on the case.

The quaternionic Stiefel manifold requires some additional explanations. Let $1, \mathbf{i}, \mathbf{j}, \mathbf{k}$ denote the standard basis in the quaternion algebra \mathbb{H} . Then, every quaternion is a linear combination $q = q_0 + q_1 \mathbf{i} + q_2 \mathbf{j} + q_3 \mathbf{k}$ with real

coefficients. The conjugate \bar{q} of the quaternion q is given by $\bar{q} = q_0 - q_1\mathbf{i} - q_2\mathbf{j} - q_3\mathbf{k}$. The product of two quaternions is defined by using the coordinate representation and the law $\mathbf{i}^2 = \mathbf{j}^2 = \mathbf{k}^2 = \mathbf{ijk} = -1$. It follows that $\overline{qq'} = \bar{q}'\bar{q}$ and $\bar{q}q = q_0^2 + q_1^2 + q_2^2 + q_3^2 = |q|^2$. The notion of conjugacy extends to \mathbb{H}^n and yields the inner product $(v, w) = \sum_{l=1}^n \bar{v}_l w_l$. It readily follows that

$$(v\alpha, w) = \bar{\alpha}(v, w), \quad (v, w\alpha) = (v, w)\alpha, \quad (v, w) = \overline{(w, v)}, \quad (11)$$

for any $\alpha \in \mathbb{H}$. Then $\text{Sp}(n)$ is the group of matrices that leave invariant the quaternionic Hermitian product (11). It follows that $\text{Sp}(n)$ consists of $n \times n$ matrices Θ with quaternionic entries that satisfy $\Theta^*\Theta = \Theta\Theta^* = I_n$. Reminiscent of the unitary group, one can show that $\text{Sp}(n)$ is isomorphic to $\text{Sp}(2n, \mathbb{C}) \cap \text{U}(2n)$ [9, page 445].

3.2. Stiefel manifolds as homogeneous spaces

All Stiefel manifolds are homogeneous manifolds, and can be realized as the quotients of Lie groups through several group actions. Below we will describe two such actions. To avoid unnecessary repetitions, we will use G_n to denote $\text{SO}(n)$ in the real case, $\text{SU}(n)$ in the complex case, and $\text{Sp}(n)$ in the quaternionic case.

Both G_n and G_k act on elements of $\text{St}_k^n(V)$, represented as $n \times k$ matrices. The first group acts by the matrix multiplication on the left, and the second group by the matrix multiplication on the right.

Let us first consider the full group action.

• $\text{St}_k^n(V)$ as a homogeneous manifold $G_n \times G_k / G_k \times G_{n-k}$. Here the full group $G = G_n \times G_k$ acts on $\text{St}_k^n(V)$ by

$$\phi((r, s), m) = rms^{-1}, \quad m \in \text{St}_k^n(V), \quad r \in G_n, \quad s \in G_k.$$

The action is transitive, and the orbit through $m = I_{nk} = \begin{pmatrix} I_k \\ 0 \end{pmatrix}$ consists of the matrices $M_{nk} = rI_{nk}s^{-1}$, $r \in G_n$, $s \in G_k$. The isotropy group K , that leaves I_{nk} fixed, consists of matrices $r \in G_n$ and $s \in G_k$ such that

$$rI_{nk} = I_{nk}s. \quad (12)$$

The matrix $r \in G_n$ can be written in block form as $r = \begin{pmatrix} R_1 & R_2 \\ R_3 & R_4 \end{pmatrix}$, with R_1 an $k \times k$ -matrix, and the remaining matrices of the corresponding sizes. Then (12) holds for r only when $R_1 = s$ and $R_2 = R_3 = 0$. Therefore,

$r = \begin{pmatrix} R_1 & 0 \\ 0 & R_4 \end{pmatrix}$. This shows that the isotropy group K is

$$K = \left\{ \left(\begin{pmatrix} S & 0 \\ 0 & T \end{pmatrix}, S \right) : S \in G_k, T \in G_{n-k} \right\} \cong G_k \times G_{n-k}. \quad (13)$$

• $\text{St}_k^n(V)$ as a homogeneous manifold G_n/G_{n-k} . The reduced group action is given by

$$\phi(g, m) = gm, \quad m \in \text{St}_k^n(V), \quad g \in G_n.$$

The orbit of the reduced action through the matrix $m = I_{nk}$ consists of matrices $M_{nk} = gI_{nk} \in \text{St}_k^n(V)$, $g \in G_n$, i.e., the first k columns of g . The isotropy group at the point I_{nk} is equal to

$$K = \left\{ \begin{pmatrix} I_k & 0 \\ 0 & H \end{pmatrix} : H \in G_{n-k} \right\} \cong G_{n-k}. \quad (14)$$

The above shows that each Stiefel manifold $\text{St}_k^n(V)$ can be represented in two ways: as the full quotient $(G_n \times G_k)/G_k \times G_{n-k}$, as well as the reduced quotient G_n/G_{n-k} .

In what follows we will write $\text{St}_k^n(V) = G/K$ with the understanding that G and K are either $G = G_n$ and $K = G_{n-k}$, or $G = G_n \times G_k$ and $K = G_k \times G_{n-k}$, depending on the context. Then, \mathfrak{g} and \mathfrak{k} will denote the Lie algebras of G and K . Similarly \mathfrak{g}_k , \mathfrak{g}_{n-k} will denote the Lie algebras of G_k and G_{n-k} .

3.2.1. Homogeneous metrics

Each Lie algebra \mathfrak{g} is endowed with a positive definite bilinear form $\langle \cdot, \cdot \rangle$. If \mathfrak{g}_n is either $\mathfrak{so}(n)$ or $\mathfrak{u}(n)$, then the form is given by

$$\langle A, B \rangle = -\frac{1}{2} \text{Tr}(AB) \quad \text{for } A, B \in \mathfrak{g}_n, \quad (15)$$

with $\text{Tr}(A)$ the trace of the matrix A . On $\mathfrak{g}_n = \mathfrak{sp}(n)$ the form is written as

$$\langle A, B \rangle = -\frac{1}{4} \text{Tr}(AB + (AB)^*). \quad (16)$$

We will often refer to the above forms on \mathfrak{g} as *the trace form*, and to the induced metric as the *trace metric*. The trace form extends to the product $\mathfrak{g}_k \times \mathfrak{g}_{n-k}$ with $\langle \cdot, \cdot \rangle = \langle \cdot, \cdot \rangle_1 + \langle \cdot, \cdot \rangle_2$, where $\langle \cdot, \cdot \rangle_1$ is the trace form in \mathfrak{g}_k and $\langle \cdot, \cdot \rangle_2$ is the trace form in \mathfrak{g}_{n-k} . The trace form on \mathfrak{g} is Ad_G invariant and satisfies (3). Thus the corresponding left-invariant Riemannian metric $\langle \cdot, \cdot \rangle_G$ is also bi-invariant.

We will now introduce three decompositions $\mathfrak{g} = \mathfrak{p} \oplus \mathfrak{k}$ that conform to (2) and induce the left-invariant horizontal distributions $\mathcal{H}_{\mathfrak{p}}$ that are relevant for the applications.

The reduced horizontal distribution is the horizontal distribution associated with the representation $\text{St}_k^n(V) = G_n/G_{n-k}$. It is induced by the orthogonal complement

$$\mathfrak{p} = \left\{ \begin{pmatrix} A & B \\ -B^* & 0 \end{pmatrix} : A \in \mathfrak{g}_k \right\} \quad (17)$$

with respect to the trace metric to the isotropy algebra

$$\mathfrak{k} = \left\{ \begin{pmatrix} 0 & 0 \\ 0 & D \end{pmatrix} : D \in \mathfrak{g}_{n-k} \right\} \quad (18)$$

of the isotropy group K in (14).

The other two horizontal distributions are associated with the representation $\text{St}_k^n(V) = (G_n \times G_{n-k})/(G_k \times G_{n-k})$.

First, is *the orthogonal horizontal distribution on $G = G_n \times G_{n-k}$* . It is induced by the orthogonal complement

$$\mathfrak{p} = \mathfrak{k}^\perp = \left\{ \left(\begin{pmatrix} A & B \\ -B^* & 0 \end{pmatrix}, -A \right) : A \in \mathfrak{g}_k \right\} \quad (19)$$

with respect to the trace metric to the isotropy algebra

$$\mathfrak{k} = \left\{ \left(\begin{pmatrix} C & 0 \\ 0 & D \end{pmatrix}, C \right) : C \in \mathfrak{g}_k, D \in \mathfrak{g}_{n-k} \right\} \quad (20)$$

of the isotropy group K in (13).

Second, *the quasi-geodesic horizontal distribution on $G = G_n \times G_{n-k}$* . This horizontal distribution is induced by the vector space

$$\mathfrak{p} = \left\{ \left(\begin{pmatrix} 0 & B \\ -B^* & 0 \end{pmatrix}, A \right) : A \in \mathfrak{g}_k \right\} \quad (21)$$

that is not orthogonal to \mathfrak{k} in (20).

Evidently \mathfrak{p} is transversal to \mathfrak{k} , and satisfies $\mathfrak{g} = \mathfrak{p} \oplus \mathfrak{k}$, because for any $\begin{pmatrix} A_1 & A_2 \\ -A_2^* & A_3 \end{pmatrix} \in \mathfrak{g}_n$ and $B \in \mathfrak{g}_k$ we have

$$\left(\begin{pmatrix} A_1 & A_2 \\ -A_2^* & A_3 \end{pmatrix}, B \right) = \left(\begin{pmatrix} 0 & A_2 \\ -A_2^* & 0 \end{pmatrix}, B - A_1 \right) + \left(\begin{pmatrix} A_1 & 0 \\ 0 & A_3 \end{pmatrix}, A_1 \right).$$

The distribution defined by the left translations of \mathfrak{p} in (21) is called *quasi-geodesic distribution* to emphasize the connection with curves called *quasi-geodesic*, whose significance was demonstrated in [12] in interpolation problems arising from applications.

Lemma 1. *In all three cases above, the reduced, orthogonal, and quasi-geodesic case, the decomposition $\mathfrak{g} = \mathfrak{p} \oplus \mathfrak{k}$ satisfies (2).*

Proof: In the reduced and the orthogonal cases $\langle [\mathfrak{p}, \mathfrak{k}], \mathfrak{k} \rangle = \langle \mathfrak{p}, [\mathfrak{k}, \mathfrak{k}] \rangle = 0$, because \mathfrak{k} is a Lie algebra, and it is orthogonal to \mathfrak{p} . Therefore, $[\mathfrak{p}, \mathfrak{k}] \subset \mathfrak{p}$. It is also true for the quasi-geodesic case by a direct calculation. So in all cases $[\mathfrak{p}, \mathfrak{k}] \subseteq \mathfrak{p}$.

We will now show that $\mathfrak{k} \subseteq [\mathfrak{p}, \mathfrak{p}]$. In the calculations below, $E_{i,j}$ denotes the $n \times n$ matrix with entry (i, j) equal to 1 and all other entries equal to 0, and $A_{i,j} = E_{i,j} - E_{j,i}$. Matrices $A_{i,j}$ satisfy the following commutator properties:

$$[A_{i,j}, A_{f,l}] = -\delta_{il}A_{j,f} - \delta_{jf}A_{i,l} + \delta_{if}A_{j,l} + \delta_{jl}A_{i,f},$$

where δ_{ij} denotes the Kronecker delta function.

Taking into consideration the structure of the matrices in \mathfrak{k} and \mathfrak{p} in the reduced case, given respectively by (18) and (17), it is clear that

$$\{A_{k+i,k+j}, 1 \leq i < j \leq n - k\}$$

is a basis for \mathfrak{k} , while

$$\{A_{i,j}, 1 \leq i < j \leq k\} \cup \{A_{i,k+j}, 1 \leq i \leq k, 1 \leq j \leq n - k\}$$

is a basis for \mathfrak{p} . Since for any $A_{k+i,k+j}$ in \mathfrak{k} , there exists $A_{l,k+i}$ and $A_{l,k+j}$ in \mathfrak{p} such that

$$A_{k+i,k+j} = [A_{l,k+i}, A_{l,k+j}], \quad (22)$$

we have proved that in the reduced case $\mathfrak{k} \subseteq [\mathfrak{p}, \mathfrak{p}]$. To show that this inclusion is also true for the other two distributions, it is enough to take into account the structure of the matrices that define the subspaces \mathfrak{k} (given by (20)) and \mathfrak{p} , given either by (19) or (21), and use (22) together with the following extra identity:

$$A_{i,j} = [A_{i,k+l}, A_{j,k+l}],$$

which is valid for all $1 \leq i < j \leq k$ and $l \in \{1, \dots, n - k\}$. ■

Thus, we obtain the following corollary.

Corollary 3. *All three horizontal distributions \mathcal{H}_p are two-step bracket generating distributions, $\mathcal{H}_p(g) + [\mathcal{H}_p, \mathcal{H}_p](g) = T_g G$ for all $g \in G$.*

Theorem 2. *The homogeneous metrics on the Stiefel manifold $\text{St}_k^n(V)$ induced by the sub-Riemannian metrics relative to the reduced horizontal distribution on G_n , and the sub-Riemannian metric on $G_n \times G_k$ relative to the quasi-geodesic distribution are equal, and they are different from the homogeneous metric induced by the sub-Riemannian metric relative to the orthogonal horizontal distribution on $G_n \times G_k$.*

Proof: We start with the proof of the first statement. Let $\dot{m}(t)$ denote the tangent vector of a curve $m(t)$ in $\text{St}_k^n(V)$. Then, $m(t) = g(t)I_{nk}$ and $m(t) = r(t)I_{nk}s^*(t)$ for some horizontal curves $g(t) \in G_n$ and $(r(t), s(t))$ in $G = G_n \times G_k$. Then

$$\dot{m}(t) = g(t)W(t)I_{nk} = r(t)(U_1(t)I_{nk} - I_{nk}U_2(t))s^*(t), \quad (23)$$

where

$$\frac{dg}{dt} = g(t)W(t), \quad \frac{dr}{dt} = r(t)U_1(t), \quad \frac{ds}{dt} = s(t)U_2(t).$$

It will be convenient to embed G_k into G_n by identifying $s \in G_k$ with $\begin{pmatrix} S & 0 \\ 0 & I_{n-k} \end{pmatrix}$, and identify I_{nk} with $\begin{pmatrix} I_k & 0 \\ 0 & 0 \end{pmatrix}$, so that all the matrices above can be written as $n \times n$ matrices in block form:

$$W = \begin{pmatrix} A & B \\ -B^* & 0 \end{pmatrix}, \quad U_1 = \begin{pmatrix} 0 & C \\ -C^* & 0 \end{pmatrix}, \quad U_2 \sim \begin{pmatrix} D & 0 \\ 0 & 0 \end{pmatrix}, \quad A, D \in \mathfrak{g}_k.$$

Let $\|\dot{m}(t)\|_1$ and $\|\dot{m}(t)\|_2$ denote the lengths of $\dot{m}(t)$ relative to the homogeneous metrics induced by the reduced and the quasi-geodesic distributions. Since the sub-Riemannian metrics on G are left-invariant, we need only to compare the norms of the horizontal vectors on the corresponding Lie algebras. Thus

$$\|W(t)\|_p^2 = \|A(t)\|^2 + \text{Tr}(B(t)B^*(t)),$$

is equal to $\|\dot{m}(t)\|_1^2$, and

$$\begin{aligned} \|(U_1(t), U_2(t))\|_p^2 &= \|U_1(t)\|^2 + \|U_2(t)\|^2 \\ &= \|D(t)\|^2 + \text{Tr}(C(t)C^*(t)) \end{aligned}$$

is equal to $\|\dot{m}(t)\|_2^2$. We need to show that $\|\dot{m}(t)\|_1 = \|\dot{m}(t)\|_2$.

Since I_{nk} commutes with s^* ,

$$m(t) = g(t) I_{nk} = r(t) I_{nk} s^*(t) = r(t) s^*(t) I_{nk},$$

and therefore, $r(t) s^*(t) h = g(t)$ for some constant $h \in K$, where K is given by (14). Now equation (23) implies that $r s^* h W h^* I_{nk} = r(U_1 - U_2) I_{nk} s^*$, or $(U_1 - U_2) I_{nk} = s^* h W h^* s I_{nk}$, because I_{nk} commutes with U_2 , and $I_{nk} h = h I_{nk} = I_{nk}$. This equality implies that

$$U_1 - U_2 = \begin{pmatrix} -D(t) & C(t) \\ -C^*(t) & 0 \end{pmatrix} = \tilde{W},$$

because both of these matrices are of the form $\begin{pmatrix} X & Y \\ -X^* & 0 \end{pmatrix}$. Here we denote $\tilde{W} = s^* h W h^* s$. Then,

$$\begin{aligned} \|\dot{m}(t)\|_1^2 &= \|W(t)\|_{\mathfrak{p}}^2 = \|\tilde{W}(t)\|_{\mathfrak{p}}^2 = -\frac{1}{2} \text{Tr}(\tilde{W}^2(t)) \\ &= -\frac{1}{2} \text{Tr} \begin{pmatrix} -D(t) & C(t) \\ -C^*(t) & 0 \end{pmatrix}^2 = \|D(t)\|^2 + \text{Tr}(C(t)C^*(t)) \\ &= \|(U_1(t), U_2(t))\|_{\mathfrak{p}}^2 = \|\dot{m}(t)\|_2^2. \end{aligned}$$

Now we prove the second statement of the proposition. In this case

$$\frac{dg}{dt} = g(t) \tilde{U}_1(t), \quad \frac{ds}{dt} = s(t) \tilde{U}_2(t)$$

with $\tilde{U}_1(t) = \begin{pmatrix} A(t) & B(t) \\ -B^*(t) & 0 \end{pmatrix}$, $\tilde{U}_2(t) = \begin{pmatrix} -A(t) & 0 \\ 0 & 0 \end{pmatrix}$. The norm of $(\tilde{U}_1, \tilde{U}_2)$ with respect to orthogonal horizontal distribution is given by

$$\|(\tilde{U}_1(t), \tilde{U}_2(t))\|_{\mathfrak{p}}^2 = \|\tilde{U}_1(t)\|^2 + \|\tilde{U}_2(t)\|^2 = 2\|A(t)\|^2 + \text{Tr}(B(t)B^*(t)).$$

A calculation similar to the one above shows that

$$\tilde{W} = s^* h W h^* s = \begin{pmatrix} 2A(t) & B(t) \\ -B^*(t) & 0 \end{pmatrix} = \tilde{U}_1 - \tilde{U}_2.$$

Therefore,

$$\begin{aligned} \|W(t)\|_{\mathfrak{p}}^2 &= \|\tilde{W}(t)\|_{\mathfrak{p}}^2 = 4\|A(t)\|^2 + \text{Tr}(B(t)B^*(t)) \\ &\neq \|(\tilde{U}_1(t), \tilde{U}_2(t))\|_{\mathfrak{p}}^2. \end{aligned}$$

■

3.3. Sub-Riemannian geodesics

3.3.1. Geodesics on $\text{St}_k^n(V)$, induced by the reduced horizontal distribution

In this case, $\mathfrak{p} = \mathfrak{k}^\perp$ where \mathfrak{k} and \mathfrak{p} are given by (18) and (17) respectively, and hence by (10), the sub-Riemannian geodesics on G_n are of the form

$$g(t) = g_0 \exp \left(t \begin{pmatrix} A & B \\ -B^* & D \end{pmatrix} \right) \begin{pmatrix} I_k & 0 \\ 0 & e^{-tD} \end{pmatrix},$$

Their projections on $\text{St}_k^n(V)$ are given by

$$m(t) = \pi(g(t)) = g_0 \exp \left(t \begin{pmatrix} A & B \\ -B^* & D \end{pmatrix} \right) I_{nk},$$

and the Riemannian geodesics are of the form

$$m(t) = g_0 e^{t\Omega} I_{nk}, \quad \Omega = \begin{pmatrix} A & B \\ -B^* & 0 \end{pmatrix}. \quad (24)$$

These geodesics are called canonical in [7] and normal in [8].

3.3.2. Geodesics on $\text{St}_k^n(V)$ induced by the orthogonal horizontal distribution

In this case \mathfrak{p} and \mathfrak{k} are given by (19) and (20), respectively. Then $P_{\mathfrak{p}} = \left(\begin{pmatrix} A & B \\ -B^* & 0 \end{pmatrix}, -A \right)$, and $P_{\mathfrak{k}} = \left(\begin{pmatrix} C & 0 \\ 0 & D \end{pmatrix}, C \right)$, which leads to

$$\begin{aligned} g(t) &= g_0 \exp(t(P_{\mathfrak{p}} + P_{\mathfrak{k}})) \exp(-tP_{\mathfrak{k}}) \\ &= g_0 \left(\exp \left(t \begin{pmatrix} A+C & B \\ -B^* & D \end{pmatrix} \right), e^{t(-A+C)} \right) \left(\begin{pmatrix} e^{-tC} & 0 \\ 0 & e^{-tD} \end{pmatrix}, e^{-tC} \right). \end{aligned}$$

The projection is given by

$$\begin{aligned} \pi(g(t)) &= g_0 \left(\exp \left(t \begin{pmatrix} A+C & B \\ -B^* & D \end{pmatrix} \right) \right) \left(\begin{pmatrix} e^{-tC} & 0 \\ 0 & e^{-tD} \end{pmatrix} I_{nk} e^{tC} e^{t(A-C)} \right) \\ &= g_0 \left(\exp \left(t \begin{pmatrix} A+C & B \\ -B^* & D \end{pmatrix} \right) I_{nk} e^{t(A-C)} \right). \end{aligned}$$

If $g_0 = (r, s) \in G_n \times G_k$, then the geodesics through the point $g_0 I_{nk} = r I_{nk} s^*$, corresponding to $P_{\mathfrak{k}} = 0$, i.e., $C = D = 0$, have the form

$$m(t) = g_0 \exp \left(t \begin{pmatrix} A & B \\ -B^* & 0 \end{pmatrix} \right) I_{nk} e^{tA} = r e^{t\Omega} I_{nk} e^{tA} s^* \quad (25)$$

3.3.3. Geodesics on $\text{St}_k^n(V)$ induced by quasi-geodesic horizontal distribution

The quasi-geodesic distribution is generated by (20) and (21). An easy calculation shows that

$$\mathfrak{p}^\perp = \left\{ \left(\begin{pmatrix} E & 0 \\ 0 & F \end{pmatrix}, 0 \right) : E \in \mathfrak{g}_k, F \in \mathfrak{g}_{n-k} \right\}.$$

Evidently, \mathfrak{p}^\perp is a Lie subalgebra of \mathfrak{g} , hence its sub-Riemannian geodesics are given by (7). Let

$$P_{\mathfrak{p}} = \left(\begin{pmatrix} 0 & B \\ -B^* & 0 \end{pmatrix}, A \right) \in \mathfrak{p}, \quad P_{\mathfrak{p}^\perp} = \left(\begin{pmatrix} E & 0 \\ 0 & F \end{pmatrix}, 0 \right) \in \mathfrak{p}^\perp. \quad (26)$$

Then

$$\begin{aligned} g(t) &= g_0 \exp(t(P_{\mathfrak{p}} + P_{\mathfrak{p}^\perp})) \exp(-tP_{\mathfrak{p}^\perp}) \\ &= g_0 \left(\exp \left(t \begin{pmatrix} E & B \\ -B^* & F \end{pmatrix} \right) \begin{pmatrix} e^{-tE} & 0 \\ 0 & e^{-tF} \end{pmatrix}, e^{tA} \right). \end{aligned} \quad (27)$$

If $g_0 = (r, s) \in G_n \times G_k$, then

$$\begin{aligned} \pi(g(t)) &= r \left(\exp \left(t \begin{pmatrix} E & B \\ -B^* & F \end{pmatrix} \right) \begin{pmatrix} e^{-tE} & 0 \\ 0 & e^{-tF} \end{pmatrix} I_{nk} e^{-tA} s^* \right) \\ &= r \left(\exp \left(t \begin{pmatrix} E & B \\ -B^* & F \end{pmatrix} \right) \begin{pmatrix} e^{-tE} e^{-tA} & 0 \\ 0 & e^{-tF} \end{pmatrix} I_{nk} s^* \right). \end{aligned} \quad (28)$$

According to Theorem 1, the geodesics on $\text{St}_n^k(V)$ are the projections of the above curves for which $P_{\mathfrak{p}} + P_{\mathfrak{p}^\perp}$ is orthogonal to \mathfrak{k} . The orthogonal complement \mathfrak{k}^\perp consists of matrices of the form $\left(\begin{pmatrix} X & Y \\ -Y^* & 0 \end{pmatrix}, -X \right)$. So, if $P_{\mathfrak{p}}$ and $P_{\mathfrak{p}^\perp}$ are as in (26), then $P_{\mathfrak{p}} + P_{\mathfrak{p}^\perp} \in \mathfrak{k}^\perp$ if and only if $E = -A$ and $F = 0$. In such a case,

$$P_{\mathfrak{p}} = \left(\begin{pmatrix} 0 & B \\ -B^* & 0 \end{pmatrix}, A \right), \quad P_{\mathfrak{p}^\perp} = \left(\begin{pmatrix} -A & 0 \\ 0 & 0 \end{pmatrix}, 0 \right),$$

and

$$P_{\mathfrak{p}} + P_{\mathfrak{p}^\perp} = \left(\begin{pmatrix} -A & B \\ -B^* & 0 \end{pmatrix}, A \right).$$

It then follows from (28) that the geodesics at $g_0 = (r, s) \in G_n \times G_k$ are of the form

$$m(t) = r e^{t\tilde{\Omega}} I_{nk} s^*, \quad \tilde{\Omega} = \begin{pmatrix} -A & B \\ -B^* & 0 \end{pmatrix}. \quad (29)$$

At first glance, formulas (29) and (24) give different curves. To show that it is not the case, let $g = r s^*$, where, whenever convenient, we identify $s \in G_k$ with $s = \begin{pmatrix} S & 0 \\ 0 & I_{n-k} \end{pmatrix} \in G_n$. Then,

$$m(t) = r e^{t\tilde{\Omega}} I_{nk} s^* = r e^{t\tilde{\Omega}} s^* I_{nk} = r s^* s e^{t\tilde{\Omega}} s^* I_{nk} = g e^{t\Omega} I_{nk},$$

where $\Omega = s \tilde{\Omega} s^* = \begin{pmatrix} -S A S^* & S B \\ -B^* S^* & 0 \end{pmatrix}$.

We are now almost ready to relate the above formalism to the quasi-geodesic curves. The following lemma will lead the way.

Lemma 2. *Curves*

$$m(t) = r \left(\exp t \begin{pmatrix} E & B \\ -B^* & F \end{pmatrix} \begin{pmatrix} e^{-tE} e^{-tA} & 0 \\ 0 & e^{-tF} \end{pmatrix} I_{nk} s^* \right)$$

in $\text{St}_k^n(V)$ that are the projections of sub-Riemannian geodesic on $G_n \times G_k$ relative to the quasi-geodesic distribution are also the projections of horizontal curves $g(t)$ in G_n that are solutions of

$$\frac{dg}{dt} = g(t) \begin{pmatrix} -\tilde{A} & e^{t\tilde{A}} e^{t\tilde{E}} \tilde{B} e^{-tF} \\ -e^{tF} \tilde{B}^* e^{-t\tilde{E}} e^{-t\tilde{A}} & 0 \end{pmatrix} \quad (30)$$

with $\tilde{A} = S A S^*$, $\tilde{E} = S E S^*$, and $\tilde{B} = S B$.

Proof: The sub-Riemannian geodesic relative to the quasi-geodesic distribution is given by (27) and its projection on the Stiefel manifold is given by (28). The latter can be written as

$$m(t) = g_0 e^{t\Phi} \Delta(t) I_{nk}, \quad g_0 = r s^*, \quad (31)$$

with

$$e^{t\Phi} = s \exp \left(t \begin{pmatrix} E & B \\ -B^* & F \end{pmatrix} \right) s^* = \exp \left(t \begin{pmatrix} \tilde{E} & \tilde{B} \\ -\tilde{B}^* & F \end{pmatrix} \right) \quad \text{and}$$

$$\Delta = s \begin{pmatrix} e^{-tE} e^{-tA} & 0 \\ 0 & e^{-tF} \end{pmatrix} s^* = \begin{pmatrix} e^{-t\tilde{E}} e^{-t\tilde{A}} & 0 \\ 0 & e^{-tF} \end{pmatrix}.$$

The curve $g(t) = g_0 e^{t\Phi} \Delta(t)$ is the curve on G_n having the derivative

$$\dot{g}(t) = g_0 e^{t\Phi} \Delta \left(\Delta^{-1} \Phi \Delta + \Delta^{-1} \dot{\Delta} \right) = g(t) \left(\Delta^{-1} \Phi \Delta + \Delta^{-1} \dot{\Delta} \right).$$

A straightforward calculation shows that

$$\Delta^{-1} \Phi \Delta + \Delta^{-1} \dot{\Delta} = \begin{pmatrix} -\tilde{A} & e^{t\tilde{A}} e^{t\tilde{E}} \tilde{B} e^{-tF} \\ e^{-tF} \tilde{B}^* e^{-t\tilde{E}} e^{-t\tilde{A}} & 0 \end{pmatrix}.$$

Therefore, $g(t)$ is a horizontal curve in G_n satisfying the conditions of the lemma. \blacksquare

We finally come to the quasi-geodesic curves.

Definition 3. *Quasi-geodesic curves through a point $m = r I_{nk} s^*$ in $\text{St}_k^n(V)$ are curves $\gamma(t)$ having the form*

$$\gamma(t) = r \exp(t\Psi) I_{nk} e^{-tA} s^*,$$

for some matrices $\Psi = \begin{pmatrix} 0 & B \\ -B^* & 0 \end{pmatrix}$, with $B \in \mathcal{M}_{k(n-k)}(V)$ and $A \in \mathfrak{g}_k$.

Alternatively, quasi-geodesic curves can be defined as curves

$$\gamma(t) = \exp(tX) m \exp(tY),$$

where $X = r\Psi r^*$, and $Y = -sAs^* \in \mathfrak{g}_k$. Indeed,

$$\begin{aligned} \gamma(t) &= r \exp(t\Psi) I_{nk} \exp(-tA) s^* = r \exp(t\Psi) r^* r I_{nk} s^* s \exp(-tA) s^* \\ &= \exp(t(r\Psi r^*)) r I_{nk} s^* \exp(-tsAs^*) = \exp(tX) m \exp(tY). \end{aligned}$$

Proposition 6. *Quasi-geodesic curves coincide with the projections of sub-Riemannian geodesics in (27) with $P_{\mathfrak{p}^\perp} = 0$. They are curves of constant geodesic curvature. A quasi-geodesic is a Riemannian geodesic on $\text{St}_k^n(V)$ if either $P_{\mathfrak{p}} = (0, A)$, or $P_{\mathfrak{p}} = \left(\begin{pmatrix} 0 & B \\ -B^* & 0 \end{pmatrix}, 0 \right)$.*

Proof: The first statement is a consequence of formula (28).

To show the second statement we will use Lemma 2. When $P_{\mathfrak{p}^\perp} = 0$, $E = F = 0$, and $m(t) = g(t) I_{nk}$, where $g(t) = g_0 \exp\left(t \begin{pmatrix} 0 & \tilde{B} \\ -\tilde{B}^* & 0 \end{pmatrix}\right) \begin{pmatrix} e^{-t\tilde{A}} & 0 \\ 0 & I \end{pmatrix}$.

Then equation (30) reduces to

$$\frac{dg}{dt} = g(t) \begin{pmatrix} -\tilde{A} & e^{t\tilde{A}}\tilde{B} \\ -\tilde{B}^*e^{-t\tilde{A}} & 0 \end{pmatrix} = g(t) \begin{pmatrix} e^{t\tilde{A}} & 0 \\ 0 & I \end{pmatrix} \begin{pmatrix} -\tilde{A} & \tilde{B} \\ -\tilde{B}^* & 0 \end{pmatrix} \begin{pmatrix} e^{-t\tilde{A}} & 0 \\ 0 & I \end{pmatrix}.$$

Since $m(t)$ is the projection of a sub-Riemannian geodesic, it is parametrized by the arc length. That implies that $U(t) = \begin{pmatrix} e^{t\tilde{A}} & 0 \\ 0 & I \end{pmatrix} \begin{pmatrix} -\tilde{A} & \tilde{B} \\ -\tilde{B}^* & 0 \end{pmatrix} \begin{pmatrix} e^{-t\tilde{A}} & 0 \\ 0 & I \end{pmatrix}$ is of unit length.

The geodesic curvature of $m(t)$ is given by $\|\frac{D}{dt}(\frac{dm}{dt})\|$ relative to the homogeneous metric, where $\frac{D}{dt}$ denotes the covariant derivative. An argument completely analogous to that in Proposition 5 shows that

$$\begin{aligned} \left\| \frac{D}{dt} \left(\frac{dm}{dt} \right) \right\| &= \left\| \frac{dU}{dt} \right\|_{\mathfrak{p}} = \left\| \left[\begin{pmatrix} A & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} -A & B \\ -B^* & 0 \end{pmatrix} \right] \right\| \\ &= \left\| \begin{pmatrix} 0 & -AB \\ -B^*A & 0 \end{pmatrix} \right\| \end{aligned}$$

Evidently, $\|\frac{D}{dt}(\frac{dm}{dt})\| = 0$ if and only if either $A = 0$, or $B = 0$. ■

The last two statements of Proposition 6 were proved earlier in [12] by direct computations without any recourse to Lie groups.

3.4. The ambient (Euclidean, Hermitian, or quaternion Hermitian) metric on the Stiefel manifolds

Each Stiefel manifold $\text{St}_k^n(V)$ is a closed subset of the vector space $\mathcal{M}_{nk}(V)$ of $n \times k$ matrices with the entries in V , $V = \mathbb{R}^n$, $V = \mathbb{C}^n$, or $V = \mathbb{H}^n$, endowed with the usual quadratic form $\langle A, B \rangle_{\mathcal{M}} = \text{Tr}(A^*B)$, $A, B \in \mathcal{M}_{nk}(V)$. We will refer to $\mathcal{M}_{nk}(V)$ together with the metric induced by $\langle \cdot, \cdot \rangle_{\mathcal{M}}$ as the *ambient manifold*.

Thus each Stiefel manifold $\text{St}_k^n(V)$ is identified with a closed submanifold defined by $\{X \in \mathcal{M}_{nk}(V) : X^*X = I_n\}$ of \mathcal{M}_{nk} . And, consequently, its tangent space $T_X \text{St}_k^n(V)$ is identified with

$$T_X \text{St}_k^n(V) = \{\dot{X} \in \mathcal{M}_{n,k} : X^* \dot{X} = -\dot{X}^* X\}, \quad X \in \text{St}_k^n(V).$$

We will now consider $\text{St}_k^n(V)$ as a Riemannian manifold with the metric given by

$$(\dot{X}_1, \dot{X}_2) = \langle \dot{X}_1, \dot{X}_2 \rangle_{\mathcal{M}} = \text{Tr}(\dot{X}_1^* \dot{X}_2)$$

This choice of a metric will be called *ambient*.

We will now show that the ambient metric can be lifted to a metric $(\cdot, \cdot)_{\mathfrak{p}}$ on the space \mathfrak{p} given by (17), which then induces yet another left-invariant sub-Riemannian structure $(\mathcal{H}_p, (\cdot, \cdot)_{\mathfrak{p}})$ on G_n . We will then extract the Riemannian geodesics relative to the ambient metric by analyzing the sub-Riemannian geodesics induced by $(\cdot, \cdot)_{\mathfrak{p}}$ by a procedure that is analogous to the one described in Section 3.3.

We have shown earlier that every tangent vector \dot{X} at a point X can be lifted to a unique horizontal vector gU above X , that is $\dot{X} = gUI_{nk}$, where $U = \begin{pmatrix} A & B \\ -B^* & 0 \end{pmatrix}$ for suitable matrices A and B . Now,

$$\begin{aligned} \text{Tr}(\dot{X}_1^* \dot{X}_2) &= \text{Tr}(I_{nk}^* U_1^* g^* g U_2 I_{nk}) = \text{Tr}(I_{nk}^* U_1^* U_2 I_{nk}) \\ &= \text{Tr}\left(\begin{pmatrix} A_1^* & -B_1 \end{pmatrix} \begin{pmatrix} A_2 \\ -B_2^* \end{pmatrix} \right) = \text{Tr}(A_1^* A_2 + B_1 B_2^*). \end{aligned}$$

Thus the ambient bilinear form lifts to

$$(U_1, U_2)_{\mathfrak{p}} = \text{Tr}(A_1^* A_2) + \text{Tr}(B_1 B_2^*) \quad (32)$$

for $U_1 = \begin{pmatrix} A_1 & B_1 \\ -B_1^* & 0 \end{pmatrix} \in \mathfrak{p}$, and $U_2 = \begin{pmatrix} A_2 & B_2 \\ -B_2^* & 0 \end{pmatrix} \in \mathfrak{p}$. We therefore have two bilinear forms on \mathfrak{p} , the trace form $\langle \cdot, \cdot \rangle_{\mathfrak{p}}$ given by (15) or (16), and the quadratic form $(\cdot, \cdot)_{\mathfrak{p}}$ given by (32). They are related by the formula

$$(U_1, U_2)_{\mathfrak{p}} = \langle U_1, DU_2 + U_2 D \rangle_{\mathfrak{p}},$$

where $D = \begin{pmatrix} I_k & 0 \\ 0 & 0 \end{pmatrix}$.

3.4.1. Sub-Riemannian problem on $(G_n, \mathcal{H}_p, (\cdot, \cdot)_{\mathfrak{p}})$

We will now obtain the sub-Riemannian geodesics associated with minimizing $\frac{1}{2} \int_0^T (g^{-1} \frac{dg}{dt}, g^{-1} \frac{dg}{dt})_{\mathfrak{p}} dt$ over the horizontal curves that satisfy $g(0) = g_1$ and $g(T) = g_2$.

The Hamiltonian equations, based on the Maximum Principle, will be obtained much in the same manner as in Section 2.4. For the moment we assume that $V = \mathbb{R}^n$ or $V = \mathbb{C}^n$. We let L denote the vector in \mathfrak{g} dual to some $l \in \mathfrak{g}^*$ with respect to the metric $\langle \cdot, \cdot \rangle$ defined in (15), and write $L = L_{\mathfrak{p}} + L_{\mathfrak{k}}$ for the decomposition relative to the factors \mathfrak{p} and \mathfrak{k} defined in (17) and (18). It follows that the regular extremals are the projections

from T^*G onto G of the integral curves $L(t)$ of the lifted Hamiltonian

$$h_U(L)(t) = -\frac{1}{2}(U(t), U(t))_{\mathfrak{p}} + \langle L_{\mathfrak{p}}, U \rangle_{\mathfrak{p}},$$

subject to the optimality condition, that the extremal control $U(t)$ and the associated dual vector $L(t)$ maximize $h_U(L(t))$ over all controls U in \mathfrak{p} . If

$$L(t) = \begin{pmatrix} A(t) & B(t) \\ -B^*(t) & C(t) \end{pmatrix}, \text{ and } U = \begin{pmatrix} u & v \\ -v^* & 0 \end{pmatrix}, \text{ then}$$

$$h_U(L(t)) = -\frac{1}{2}\text{Tr}(u^*u) - \frac{1}{2}\text{Tr}(vv^*) - \frac{1}{2}\text{Tr}(A(t)u) + \frac{1}{2}\text{Tr}(B^*(t)v + B(t)v^*).$$

It follows that $h_U(L(t))$ attains the maximum relative to the control functions precisely when $2u(t) = A(t)$, and $v(t) = B(t)$. Therefore, the extremal curves $(g(t), L(t))$ are the integral curves of the Hamiltonian system generated by

$$H = \frac{1}{2}(U, U)_{\mathfrak{p}} = \frac{1}{4}\|A\|^2 + \frac{1}{2}\text{Tr}(BB^*), \quad (33)$$

i.e., they are the solutions of the system $\frac{dg}{dt} = g(t)U(t)$, $\frac{dL}{dt} = [U(t), L(t)]$. where $dH = \begin{pmatrix} \frac{1}{2}A & B \\ -B^* & 0 \end{pmatrix} = U$. Hence,

$$\begin{pmatrix} \dot{A} & \dot{B} \\ -\dot{B}^* & \dot{C} \end{pmatrix} = \begin{pmatrix} 0 & \frac{1}{2}AB - BC \\ \frac{1}{2}B^*A - CB^* & 0 \end{pmatrix}.$$

It follows that A and C are constant and that

$$B(t) = \exp\left(\frac{t}{2}A\right)B(0)\exp(-tC).$$

This yields

$$U(t) = \begin{pmatrix} \frac{1}{2}A & e^{\frac{1}{2}tA}B(0)e^{-tC} \\ -e^{tC}B^*(0)e^{-\frac{1}{2}tA} & 0 \end{pmatrix} = e^{tQ} \begin{pmatrix} \frac{1}{2}A & B(0) \\ -B^*(0) & 0 \end{pmatrix} e^{-tQ},$$

where $Q = \begin{pmatrix} \frac{1}{2}A & 0 \\ 0 & C \end{pmatrix}$.

The extremal curve $g(t)$ in G_n is a solution of

$$\frac{dg}{dt} = g(t)(e^{tQ}Pe^{-tQ}),$$

with $P = \begin{pmatrix} \frac{1}{2}A & B(0) \\ -B^*(0) & 0 \end{pmatrix}$ and $Q = \begin{pmatrix} \frac{1}{2}A & 0 \\ 0 & C \end{pmatrix}$. It then follows that the sub-Riemannian geodesic

$$g(t) = g_0 e^{t(P+Q)} e^{-tQ}$$

projects on the Stiefel manifold as

$$X(t) = \pi(g(t)) = g_0 e^{t(P+Q)} e^{-tQ} I_{nk} = g_0 e^{t(P+Q)} I_{nk} e^{-\frac{t}{2}A}, \quad (34)$$

with $P + Q = \begin{pmatrix} A & B(0) \\ -B^*(0) & C \end{pmatrix}$, and $A \in \mathfrak{g}_k$.

For the Riemannian geodesics on the Stiefel manifold we set $C = 0$, because of the transversality conditions.

Let us now show that $X(t) = g_0 e^{t(P+Q)} e^{-tQ} I_{nk}$ with the term $P + Q = \begin{pmatrix} A & B \\ -B^* & 0 \end{pmatrix} \in \mathfrak{p}$ and $Q = \begin{pmatrix} \frac{A}{2} & 0 \\ 0 & 0 \end{pmatrix}$ satisfies the Euler-Lagrange equation

$$\ddot{X} + X\dot{X}^*\dot{X} = 0 \iff \dot{X} = Y, \quad \dot{Y} = -X(Y^*Y), \quad (35)$$

found in [7] and [8] for the case $V = \mathbb{R}^n$.

We have $X(t) = g(t)I_{nk}$, $\dot{X}(t) = Y(t) = g(t)U(t)I_{nk}$, where $g(t) = g_0 e^{t(P+Q)} e^{-tQ}$, and $U(t) = e^{tQ} \begin{pmatrix} \frac{1}{2}A & B \\ -B^* & 0 \end{pmatrix} e^{-tQ}$. Then,

$$\begin{aligned} \dot{Y} &= gU^2I_{nk} + g\dot{U}I_{nk} = g(e^{tQ}P^2e^{-tQ})I_{nk} + g(e^{tQ}[P, Q]e^{-tQ})I_{nk} \\ &= g e^{tQ}(P^2 + [P, Q])e^{-tQ}I_{nk} \\ &= g \begin{pmatrix} e^{t\frac{A}{2}}(\frac{1}{4}A^2 - BB^*)e^{-t\frac{A}{2}} & 0 \\ 0 & 0 \end{pmatrix} = g \begin{pmatrix} \frac{1}{4}A^2 - e^{t\frac{A}{2}}BB^*e^{-t\frac{A}{2}} & 0 \\ 0 & 0 \end{pmatrix}. \end{aligned}$$

On the other hand,

$$X(Y^*Y) = -gI_{nk}U^2I_{nk} = -g \begin{pmatrix} \frac{1}{4}A^2 - e^{t\frac{A}{2}}BB^*e^{-t\frac{A}{2}} & 0 \\ 0 & 0 \end{pmatrix}.$$

Therefore, $X(t)$ in (34) satisfies the Euler-Lagrange equation (35) when $C = 0$.

The calculations in the case $V = \mathbb{H}^n$ are similar. We obtain

$$\begin{aligned} h_U(L(t)) &= -\frac{1}{2}\mathrm{Tr}(u^*u) - \frac{1}{2}\mathrm{Tr}(vv^*) - \frac{1}{4}\mathrm{Tr}(Au + (Au)^*) \\ &\quad + \frac{1}{4}\mathrm{Tr}(Bv^* + (Bv^*)^* + B^*v + (B^*v)^*). \end{aligned}$$

The maximum is achieved at $2u = A$ and $v = B$ giving the Hamiltonian (33) for the corresponding metric. The rest of the calculations, identical to the ones above, show that the quaternionic geodesics are given by (34), with $C = 0$.

Observe now that all the homogeneous metrics discussed above coalesce into a single metric in the extreme cases $k = n$ and $k = 1$, and in both cases agree with the ambient metric. This is obvious in the case that $k = n$, for then $\text{St}_n^n(V)$ is equal to G_n , and the homogeneous metric is equal to the bi-invariant metric on G_n .

In the case $k = 1$, the Stiefel manifolds $\text{St}_1^n(V)$ is the unit sphere, S^{n-1} in the real case, S^{2n-1} in the complex case, and S^{4n-1} in the quaternionic case. To see that the homogeneous metric coincides with the metric inherited from the ambient space V , note that the matrix $\Omega = \begin{pmatrix} A & B \\ -B^* & 0 \end{pmatrix}$ in (24) and (25) is equal to $\begin{pmatrix} 0 & b \\ -b^* & 0 \end{pmatrix}$ where b is a row vector when $k = 1$. Hence

$$m(t) = \exp \left(t \begin{pmatrix} 0 & b \\ -b^* & 0 \end{pmatrix} \right) e_1 = (I \cos \|b\|t + \frac{1}{\|b\|} \begin{pmatrix} 0 & b \\ -b^* & 0 \end{pmatrix} \sin \|b\|t) e_1.$$

Therefore, $m(t)$ is a solution of

$$\ddot{m}(t) + \|b\|^2 m(t) = 0.$$

It may be somewhat surprising that in all other cases, $1 < k < n$, the metric on $\text{St}_k^n(V)$ inherited from the ambient space $\mathcal{M}_{nk}(V)$ is less natural than the homogeneous metric on $\text{St}_k^n(V)$ relative to the reduced action of G_n .

4. Grassmann manifolds $\text{Gr}_k^n(V)$

We will now demonstrate the relevance of the sub-Riemannian structures on Lie groups, as described in the first part of this paper, to the canonical Riemannian structure of the Grassmann manifolds $\text{Gr}_k^n(V)$. We will also make use of the fact that $\text{St}_k^n(V)$ is a principal G_k bundle over $\text{Gr}_k^n(V)$ to examine the geometric properties of the projections to $\text{Gr}_k^n(V)$ of the sub-Riemannian geodesics in $\text{St}_k^n(V)$.

Recall that $\text{Gr}_k^n(V)$ is the set of all k -dimensional vector subspaces of an n -dimensional vector space V . We will continue with our notations from above, with V one of \mathbb{R}^n , \mathbb{C}^n , or \mathbb{H}^n endowed with its usual metric, except that for the moment G_n will denote $O(n)$ in the real case, $U(n)$ in the complex case,

rather than $\mathrm{SO}(n)$ and $\mathrm{SU}(n)$ as before, while in the quaternionic case G_n will be $\mathrm{Sp}(n)$, the same as before.

Then $\mathrm{Gr}_k^n(V)$ can be embedded into G_n by identifying each vector space W in $\mathrm{Gr}_k^n(V)$ with the orthogonal reflection R_W defined by

$$R_W(x) = \begin{cases} x & \text{if } x \in W, \\ -x, & \text{if } x \in W^\perp. \end{cases}$$

Group G_n acts on Grassmann manifolds $\mathrm{Gr}_k^n(V)$ under the action

$$(\mathcal{O}, W) \rightarrow \mathcal{O}W = \{\mathcal{O}w : w \in W\}, \quad \mathcal{O} \in G_n.$$

The action of G_n on $\mathrm{Gr}_k^n(V)$ can be also expressed in terms of the reflections R_W by the following:

$$(\mathcal{O}, R_W) \rightarrow \mathcal{O}R_W\mathcal{O}^*, \quad \mathcal{O} \in G_n. \quad (36)$$

It is easy to verify that this action is transitive. Therefore, $\mathrm{Gr}_k^n(V)$ can be realized as the quotient G_n/K , where

$$K = \left\{ \begin{pmatrix} A & 0 \\ 0 & C \end{pmatrix}, A \in G_k, C \in G_{n-k} \right\} \cong G_k \times G_{n-k} \quad (37)$$

is the isotropy group of $R_{W_0} = \begin{pmatrix} I_k & 0 \\ 0 & -I_{n-k} \end{pmatrix}$ associated with the vector space W_0 spanned by the standard vectors e_1, \dots, e_k .

For our purposes it is desirable to work with connected Lie groups. So, from now on we assume that G and K are connected, that is G is equal to $\mathrm{SO}(n)$, $\mathrm{SU}(n)$ or $\mathrm{Sp}(n)$, K is modified accordingly, and the quotient G/K is the oriented Grassmannians instead.

Alternatively, the decomposition $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ could have been obtained through the involutive automorphism $\sigma(g) = DgD^{-1}$ where we denote $D = \begin{pmatrix} I_k & 0 \\ 0 & -I_{n-k} \end{pmatrix}$. Then K is equal to the subgroup of fixed points of σ : $K = \{\sigma(g) = g : g \in G\}$. Note that D can be also seen as the orthogonal reflexion R_{W_0} across W_0 the linear span of e_1, \dots, e_k .

In general, an involutive automorphism $\sigma \neq \mathrm{Id}$ on a Lie group G is an automorphism that satisfies $\sigma^2 = \mathrm{Id}$. It follows that the tangent map σ_* at the group identity is a Lie algebra automorphism that satisfies $\sigma_*^2 = \mathrm{Id}$. Hence $(\sigma_* - \mathrm{Id})(\sigma_* + \mathrm{Id}) = 0$, and therefore, $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$, where

$$\mathfrak{p} = \{A \in \mathfrak{g} : \sigma_*(A) = -A\}, \quad \text{and} \quad \mathfrak{k} = \{A \in \mathfrak{g} : \sigma_*(A) = A\}.$$

The subspaces \mathfrak{p} and \mathfrak{k} are orthogonal relative to the Killing form and satisfy Cartan relations

$$\mathfrak{g} = \mathfrak{p} \oplus \mathfrak{k}, \quad [\mathfrak{p}, \mathfrak{k}] \subseteq \mathfrak{p}, \quad [\mathfrak{p}, \mathfrak{p}] \subseteq \mathfrak{k}, \quad [\mathfrak{k}, \mathfrak{k}] \subseteq \mathfrak{k}.$$

On semisimple Lie algebras $[\mathfrak{k}, \mathfrak{p}] = \mathfrak{p}$, and on simple Lie algebras $[\mathfrak{p}, \mathfrak{p}] = \mathfrak{k}$, and therefore $\mathfrak{p} + [\mathfrak{p}, \mathfrak{p}] = \mathfrak{g}$, see [10].

Our case here is a particular case of this general situation since the trace form is a scalar multiple of the Killing form. Moreover, $[\mathfrak{p}, \mathfrak{p}] = \mathfrak{k}$, as can be easily verified. So we are in the situation where $\mathfrak{p} + [\mathfrak{p}, \mathfrak{p}] = \mathfrak{g}$.

Therefore, the left-invariant distribution $\mathcal{H}_{\mathfrak{p}}$ with values in \mathfrak{p} defines a natural sub-Riemannian problem on G_n :

Find the sub-Riemannian geodesics on G_n and identify those that project on the Riemannian geodesics in the Grassmannian $\text{Gr}_k^n(V)$.

According to Theorem 1, the sub-Riemannian geodesics are given by

$$g(t) = g_0 e^{t(P_{\mathfrak{p}}+P_{\mathfrak{k}})} e^{-tP_{\mathfrak{k}}} = g_0 \exp \left(t \begin{pmatrix} A & B \\ -B^* & C \end{pmatrix} \right) \begin{pmatrix} e^{-tA} & 0 \\ 0 & e^{-tC} \end{pmatrix},$$

and their projections on $\text{Gr}_k^n(V)$, obtained by (36), are of the form

$$R(t) = g_0 \exp \left(t \begin{pmatrix} A & B \\ -B^* & C \end{pmatrix} \right) D \exp \left(-t \begin{pmatrix} A & B \\ -B^* & C \end{pmatrix} \right) g_0^*, \quad (38)$$

Since $\mathfrak{p}^{\perp} = \mathfrak{k}$, the curves in (38) have constant geodesic curvature in $\text{Gr}_k^n(V)$ by Proposition 5. The Riemannian geodesics on $\text{Gr}_k^n(V)$ are given by Corollary 2,

$$R(t) = g_0 \exp \left(t \begin{pmatrix} 0 & B \\ -B^* & 0 \end{pmatrix} \right) D \exp \left(-t \begin{pmatrix} 0 & B \\ -B^* & 0 \end{pmatrix} \right) g_0^*. \quad (39)$$

Equation (39) can be expressed in the form

$$R(t) = \exp(tP)R_0 \exp(-tP),$$

where $P = g_0 \begin{pmatrix} 0 & B \\ -B^* & 0 \end{pmatrix} g_0^*$. It is easy to verify that $R_0P + PR_0 = 0$. The converse is also true: if $P \in \mathfrak{g}_n$ satisfies $R_0P + PR_0 = 0$, then $P = g_0 \begin{pmatrix} 0 & B \\ -B^* & 0 \end{pmatrix} g_0^*$.

Let us now note that the involutive automorphism $\sigma(g) = DgD^{-1}$ is an isometry for the above sub-Riemannian structure on G since

$$\langle \sigma_*(A), \sigma_*(B) \rangle_{\mathfrak{p}} = \langle -A, -B \rangle_{\mathfrak{p}} = \langle A, B \rangle_{\mathfrak{p}}.$$

We will presently show that this isometry accounts for the geodesic symmetry of the Riemannian Grassmannian manifolds.

To elaborate, first note that $\sigma(e^{tA}) = e^{t\sigma_*(A)}$ for any $A \in \mathfrak{g}$. Next, let $F_g: G \rightarrow G$ be the mapping defined by $F_g(h) = g\sigma(g^{-1}h)$ at each $g \in G$. It follows that F_g is an isometry for the sub-Riemannian structure and satisfies $F_g(g) = g$. If $g_0 e^{t(P_{\mathfrak{p}}+P_{\mathfrak{k}})}e^{-tP_{\mathfrak{k}}}$ is a sub-Riemannian geodesic at g_0 then

$$\begin{aligned} F_{g_0}(g(t)) &= g_0 \sigma(e^{t(P_{\mathfrak{p}}+P_{\mathfrak{k}})}e^{-tP_{\mathfrak{k}}}) = g_0 e^{t\sigma_*(P_{\mathfrak{p}}+P_{\mathfrak{k}})}e^{-t\sigma_*(P_{\mathfrak{k}})} \\ &= g_0 e^{t(-P_{\mathfrak{p}}+P_{\mathfrak{k}})}e^{-tP_{\mathfrak{k}}}. \end{aligned}$$

It follows that F_g maps the sub-Riemannian geodesics at g onto the sub-Riemannian geodesics at g . The sub-Riemannian geodesics that project onto the Riemannian geodesics are given by $P_{\mathfrak{k}} = 0$, and we have

$$F_g(g e^{tP_{\mathfrak{p}}}) = g e^{-tP_{\mathfrak{p}}}.$$

It follows that $S_{\pi(g_0)}(\pi(g)) = \pi \circ F_{g_0}(g)$ is an isometry that satisfies $S_p(\gamma(t)) = \gamma(-t)$ for any geodesic curve $\gamma(t)$ with $\gamma(0) = p$.

Any Riemannian space M in which the map $S_p: \gamma(t) \mapsto \gamma(-t)$ is an isometry for any geodesic γ is called *symmetric Riemannian space* [6, 9]. The above shows that the oriented Grassmannian manifolds $\text{Gr}_k^n(V)$ with the above homogeneous metric belongs to the class of symmetric Riemannian spaces.

4.1. Relation between Stiefel and Grassmann manifolds

Every k -dimensional subspace W of V is in one to one correspondence with the orthogonal reflection R_W and the orthogonal projection Π_W defined by

$$\Pi_W(x) = \begin{cases} x, & \text{if } x \in W \\ 0, & \text{if } x \in W^\perp. \end{cases}$$

The map

$$W \rightarrow \Pi_W \in \{A \in \mathfrak{gl}(V) : A^* = A, A^2 = A, \dim(\ker(A)) = n - k\}$$

defines a matrix representation of $\text{Gr}_k^n(V)$ in terms of the orthogonal projections. The passage from the reflections to the projections is given by a

simple formula

$$R_W = 2\Pi_W - I.$$

Therefore $R_W = gR_{W_0}g^*$ corresponds to $2\Pi_W - I = g(2\Pi_{W_0} - I)g^*$, or $\Pi_W = g\Pi_{W_0}g^*$. In particular, $\Pi_{W_0} = \begin{pmatrix} I_k & 0 \\ 0 & 0 \end{pmatrix}$ when $R_{W_0} = D$. In this representation the geodesic equations (39) become $\Pi(t) = e^{tP}\Pi_0e^{-tP}$, where $\Pi_0P + P\Pi_0 = P$.

There is a natural projection from the Stiefel manifold to the Grassmann manifold, because every point $q = [v_1, \dots, v_k]$ in $\text{St}_k^n(V)$ can be projected to the vector space W spanned by v_1, \dots, v_k . In terms of the orthogonal projections, the projection $\Pi(q) = W$ is given by $\Pi(q) = \sum_{i=1}^k v_i \otimes v_i^*$. When q is regarded as an $n \times (n - k)$ matrix with columns v_1, \dots, v_k , then $\Pi(q) = \sum_{i=1}^k v_i \otimes v_i^* = qq^*$. This identification then yields

$$\Pi(q) = qq^* \in \text{Gr}_k^n(V), \quad q \in \text{St}_k^n(V).$$

Evidently, $\Pi^{-1}(\Pi(q)) = \{qh, h \in G_k\}$. Therefore, Π is a surjection, and $\text{St}_k^n(V)$ is a principal G_k bundle over $\text{Gr}_k^n(V)$ relative to the action of $h \in G_k$ given by $\phi(h, q) = qh$, see for instance [1, 14].

Let us now go back to the curves on $\text{St}_k^n(V)$ that are the projections of various sub-Riemannian geodesics. Equations (31) capture all these curves. They are of the form

$$q(t) = g_0 e^{t\Phi} \Delta(t) I_{nk}, \quad g_0 = rs^*, \quad (40)$$

with

$$e^{t\Phi} = s \exp \left(t \begin{pmatrix} E & B \\ -B^* & F \end{pmatrix} \right) s^* = \exp \left(t \begin{pmatrix} \tilde{E} & \tilde{B} \\ -\tilde{B}^* & \tilde{F} \end{pmatrix} \right), \quad \text{and}$$

$$\Delta = s \begin{pmatrix} e^{-tE} e^{-tA} & 0 \\ 0 & e^{-tF} \end{pmatrix} s^* = \begin{pmatrix} e^{-t\tilde{E}} e^{-t\tilde{A}} & 0 \\ 0 & e^{-t\tilde{F}} \end{pmatrix},$$

where $\tilde{A} = SAS^*$, $\tilde{E} = SES^*$, $\tilde{B} = SB$, and $s = \begin{pmatrix} S & 0 \\ 0 & I_{n-k} \end{pmatrix}$.

The curves in (40) for arbitrary Φ correspond to the projection of sub-Riemannian curves relative to the quasi-geodesic distribution. The case $E = F = 0$ corresponds to the orthogonal distribution, and $A = E = F = 0$ corresponds to the projection of the sub-Riemannian geodesics relative to

the reduced orthogonal distribution. The projection of equations (40) on the Grassmannians is given by

$$\Pi(q(t)) = g_0 e^{t\tilde{\Phi}} \begin{pmatrix} I_k & 0 \\ 0 & 0 \end{pmatrix} e^{-t\tilde{\Phi}} g_0^*, \quad \tilde{\Phi} = \begin{pmatrix} \tilde{E} & \tilde{B} \\ -\tilde{B}^* & F \end{pmatrix},$$

or in terms of the orthogonal reflections, by

$$R(t) = g_0 e^{t\tilde{\Phi}} \begin{pmatrix} I_k & 0 \\ 0 & -I_{n-k} \end{pmatrix} e^{-t\tilde{\Phi}} g_0^*. \quad (41)$$

Proposition 7. *The projections of sub-Riemannian geodesics on $\text{St}_k^n(V)$ project onto curves of constant curvature in $\text{Gr}_k^n(V)$. Their curvature is zero precisely when $E = F = 0$. In particular, the quasi-geodesic curves project onto Riemannian geodesics in $\text{Gr}_k^n(V)$.*

Proof: Equations (41) are of the same form as (38), and curves in (38) have constant geodesic curvature by Proposition 5. These equations reduce to the geodesics when $E = 0$ and $F = 0$, and that case corresponds to the quasi-geodesic curves. ■

Acknowledgements: This work was partially supported by ISP project 239033/F20 of Norwegian Research Council and the work of the second author was also partially supported by joint BFS-TSF mathematics program.

References

1. C. Autenried, I. Markina, *Sub-Riemannian geometry of Stiefel manifolds*. SIAM J. Control Optim. **52** (2014), no. 2, 939-959.
2. A. Agrachev, Y. Sachkov, *Control theory from the geometric point of view*, Encyclopedia of Mathematical Sciences, **87** Springer-Verlag, New York, 2004.
3. U. Boscain, F. Rossi, *Invariant Carnot-Carathéodory metrics on S^3 , $SO(3)$, $SL(2)$, and lens spaces*. SIAM J. Control Optim. **47** (2008), no. 4, 1851-1878.
4. Z. M. Balogh, J. T. Tyson, B. Warhurst, *Sub-Riemannian vs. Euclidean dimension comparison and fractal geometry on Carnot groups*. Adv. Math. **220** (2009), no. 2, 560-619.
5. W. L. Chow, *Über Systeme von linearen partiellen Differentialgleichungen erster Ordnung*. Math. Ann. **117** (1939) 98-105.
6. P. Eberlein, *Geometry of Nonpositively Curved Manifolds Lie groups*. Chicago Lectures in Mathematics, The University of Chicago Press, Chicago, 1996.
7. A. Edelman, T. A. Arias, S. T. Smith, *The geometry of algorithms with orthogonality constraints*, SIAM J. Matrix Anal. Appl., **20**, no. 2 (1998) 303-353.
8. Y. Fedorov, B. Jovanovic, *Geodesic flows and Neumann systems on Stiefel varieties. Geometry and integrability*, Math. Z. **270**, no. 3-4, (2012) 659-698.
9. S. Helgason, *Differential geometry, Lie groups and symmetric spaces*, Academic Press, New York, 1978.

10. V. Jurdjevic, *Optimal control and geometry: integrable systems*, Cambridge University Press, Cambridge Studies in Advanced Mathematics, 2016, Cambridge, UK.
11. V. Jurdjevic, K. A. Krakowski, F. Silva Leite, *The geometry of quasi-geodesics on Stiefel manifolds*, (6 pages). To appear in Proc. International Conference on Automatic Control and Soft Computing, June 4-6, 2018, Azores - Portugal.
12. K. A. Krakowski, L. Machado, F. Silva Leite, J. Batista, *A modified Casteljau algorithm to solve interpolation problems on Stiefel manifolds*, Journal of Computational and Applied Mathematics, Volume 311, (2017) 84-99.
13. I. Kolář, P. Michor, L. Slovák, *Natural operations in differential geometry*. Springer-Verlag, Berlin, 1993. pp. 434.
14. R. Montgomery, *A tour of Subriemannian geometries, their geodesics and applications*, Amer Math. Soc., 2002, Providence, Rhode Island.
15. P. K. Rashevskij, *About connecting two points of complete nonholonomic space by admissible curve*, Uch. Zapiski Ped. Inst. K. Liebknecht **2** (1938), 83–94.
16. S. Sternberg, *Lectures on differential geometry*, Prentice- Hall, Inc, Englewood Cliffs, N.J. 1964.
17. F. Warner, *Foundations of differentiable manifolds and Lie groups*. Graduate Texts in Mathematics, 94. Springer-Verlag, New York-Berlin, 1983. 272 pp.

VELIMIR JURDJEVIC

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, TORONTO, ONTARIO M5S 3G3, CANADA

E-mail address: jurdj@math.utoronto.ca

IRINA MARKINA

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BERGEN,

P.O. BOX 7803, BERGEN N-5020, NORWAY

E-mail address: irina.markina@uib.no

FÁTIMA SILVA LEITE

INSTITUTE OF SYSTEMS AND ROBOTICS, UNIVERSITY OF COIMBRA - PÓLO II, PINHAL DE MARROCOS, 3030-290 COIMBRA, PORTUGAL, AND DEPARTMENT OF MATHEMATICS, UNIVERSITY OF COIMBRA, LARGO D. DINIS, 3001-454 COIMBRA, PORTUGAL

E-mail address: fleite@mat.uc.pt