

SYMPLECTIC KEYS AND DEMAZURE ATOMS IN TYPE C

JOÃO MIGUEL SANTOS

ABSTRACT: We compute, mimicking the Lascoux-Schützenberger type A combinatorial procedure, left and right keys for a Kashiwara-Nakashima tableau in type C . These symplectic keys have a similar role as the keys for semistandard Young tableaux. More precisely, our symplectic keys give a tableau criterion for the Bruhat order on the hyperoctahedral group and cosets, and describe Demazure atoms and characters in type C . The right and the left symplectic keys are related through the Lusztig involution. A type C Schützenberger evacuation is defined to realize that involution.

KEYWORDS: Keys, Demazure atom, Demazure character and Demazure crystal graph in type C .

MATH. SUBJECT CLASSIFICATION (2000): 05E05, 05E10, 17B37.

1. Introduction

The irreducible characters of the general linear group $GL(n)$, over \mathbb{C} , the Schur functions, are combinatorially expressed as sums on semistandard Young tableaux with entries $\leq n$ [30]. When restricting to the symplectic group $Sp(2n)$, two different types of symplectic tableaux have been proposed. King showed that the irreducible symplectic characters, the symplectic Schur polynomials, can be seen as a sum on a family of tableaux that are known as King tableaux [14], and De Concini has proposed the ones known as De Concini tableaux [6]. Kashiwara and Nakashima [12] described symplectic tableaux, which are just a variation of De Concini tableaux, with a crystal graph structure. That crystal structure allows a plactic monoid compatible with insertion and sliding algorithms, and Robinson-Schensted type correspondence, studied by Lecouvey in terms of crystal isomorphisms [18, 19]. The generalization of the notion of plactic monoid for finite Cartan types was first introduced by Littelmann using his path model [25]. Symplectic Kashiwara-Nakashima tableaux are the ones that we work with, in the corresponding ambient plactic monoid. We however note that very recently Lee has endowed King tableaux with a crystal structure [22].

Kashiwara [11] and Littelmann [24] have shown that Demazure characters [7], for any Weyl group, can be lifted to certain subsets of the crystal \mathfrak{B}^λ for a given dominant weight λ , a normal crystal with highest weight λ [5], called Demazure crystal. That is, a Demazure character (key polynomials) is the generating function of the weights over a Demazure crystals. In type C_n , we consider \mathfrak{B}^λ to be crystal of C_n -Kashiwara-Nakashima tableaux of shape λ , and Demazure characters are indexed by integer vectors in the orbit of the partition λ under the action of the Weyl group, the hyperoctahedral group B_n . They are certain non symmetric Laurent polynomials, with respect to the action of the Weyl group, which can be seen as *partial* symplectic characters, *i.e.*, sums of a certain portion of monomials in a symplectic Schur polynomial.

In type A_{n-1} , the Demazure crystals are certain subsets of the crystal \mathfrak{B}^λ , the crystal of all semistandard Young tableaux of shape λ , with entries $\leq n$. Lascoux and Schützenberger [17] identified the tableaux with nested columns as keys tableaux, and defined the right key map that sends tableaux to key tableaux. Their right key map gives a decomposition of \mathfrak{B}^λ into non intersecting subsets $\mathfrak{U}(v)$, each contain a unique key, in bijection with the vectors v in the orbit of λ , under the action of the Weyl group, \mathfrak{S}_n [17, Theorem 3.8]. They have called *standard bases* to the sum of monomial weights over $\mathfrak{U}(v)$, which, after Mason [26], are coined Demazure atoms. The decomposition describes what tableaux contribute to the Demazure crystal \mathfrak{B}_v , as a union of Demazure atoms, over an interval in the Bruhat order, on the classes modulo the stabilizer of λ . This order, induced on the orbit of λ , gives $\mathfrak{B}_v = \bigsqcup_{\lambda \leq u \leq v} \mathfrak{U}(u)$.

Our work has been motivated by the questions raised in a presentation by Azenhas [2], in *The 69th Séminaire Lotharingien de Combinatoire*. In those questions, Azenhas identified some type C_n Kashiwara-Nakashima tableaux as key tableaux, which match our identification, but it lacks a construction of the right key map, thus lacking a proof of the combinatorial description of type C Demazure characters. Note also that, during the preparation of this paper, Jaco and Lecouvey informed us about their paper [10], where they find the same key in type C , but their approach is different from ours.

Inspired by the Lascoux-Schützenberger's construction of the left and right keys of a semistandard Young tableau [17], we give a similar construction in type C_n . Our construction of the left and right keys of a Kashiwara-Nakashima tableau, in type C_n , is based on frank words in type C , that we

introduce in Section 4, and Sheats symplectic *jeu de taquin*. Our Theorem 17 is the type C analogue of [17, Theorem 3.8]. We also show, in Section 5, that both keys, left and right, are related via the Schützenberger involution in type C , or Lusztig involution, realized here in an explicit way, using symplectic insertion or sliding operations.

In [23], using the model of alcove paths, Lenart defined an initial key and a final key, for any Lie type, related via the Lusztig involution, which, in type C , have a similar behaviour to the left and right keys defined here. There is a crystal isomorphism between the alcove path model and the Kashiwara-Nakashima tableau model in types A and C [20, 21]. Since right and left keys in type C are explicitly related through the Schützenberger involution in type C , or Lusztig involution, the left and right key maps in types A and C coincide in the aforesaid approaches or models.

The paper is organized as follows. In Section 2, we discuss the Weyl group of type C , the signed permutation group B_n , the Bruhat order on B_n and on its cosets, modulo the stabilizer of λ , the Kashiwara-Nakashima tableaux and the symplectic key tableaux. Those key tableaux are used in Proposition 6 to explicitly construct the minimal length coset representatives. We recall some results from Bjorner and Brenti's book [4] and Proctor [27], that lead to a tableau criterion, in theorems 5 and 7, for the Bruhat order on B_n and on its cosets, using the symplectic key tableaux. In Section 3, we recall the Baker-Lecouvey insertion, the Sheats symplectic *jeu de taquin* and use them to discuss the plactic and coplactic monoids and the Robinson-Schensted type C correspondence. These monoids have a natural interpretation in the type C Kashiwara crystal, for a $U_q(sp_{2n})$ -module, in terms of connected components and crystal isomorphic connected components. In Section 4, we extend the concept of frank word, in type A , to type C and, with the help of symplectic *jeu de taquin*, we present, in Theorem 14, our right and left key maps. Using the right key map, we describe the tableaux that contribute to a Demazure atom and to a Demazure crystal, which is our main result, Theorem 17. In Section 5, we develop a type C evacuation within the plactic monoid, an analogue of the J -operation discussed by Schützenberger for semistandard Young tableaux in [28]. Proposition 21 shows that the evacuation of the right key of a Kashiwara-Nakashima tableau is the left key of the evacuation of the same tableau. This is an explicit realization of Lusztig involution using insertion and sliding operations in type C .

2. Weyl group of type C , Bruhat order and symplectic key tableau

Fix $n \in \mathbb{N}_{>0}$. Define the sets $[n] = \{1, \dots, n\}$ and $[\pm n] = \{1, \dots, n, \bar{n}, \dots, \bar{1}\}$ where \bar{i} is just another way of writing $-i$. In the second set we will consider the following order of its elements: $1 < \dots < n < \bar{n} < \dots < \bar{1}$ instead of the usual order.

Consider the group B_n , with generators s_i , $1 \leq i \leq n$, having the following presentation, regarding the relations among the generators,

$$B_n := \langle s_1, \dots, s_n \mid s_i^2 = 1, 1 \leq i \leq n; (s_i s_{i+1})^3 = 1, 1 \leq i \leq n-2; \\ (s_{n-1} s_n)^4 = 1; (s_i s_j)^2 = 1, 1 \leq i < j \leq n, |i-j| > 1 \rangle,$$

known as hyperoctahedral group or signed symmetric group. This group is a Coxeter group and we consider the (strong) Bruhat order on its elements [4]. The elements of B_n can be seen as odd bijective maps from $[\pm n]$ to itself, i.e., for all $\sigma \in B_n$ we have $\sigma(i) = -\sigma(-i)$, $i \in [\pm n]$. The subgroup with the generators s_1, \dots, s_{n-1} is the symmetric group \mathfrak{S}_n and its elements can be seen as bijections from $[n]$ to itself. Both groups can also be seen as groups of $n \times n$ matrices. The elements of the symmetric group can be identified with the permutation matrices, and if we allow the non-zero entries to be either 1 or -1 , we have the elements of B_n . Hence B_n has $2^n n!$ elements. The groups \mathfrak{S}_n and B_n are the Weyl groups for the root systems of types A_{n-1} and C_n , respectively.

Let $\sigma, \rho \in B_n$. We set that we multiply the elements of B_n , or \mathfrak{S}_n , from left to right. Thus the image of $i \in [\pm n]$ under $\sigma \in B_n$ is written $(i)\sigma$ to accomplish our convention of composition of maps, left to right, in B_n , $(i)\sigma\rho := [(i)\sigma]\rho$. We call to $[a_1 a_2 \dots a_n]$, where $a_i = (i)\sigma$ for $i \in [n]$, the window notation of σ , and write $\sigma = [a_1 a_2 \dots a_n]$. Since we multiply the elements of B_n , or \mathfrak{S}_n , from left to right, the elements of B_n act on vector in \mathbb{Z}^n on the right. Given a vector $v \in \mathbb{Z}^n$, we have that s_i , with $i \in [n-1]$, acts on v swapping the i -th and the $(i+1)$ -th entries, and s_n acts on v , vs_n , changing the sign of the last entry. Note that the window notation of $s_i\sigma$ is obtained after applying s_i to the window notation of σ , if we see it as a vector. Since we multiply the elements of B_n , or \mathfrak{S}_n , from left to right, they act on vectors on the right. Ignoring signs, $v\sigma = (v_{(1)\sigma^{-1}}, \dots, v_{(n)\sigma^{-1}})$, with $v = (v_1, \dots, v_n)$. The i -th letter of $v\sigma$ changes its sign if and only if

\bar{i} appears in σ . Hence $v\sigma = (sgn((1)\sigma^{-1})v_{|(1)\sigma^{-1}|}, \dots, sgn((n)\sigma^{-1})v_{|(n)\sigma^{-1}|})$, where $sgn(x) = 1$ if x is positive and -1 if x is negative, for $x \in [\pm n]$.

Example 1. Consider $v = (1, 2, 3) \in \mathbb{Z}^3$ and $\sigma = [2\bar{3}1] = s_2s_3s_1 = [(1)s_2s_3s_1, (2)s_2s_3s_1, (3)s_2s_3s_1] \in B_3$. So

$$\begin{aligned} (1, 2, 3)\sigma &= (1, 2, 3)s_2s_3s_1 = (1, 3, 2)s_3s_1 = (1, 3, \bar{2})s_1 = (3, 1, \bar{2}) \\ &= (sgn((1)\sigma^{-1})v_{|(1)\sigma^{-1}|}, sgn((2)\sigma^{-1})v_{|(2)\sigma^{-1}|}, sgn((3)\sigma^{-1})v_{|(3)\sigma^{-1}|}) \\ &= (1 \times 3, 1 \times 1, -1 \times 2). \end{aligned}$$

2.1. Bruhat order on B_n . The length of $\sigma \in B_n$, $\ell(\sigma)$, is the least number of generators of B_n needed to go from $[1\ 2\ \dots\ n]$, the identity map, to σ . Any expression of σ as a product of $\ell(\sigma)$ generators of B_n is called reduced. We say that two letters of the window notation of σ form an inversion if the bigger letter appears first. Next proposition gives a way to compute $\ell(\sigma)$ that only requires to look at the window notation of σ . This is a variation of the length formula presented on [4, Proposition 8.1.1], where the authors consider the usual ordering of the alphabet $[\pm n]$ and the generator that changes the sign of an entry of the window notation acts on the first entry instead of the last one.

Proposition 1. Consider $\sigma \in B_n$. Then

$$\ell(\sigma) = \#\{\text{inversions of } \sigma\} + \sum_{\bar{i} \text{ appears in } \sigma} (n + 1 - i).$$

The (signed) permutation $\sigma = [2\bar{3}1]$ has two inversions: 2, 1 and $\bar{3}$, 1 and $\ell(\sigma) = 3$.

Remark.

- If \bar{i} does not appear in the window presentation of σ , for all $i \in [n]$, we may identify σ , in one-line notation, with $(1)\sigma \dots (n)\sigma \in \mathfrak{S}_n$ and $\ell(\sigma) = \#\{\text{inversions of } \sigma\}$ [4, Proposition 1.5.2].
- We have that $\ell(s_i\sigma) > \ell(\sigma)$ if $i = n$ and $\sigma(n)$ is positive, or, $i \neq n$ and $\sigma(i) < \sigma(i + 1)$.

The Bruhat order on the set of the elements of B_n can be defined in the following way:

Definition 1. [4] Let $w = \sigma_1 \dots \sigma_{\ell(w)}$, where σ_i are generators of B_n , and u be two elements in B_n . Then $u \leq w$ in the Bruhat order if


$$\exists 1 \leq i_1 < i_2 \cdots < i_{\ell(u)} \leq \ell(w) \text{ such that } u = \sigma_{i_1} \sigma_{i_2} \cdots \sigma_{i_{\ell(u)}}.$$

By definition, if $u \leq w$ then $\ell(u) \leq \ell(w)$, but the reverse is not true. If $\sigma(n)$ is positive and $i = n$, or, $\sigma(i) < \sigma(i + 1)$ and $i \neq n$, we can also say that $s_i \sigma > \sigma$.


The combinatorics of crystal graphs in type C and the Bruhat order combinatorics on B_n and cosets are strongly related. In subsections 2.3 and 2.4, we give a tableau criteria for the Bruhat order on B_n and on cosets, respectively. For this aim, we recall Kashiwara-Nakashima (KN) tableaux in type C and define symplectic key tableau.

2.2. Kashiwara-Nakashima tableau in type C . This subsection focuses on the notion of symplectic tableaux introduced by Kashiwara and Nakashima to label the vertices of the type C crystal graphs [13], which are a variation of the De Concini tableaux [6]. (See [29] for more details.)

A vector $\lambda = (\lambda_1, \dots, \lambda_n) \in \mathbb{Z}^n$ is a partition of $|\lambda| = \sum_{i=1}^n \lambda_i$ if $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n \geq 0$. The Young diagram of shape λ is an array of boxes, left justified, in which the i -th row, from top to bottom, has λ_i boxes. We identify a partition with its Young diagram. For example, the Young diagram

of shape $\lambda = (2, 2, 1)$ is .

Given μ and ν two partitions with $\nu \leq \mu$ entrywise, we write $\nu \subseteq \mu$. The Young diagram of shape μ/ν is obtained after removing the boxes of the Young diagram of ν from the Young diagram of μ . For example, the Young

diagram of shape $\mu/\nu = (2, 2, 1)/(1, 0, 0)$ is .

Definition 2. Let $\nu \subseteq \mu$ be two partitions and A a completely ordered alphabet. A semistandard skew tableau of shape μ/ν on the alphabet A is a filling of the diagram μ/ν with letters from A , such that the entries are strictly increasing in each column and weakly increasing in each row. When $|\nu| = 0$ then we obtain a semistandard Young tableau of shape μ .

Denote by $SSYT(\mu/\nu, A)$ the set of all semistandard Young skew tableaux T of shape μ/ν , with entries in A . When $A = [n]$ we write $SSYT(\mu/\nu, n)$.

When considering tableaux with entries in $[\pm n]$, it is usual to have some extra conditions besides being semistandard. We will use a family of tableaux known as *Kashiwara-Nakashima* tableaux. From now on we consider tableaux on the alphabet $[\pm n]$.

A *column* is a strictly increasing sequence of numbers in $[\pm n]$ and it is usually displayed vertically. A column is said to be *admissible* if the following *one column condition* (1CC) holds for that column:

Definition 3 (1CC). Let C be a column. The 1CC holds for C if for all pairs i and \bar{i} in C , where i is in the a -th row counting from the top of the column, and \bar{i} in the b -th row counting from the bottom, we have $a + b \leq i$.

If a column C satisfies the 1CC then C has at most n letters.

If 1CC doesn't hold for C we say that C breaks the 1CC at z , where z is the minimal positive integer such that z and \bar{z} exist in C and there are more than z numbers in C with absolute value less or equal than z .

Example 2. The column $\begin{array}{|c|} \hline 1 \\ \hline 2 \\ \hline \bar{1} \\ \hline \end{array}$ breaks the 1CC at 1.

The following definition states conditions to when C can be split:

Definition 4. Let C be a column and let $I = \{z_1 > \dots > z_r\}$ be the set of unbarred letters z such that the pair (z, \bar{z}) occurs in C . The column C can be split when there exists a set of r unbarred letters $J = \{t_1 > \dots > t_r\} \subseteq [n]$ such that:

- (1) t_1 is the greatest letter of $[n]$ satisfying $t_1 < z_1$, $t_1 \notin C$, and $\bar{t}_1 \notin C$,
- (2) for $i = 2, \dots, r$, we have that t_i is the greatest letter of $[n]$ satisfying $t_i < \min(t_{i-1}, z_i)$, $t_i \notin C$, and $\bar{t}_i \notin C$.

The 1CC holds for a column C if and only if C can be split [29, Lemma 3.1]. If C can be split then we define right column of C , rC , and the left column of C , ℓC , as follows:

- (1) rC is the column obtained by changing in C , \bar{z}_i into \bar{t}_i for each letter $z_i \in I$ and by reordering if necessary,
- (2) ℓC is the column obtained by changing in C , z_i into t_i for each letter $z_i \in I$ and by reordering if necessary.

If C is admissible then $\ell C \leq C \leq rC$ by entrywise comparison. If C doesn't have symmetric entries, then C is admissible and $\ell C = C = rC$. In the next definition we give conditions for a column C to be coadmissible.

Definition 5. We say that a column C is coadmissible if for every pair i and \bar{i} on C , where i is on the a -th row counting from the top of the column, and \bar{i} on the b -th row counting from the top, then $b - a \leq n - i$.

Note that, unlike in Definition 3, in the last definition b is counted from the top of the column.

Given an admissible column C , consider the function Φ that sends C to the column of the same size in which the unbarred entries are taken from ℓC and the barred entries are taken from rC . The column $\Phi(C)$ is a coadmissible column and the algorithm to form $\Phi(C)$ from C is reversible [18, Section 2.2]. In particular, every column on the alphabet $[n]$ is simultaneously admissible and coadmissible.

Example 3. Let $C = \begin{array}{|c|} \hline 2 \\ \hline 3 \\ \hline 3 \\ \hline \end{array}$ be an admissible column. Then $\ell C = \begin{array}{|c|} \hline 1 \\ \hline 2 \\ \hline 3 \\ \hline \end{array}$ and $rC = \begin{array}{|c|} \hline 2 \\ \hline 3 \\ \hline 1 \\ \hline \end{array}$. So $\Phi(C) = \begin{array}{|c|} \hline 1 \\ \hline 2 \\ \hline 1 \\ \hline \end{array}$ is coadmissible.

Let T be a skew tableau with all of its columns admissible. The split form of a skew tableau T , $spl(T)$, is the skew tableau obtained after replacing each column C of T by the two columns $\ell C rC$. The tableau $spl(T)$ has double the amount of columns of T .

Definition 6. A semistandard skew tableau T is a Kashiwara-Nakashima (KN) skew tableau if its split form is a semistandard skew tableau. We define $\mathcal{KN}(\mu/\nu, n)$ to be the set of all KN tableau of shape μ/ν in the alphabet $[\pm n]$. When $\nu = 0$ we obtain $\mathcal{KN}(\mu, n)$.

Example 4. The split of the tableau $T = \begin{array}{|c|c|} \hline 2 & 2 \\ \hline 3 & 3 \\ \hline 3 & \\ \hline \end{array}$ is the tableau $spl(T) =$

$\begin{array}{|c|c|c|c|} \hline 1 & 2 & 2 & 2 \\ \hline 2 & 3 & 3 & 3 \\ \hline 3 & 1 & & \\ \hline \end{array}$. Hence $T \in \mathcal{KN}((2, 2, 1), 3)$.

If T is a tableau without symmetric entries in any of its columns, i.e., for all $i \in [n]$ and for all columns C in T , i and \bar{i} do not appear simultaneously in the entries of C , then in order to check whether T is a KN tableau it is enough to check whether T is semistandard in the alphabet $[\pm n]$. In particular $SSYT(\mu/\nu, n) \subseteq \mathcal{KN}(\mu/\nu, n)$.

The weight of a word w on the alphabet $[\pm n]$ is defined to be the vector $\text{wt}(w) \in \mathbb{Z}^n$ where the entry i is obtained by adding the multiplicity of the letter i and subtracting the multiplicity of the letter \bar{i} , for $i \in [n]$. If T is a skew tableau, the column reading of T , $cr(T)$, is the word read in T in the Japanese way, column reading top to bottom and right to left. The length of w is the total number of letters in w . The weight of a KN tableau T is the vector $\text{wt } T := \text{wt}(cr(T)) = (t_1 - t_{\bar{1}}, t_2 - t_{\bar{2}}, \dots, t_n - t_{\bar{n}}) \in \mathbb{Z}^n$, where t_α is the number of α 's in T , with $\alpha \in [\pm n]$.

Example 5. Let $T = \begin{array}{|c|c|} \hline 2 & 2 \\ \hline 3 & 3 \\ \hline 3 & \\ \hline \end{array}$ and $n = 3$. Then $cr(T) = 2323\bar{3}$ and $\text{wt}(T) = \text{wt}(cr(T)) = (0, 2, 1)$.

In Section 3.2, we recall a way to go from a word in the alphabet $[\pm n]$ to a KN tableau, the Baker-Lecouvey insertion.

2.3. Key tableaux in type C and the Bruhat order on B_n .

Definition 7. A key tableau in type C_n is a KN tableau in $\mathcal{KN}(\lambda, n)$, in which the set of elements of each column is contained in the set of elements of the previous column and the letters i and \bar{i} do not appear simultaneously as entries, for any $i \in [n]$.

Example 6. The KN tableau $T = \begin{array}{|c|c|} \hline 2 & 2 \\ \hline 3 & \bar{1} \\ \hline \bar{1} & \\ \hline \end{array}$ is a key tableau.

The set of key tableaux in type A is the subset of the key tableaux in type C consisting of the tableaux having only positive entries, hence they are SSYT for the alphabet $[n]$.

Every vector v of \mathbb{Z}^n is in the B_n -orbit of exactly one partition, λ_v , which is the one obtained by sorting the absolute values of all entries of v . Given a partition $\lambda \in \mathbb{Z}^n$, the B_n -orbit of λ is the set $\lambda B_n := \{\lambda\sigma \mid \sigma \in B_n\}$. For instance, the vector $v = (1, \bar{3}, 0, 3, \bar{2})$ is in the B_5 -orbit of $\lambda = (3, 3, 2, 1, 0)$.

Proposition 2. Let λ be a partition and $v \in \lambda B_n$. There is exactly one key tableau $K(v)$ whose weight is v . In addition, the shape of the key tableau $K(v)$ is λ . When $v = \lambda$, $K(\lambda)$ is the only KN tableau of weight and shape λ , also called Yamanouchi tableau of shape λ .

Proof: Existence: Given $v = (v_1, \dots, v_n) \in \mathbb{Z}^n$ there exists a key tableau K of weight v by putting in the first $|v_i|$ columns the letter i if $v_i \geq 0$ or \bar{i} if $v_i \leq 0$, and then sorting the columns properly. Clearly the columns of K are nested and it is a KN tableau without symmetric entries, hence it is a key tableau. Also, its shape is $\lambda_v = \lambda$.

Uniqueness: Since the key tableau doesn't have symmetric entries then, for all $i \in [n]$, we have that in K the letter $\text{sgn}(v_i)i$ appears $|v_i|$ times in its entries. In order to the columns of K be nested we have that these $|v_i|$ entries appear in the first $|v_i|$ columns, hence we have determined exactly which letters appear in which column of K and now we just have to order them correctly. So the key tableau K with weight v is unique. When $v = \lambda$, $K(\lambda)$ has only i 's in the row i , for $i \in [n]$. ■

Example 7. Let $v = (1, \bar{3}, 0, 3, \bar{2})$. Then $K(v) =$

1	4	4
4	$\bar{5}$	$\bar{2}$
$\bar{5}$	$\bar{2}$	
$\bar{2}$		

.

Hence there is a bijection between vectors in λB_n and the key tableaux in type C on the alphabet $[\pm n]$ with shape λ , given by the map $v \mapsto K(v)$. If $\sigma \in B_n$ we put $K(\sigma) := K(\Lambda_{(n)}\sigma)$, where $\Lambda_{(n)} = (n, n-1, \dots, 1)$ is the staircase partition in \mathbb{Z}^n . One has a natural bijection between B_n and the B_n -orbit of $\Lambda_{(n)}$.

Proposition 3. If $\sigma \in B_n$ has the letter α in the j -th position then α appears in the first $n+1-j$ columns of the corresponding key tableau, $K(\sigma)$.

Proof: Put $\Lambda := \Lambda_{(n)}$. Ignoring signs, $\Lambda\sigma = (\Lambda_{(1)\sigma^{-1}}, \dots, \Lambda_{(n)\sigma^{-1}})$, with $\Lambda = (n, \dots, 1)$ and the i -th letter of $\Lambda\sigma$ has negative sign if and only if \bar{i} appears in σ . If α is positive, then in the position α of $\Lambda\sigma$ will appear $\Lambda_j = n+1-j$. If α is negative, then in the position $-\alpha$ will appear $\bar{\Lambda}_j = \overline{n+1-j}$. ■

We now append 0 to the alphabet $[\pm n]$, obtaining $[\pm n] \cup \{0\}$, where $n < 0 < \bar{n}$, and, for all $\sigma \in B_n$, we put $(0)\sigma := 0$. Given an element $\sigma \in B_n$ consider the map

$$[\pm n] \cup \{0\} \times [\pm n] \cup \{0\} \rightarrow \mathbb{N}_0$$

$$(i, j) \mapsto |\{a \leq i : \sigma(a) \geq j\}| := \sigma[i, j].$$

This map, originally defined in [4], produces a table which is related to key tableaux in type C . See example below:

Example 8. Let $\sigma = [\overline{3} \overline{1} 2 4]$. Then $(4, 3, 2, 1)\sigma = (\overline{3}, 2, \overline{4}, 1)$ and

$$K(\sigma) = \begin{array}{|c|c|c|c|} \hline 2 & 2 & \overline{3} & \overline{3} \\ \hline 4 & \overline{3} & \overline{1} & \\ \hline \overline{3} & \overline{1} & & \\ \hline \overline{1} & & & \\ \hline \end{array}$$

The family of numbers $\sigma[i, j]$ where $(i, j) \in [\pm n] \cup \{0\} \times [\pm n] \cup \{0\}$ originates the following table, where i indexes the columns, left to right, and j indexes the rows, top to bottom. We add a row of zeros at the bottom for convenience:

	1	2	3	4	0	$\overline{4}$	$\overline{3}$	$\overline{2}$	$\overline{1}$
1	1	2	3	4	5	6	7	8	9
2	1	2	3	4	5	6	7	7	8
3	1	2	2	3	4	5	6	6	7
4	1	2	2	3	4	5	6	6	6
0	1	2	2	2	3	4	5	5	5
$\overline{4}$	1	2	2	2	2	3	4	4	4
$\overline{3}$	1	2	2	2	2	2	3	3	3
$\overline{2}$	0	1	1	1	1	1	2	2	2
$\overline{1}$	0	1	1	1	1	1	1	1	1
	0	0	0	0	0	0	0	0	0

To go from the table to the key tableau note that, for $i \in [n]$, the i -th column of the table encodes the $(n+1-i)$ -th column of the tableau, in the sense that if we look to the i -th column of the table, from bottom to top, if the entry of the table increases in one unity then the index of the row associated to that entry exists in the $(n+1-i)$ -th column of the tableau. Knowing the entries in a column of a tableau, its ordering is unique. The columns of the tableau constructed this way are nested because the indices in which the column i increases are $(j)\sigma$, for $j \leq i$. So the tableau taken from the table is the key tableau $K(\sigma)$. It is also possible to construct the table from the key tableau and that we only need the first n columns of the table.

We then have the following result:

Proposition 4. Consider $\sigma, \rho \in B_n$. $K(\sigma) \geq K(\rho)$ entrywise if and only if $\sigma[i, j] \geq \rho[i, j]$, where $i \in [n]$, and $j \in [\pm n]$.

In [4, Theorem 8.1.8] it is proved that, for $\sigma, \rho \in B_n$, $\sigma \leq \rho$ in the Bruhat order if and only if $\sigma[i, j] \leq \rho[i, j]$ for all $i, j \in [\pm n]$. But the result in [4, Theorem 8.1.7] implies that we only need to compare $\sigma[i, j]$ and $\rho[i, j]$ for

$i \in [n]$. Henceforth, we have the following criterion for the Bruhat order on B_n :

Theorem 5. *Consider $\sigma, \rho \in B_n$. $K(\sigma) \geq K(\rho)$ entrywise if and only if $\sigma \geq \rho$ in the Bruhat order.*

Remark. *In [4, Chapter 8.1] the authors use the same alphabet as here, but with the usual ordering on the integers. So, to translate the results from there to here, it is needed to apply the ordering isomorphism defined by: $i \mapsto \overline{n-i+1}$ if $i \in [n]$; $i \mapsto n+i+1$ if $i \in -[n]$; $0 \mapsto 0$. Using the usual ordering, the authors give a tableau criterion for the Bruhat order in Exercise 6, pp. 287–288, which is effectively the transpose version of the tableau criterion presented here. Also note that the generators used in [4, Chapter 8.1] are the same used here, although with different indexation. Our generator s_i corresponds to the generator s_{n-i} in [4, Chapter 8.1], for all $i \in [n]$.*

2.4. The Bruhat order on cosets of B_n . Consider a partition $\lambda \in \mathbb{Z}^n$. Let $W_\lambda = \{\rho \in B_n \mid \lambda\rho = \lambda\}$ be the stabilizer of λ , under the action of B_n . Since λ is a partition, W_λ is a subgroup of B_n generated by some of the generators of B_n . Let $J \subseteq [n]$ be the set of the indices of the generators of W_λ , i.e. $W_\lambda = \langle s_j, j \in J \rangle$, and J^c the complement of this set in $[n]$. Let $W_\lambda \backslash B_n = \{W_\lambda\sigma : \sigma \in B_n\}$ be the set of right cosets of B_n determined by the subgroup W_λ . Each coset $W_\lambda\sigma$ returns a unique vector v when acting on λ , and has a unique minimal length element σ_v , such that $v = \lambda\sigma_v$. Reciprocally, given a vector $v \in \lambda B_n$, there is a unique minimal length element $\sigma_v \in B_n$ such that $v = \lambda\sigma_v$. We have then a bijection between the vectors in λB_n and the right cosets of B_n , determined by the subgroup W_λ , given by $v \mapsto W_\lambda\sigma_v$. The set J^c detects the minimal length coset representatives of $W_\lambda \backslash B_n$: σ is a minimal coset representative of $W_\lambda \backslash B_n$ if and only if all its reduced decompositions start with a generator $s_i \in J^c$ [4]. However key tableaux, $K(v)$, $v \in \lambda B_n$, may be used to explicitly construct the minimal length coset representatives of $W_\lambda \backslash B_n$. Given a vector $v \in \lambda B_n$, we show that there is a unique minimal length element $\sigma_v \in B_n$ such that $v = \lambda\sigma_v$ and we show how to obtain σ_v explicitly. The next proposition is a generalization of what Lascoux does in [15] for vectors in \mathbb{N}^n (hence $\sigma_v \in \mathfrak{S}_n$).

Proposition 6. *Let $v \in \lambda B_n$ and T the tableau obtained after adding the*

column $C = \begin{array}{|c|} \hline 1 \\ \hline 2 \\ \hline \vdots \\ \hline n \\ \hline \end{array}$ to the left of $K(v)$. The minimal length element $\sigma \in$

B_n , modulo W_λ , is given the reading word of T where entries with the same absolute value are read just once.

Proof: Consider $\lambda = (\lambda_1, \dots, \lambda_n)$. In this proof we will write λ as $(\lambda_1^{a_{\lambda_1}}, (\lambda_1 - 1)^{a_{\lambda_1 - 1}}, \dots, 1^{a_1}, 0^{a_0})$ where a_i is the multiplicity of i in λ , for $0 \leq i \leq \lambda_1$.

Note that $\sum_{i=0}^{\lambda_1} a_i = n$.

Let $\sigma = [\alpha_1 \dots \alpha_n] \in B_n$ read from T . Let's prove that α_j appears λ_j times in $K(v)$: If $j = 1$ then α_1 appears in all columns of $K(v)$, because it was the first letter read and the columns are nested. Hence it appears λ_1 times. Also, the $|\alpha_1|$ -th entry of $\lambda\sigma$ is $\text{sgn}(\alpha_1)\lambda_1$ which is the weight of $|\alpha_1|$ in $K(v)$. For $j > 1$, proceeding inductively, we have that α_j appears in all columns of $K(v)$ not fully occupied by α_i , with $i < j$, hence it appears λ_j times. Also, the $|\alpha_j|$ -th entry of $\lambda\sigma$ is $\text{sgn}(\alpha_j)\lambda_j$, which is the weight of $|\alpha_j|$ in $K(v)$. This makes sense even if $\lambda_j = 0$. So we have that $\lambda\sigma = v$.

We only have to see that σ is the minimal length element of the set $\{\rho \in B_n \mid \lambda\rho = v\}$. The subset of elements B_n that applied to λ returns v is the coset $W_\lambda\sigma$. Looking at σ , this allows us to swap α_i and α_j in σ if $\lambda_i = \lambda_j$ and to change the sign of α_i if $\lambda_i = 0$. Since for each column the reading to obtain σ is ordered from the least to the biggest, we have that between these elements of B_n , σ has minimal number of inversions and the letter α_j is unbarred if $\lambda_j = 0$ because α_j will only be added to σ when reading the column C . Hence, by Proposition 1, σ is the minimal length element of $W_\lambda\sigma$. \blacksquare

Given a partition $\lambda \in \mathbb{Z}^n$ we identify each coset $W_\lambda\sigma$ with its minimal length representative σ_v , where $v = \lambda\sigma \in \lambda B_n$. Under this identification, we now induce the Bruhat order in the B_n -orbit of λ and in the coset space of $W_\lambda \setminus B_n$.

Definition 8. Consider the vectors $v, w \in \lambda B_n$, where λ is a partition. We say that $v \leq w$, in the Bruhat order, if $\sigma_v \leq \sigma_w$.

Let $v \in \lambda B_n$. If $K := K(v)$ is the key tableau with weight v , consider the tableau \tilde{K} obtained from K after erasing the minimal number of columns in order to have a tableau with no duplicated columns. Call \tilde{v} and $\tilde{\lambda}$ to the

weight and the shape of \widetilde{K} , respectively. If K and K' are two key tableaux with shape λ , we have that $K \geq K'$ (by entrywise comparison) if and only if $\widetilde{K} \geq \widetilde{K}'$. Note that to recover K from \widetilde{K} we just have to know λ , and that $\widetilde{K} = K(\tilde{v})$.

It is also possible to obtain \tilde{v} from v without having to look at the key tableau. If i is positive, i and \bar{i} do not appear in v but $i+1$ or $\overline{i+1}$ appears then change all appearances of $i+1$ and $\overline{i+1}$ to i and \bar{i} , respectively, and repeat this as many times as possible, obtaining the vector \tilde{v} . The set of the absolute values of its entries is a set of consecutive integers starting either in 0 or 1. Hence the key tableau associated to it doesn't have repeated columns.

Due to Proposition 6 we have that $\sigma_{\tilde{v}} = \sigma_v$ and $\tilde{v} = \widetilde{\lambda}_v \sigma_{\tilde{v}} = \widetilde{\lambda}_v \sigma_v$.

Example 9. Consider $v = (1, 0, \bar{3}, 3, \bar{5}) \in (5, 3, 3, 1, 0)B_5$. Hence $K(v) =$

1	4	4	5	5
4	5	5		
5	3	3		
3				

has shape $\lambda = (5, 3, 3, 1, 0)$, weight v and $\sigma_v = [\bar{5} 4 \bar{3} 1 2]$. Now

note that $\tilde{v} = (1, 0, \bar{2}, 2, \bar{3})$, hence $K(\tilde{v}) =$

1	4	5
4	5	
5	3	
3		

$= \widetilde{K(v)}$ has shape

$(3, 2, 2, 1, 0) = \widetilde{\lambda}$ and $\sigma_{\tilde{v}} = [\bar{5} 4 \bar{3} 1 2] = \sigma_v$.

Recall J and J^c defined above. Note that the set J is the same for λ and $\widetilde{\lambda}$. If $i \in J^c$ and $i = n$ then all entries of λ are different from 0, which implies $K(v)$ (and $\widetilde{K(v)}$) having columns of length n ; if $i \in J^c$ and $i < n$ then $\lambda_i > \lambda_{i+1}$, hence $K(v)$ will have exactly i rows with length greater than λ_{i+1} , hence $K(v)$ (and $\widetilde{K(v)}$) will have columns of length i . Since $\widetilde{K(v)}$ doesn't have repeated columns, J^c have exactly the information of what columns length exist in $\widetilde{K(v)}$. Theorem 3BC of Proctor's Ph.D. thesis [27] states that given a partition λ there is a poset isomorphism between the poset formed by the key tableaux of shape $\widetilde{\lambda}$ (ordered by entrywise comparison) and the poset formed by the Bruhat order in the vectors of the orbit $\widetilde{\lambda}B_n = \{\widetilde{\lambda}\sigma : \sigma \in B_n\}$.

The following theorem gives a tableau criterion for the Bruhat order on vector the same B_n -orbit and for the corresponding B_n -coset space.

Theorem 7. Let $v, u \in \lambda B_n$. Then $\sigma_v \leq \sigma_u$ if and only if $K(v) \leq K(u)$.

Proof: We have that

$$\sigma_v \leq \sigma_u \stackrel{(1)}{\Leftrightarrow} v \leq u \stackrel{(2)}{\Leftrightarrow} \tilde{v} \leq \tilde{u} \stackrel{(3)}{\Leftrightarrow} K(\tilde{v}) \leq K(\tilde{u}) \Leftrightarrow \widetilde{K(v)} \leq \widetilde{K(u)} \stackrel{(4)}{\Leftrightarrow} K(v) \leq K(u),$$

where (1) holds by Definition 8. Note that in (2) we also need to record λ , because it is needed in (4) to recover the shape of $K(v)$ from the shape $\widetilde{K(v)}$. Finally the equivalence (3) is an application of Theorem 3BC of Proctor's Ph.D. thesis [27]. \blacksquare

$$K(3, \bar{3}, 0, 0, \bar{2}) = \begin{array}{|c|c|c|} \hline 1 & 1 & 1 \\ \hline \bar{5} & \bar{5} & 2 \\ \hline 2 & 2 & \\ \hline \end{array} \leq K(\bar{3}, 2, 0, \bar{3}, 0) = \begin{array}{|c|c|c|} \hline 2 & 2 & \bar{4} \\ \hline \bar{4} & \bar{4} & \bar{1} \\ \hline \bar{1} & \bar{1} & \\ \hline \end{array} \text{ and } \sigma_v = [1\bar{2}\bar{5}34] \leq \\ \sigma_u = [4\bar{1}235]$$

3. Type C crystal graphs and symplectic plactic monoid

We recall two equivalence relations of words in the alphabet $[\pm n]$, the type C Knuth equivalence, or (symplectic) plactic equivalence, and the (symplectic) coplactic equivalence. On the basis of these two equivalence relations is the Robinson-Schensted type C correspondence, in which each word is uniquely parametrized by a KN tableau and an oscillating tableau of the same final shape. This bijection has a natural interpretation in terms of crystal connectivity and crystal isomorphic connected components in Kashiwara theory of crystal graphs [5, 12, 18, 19]. For this aim and reader convenience, we begin to recall the Sheats symplectic *jeu de de taquin* and Baker-Lecouvey insertion.

3.1. Sheats symplectic *jeu de de taquin*. The symplectic *jeu de de taquin* [18, 29] is a procedure that allows us to change the shape of a KN skew tableau and eventually rectify it.

To explain how the symplectic *jeu de de taquin* behaves, we need to look to how it works on 2-column KN skew tableaux. Let T be a 2-column KN skew tableau with splittable columns C_1 and C_2 such that C_1 has an empty cell.

Consider the tableau $spl(T)$ such that the columns ℓC_1 and rC_1 have an empty cell in the same row as C_1 . Call α to the entry under the empty cell of rC_1 and β to the entry right of the empty cell of rC_1 .

If $\alpha \leq \beta$ or β does not exist, then the empty cell of T will change its position with the cell beneath it. This is a vertical slide.

If the slide is not vertical, then it is horizontal. So we have $\alpha > \beta$ or α does not exist. Call C'_1 and C'_2 to the columns after the slide. In this case we have two subcases, depending on the sign of β :

- (1) If β is barred we are moving a barred letter from ℓC_2 to rC_1 . Remember that ℓC_2 has the same barred part as C_2 and that rC_1 has the same barred part as $\Phi(C_1)$. So, looking at T , we will have an horizontal slide of the empty cell, $C'_2 = C_2 \setminus \{\beta\}$ and $C'_1 = \Phi^{-1}(\Phi(C_1) \cup \{\beta\})$. In a sense, β went from C_2 to $\Phi(C_1)$.
- (2) If β is unbarred we have a similar story, but this time β will go from $\Phi(C_2)$ to C_1 , hence $C'_1 = C_1 \cup \{\beta\}$ and $C'_2 = \Phi^{-1}(\Phi(C_2) \setminus \{\beta\})$. Although in this case it may happen that C'_1 is no longer admissible. In this case, if the 1CC breaks at i , we erase both i and \bar{i} from the column and remove a cell from the bottom and from the top column, and place all the remaining cells orderly.

Eventually the empty cell will be a cell such that α and β do not exist. In this case we redefine the shape to not include this cell and the *jeu de taquin* ends. An entry of the tableau without cells under it or to the right of it is called an inner corner.

Given a KN skew tableau T of shape μ/ν , the rectification of T consists in playing the *jeu de taquin* until we get a tableau of shape λ , for some partition λ . The rectification is independent of the order in which the inner corners of ν are filled [18, Corollary 6.3.9].

Example 10. Consider the KN skew tableau $T = \begin{array}{|c|} \hline 2 \\ \hline 1 & 3 \\ \hline 2 & 1 \\ \hline \end{array}$. We want to rectify it via symplectic *jeu taquin*. We start by splitting and conclude that the first two slides are vertical, obtaining $\begin{array}{|c|c|c|c|} \hline 1 & 1 & 2 & 2 \\ \hline 2 & 2 & 3 & 3 \\ \hline 1 & 1 & & \\ \hline \end{array}$. Now we do an horizontal slide in which we take $\bar{1}$ from the second column of T and adding it to the coadmissible column of the first column of T , obtaining the tableau $\begin{array}{|c|} \hline 2 & 2 \\ \hline 3 & 3 \\ \hline 3 \\ \hline \end{array}$.

Remark. If the columns C_1 and C_2 do not have negative entries then the symplectic *jeu de taquin* coincides with the *jeu de taquin* known for SSYT.

3.2. Baker-Lecouvey insertion. The Baker-Lecouvey insertion [3, 18] is a bumping algorithm that given a word in the alphabet $[\pm n]$ returns a KN

tableau. Let w be a word in the alphabet $[\pm n]$, we call $P(w)$ to the tableau obtained after inserting w . This insertion is similar to the usual column insertion for SSYT tableau. In fact both have the same behavior unless one the following three cases happens:

Suppose that we are inserting the letter α in the column C of the KN tableau and

- (1) $\bar{y} \in C$ is the smallest letter bigger or equal than α and $y \in C$, for some $y \in [n]$: there is in C a maximal string of consecutive decreasing integers $y, y-1, \dots, u+1$ starting in the entry y in the column C . Then the bump consists of replacing the entry \bar{y} with α and subtracting 1 to the entries $y, y-1, \dots, u+1$. The entry \bar{u} is then inserted in the next column to the right. This is known as the Type I special bump.
- (2) if $\alpha = x$ and $\bar{x} \in C$, for some $x \in [n]$: there is a maximal string of consecutive decreasing entries $\bar{x}, \overline{x+1}, \dots, \overline{v-1}$ starting in the entry \bar{x} in C . Let β be the next entry above $\overline{v-1}$. Then we have two subcases:
 - (a) If $v \leq \beta \leq \overline{v+1}$ then suppose δ is the smallest entry in C which is bigger or equal than v . Then this bump consists of deleting the entry \bar{x} , shifting the entries $\overline{x+1}, \dots, \overline{v-1}$ down one position, inserting \bar{v} where $\overline{v-1}$ was, and replacing δ with v . The entry δ is then bumped into the next column. This is known as the Type IIa special bump.
 - (b) If $\beta \leq v-1$ or β doesn't exist then there is a maximal string (possibly empty) of consecutive integers $v-1, \dots, u+1$ above the entry $\overline{v-1}$. The string is not empty only when $\beta = v-1$, or else the string is empty and $u = v-1$. The bump consists of deleting the entry \bar{x} , shifting the entries $\overline{x+1}, \dots, u+1$ down one position, and inserting an entry u where $u+1$ (or $\overline{v-1}$, if $\beta \neq v-1$) was. The entry \bar{u} is then bumped into the next column. This is known as the Type IIb special bump.
- (3) after adding α in the bottom of the column C , the 1CC breaks at i : then we will slide out the cells that contain \bar{i} and i via symplectic *jeu de taquin*.

In the case 3 of the Baker-Lecouvey insertion we will be removing a cell from the tableau instead of adding. Despite the length of $cr(P(w))$ might

be less than the length of w , the weight is preserved during Baker-Lecouvey insertion, $\text{wt}(w) = \text{wt}(P(w))$.

Remark. *The Baker-Lecouvey insertion is different from what we would have if we use the SSYT column insertion. However, if the word w doesn't have symmetric letters, then the insertion works just like the column insertion for SSYT. Apart from this case, if we were to use SSYT column insertion, the final tableau may not even be a KN tableau. For instance, consider the word $w = 2\bar{1}1$. The Baker-Lecouvey insertion of w creates the sequence of tableaux*

$\boxed{2} \quad \boxed{\frac{2}{\bar{1}}} \quad \boxed{\frac{2}{2} \frac{2}{2}} = P(2\bar{1}1)$. The SSYT column insertion of w results in the tableau $\boxed{\frac{1}{1} \frac{2}{1}}$, which is not a KN tableau because the first column is not admissible.

Example 11. *Consider the word $w = 23\bar{2}\bar{3}1$. We now insert all five letters of*

w , obtaining the following sequence of tableaux: $\boxed{2} \quad \boxed{\frac{2}{3}} \quad \boxed{\frac{2}{3} \frac{2}{2}} \quad \boxed{\frac{1}{3} \frac{\bar{1}}{3}} \quad \boxed{\frac{1}{3} \frac{1}{3} \frac{\bar{1}}{3}} =$

$P(w)$. Note that the insertion of the fourth letter, $\bar{3}$, causes a type I special bump on the first column and the insertion of the fifth letter, 1 , causes a type IIb special bump on the second column.

Proposition 8. [18, Corollary 6.3.9] *Let $T \in \mathcal{KN}(\mu/\nu, n)$. The tableau obtained after rectifying T via symplectic jeu de taquin coincides with $P(\text{cr}(T))$. Moreover, the insertion of $w = w_1 \dots w_k$, $P(w)$, is the rectification of the tableau with diagonal shape $\Lambda_{(n)}/\Lambda_{(n-1)}$ and column reading w .*

In particular we have that if we insert $\text{cr}(T)$ we obtain T again. This implies that during the insertion of $\text{cr}(T)$ the case 3 of the Baker-Lecouvey insertion cannot happen. In Example 11, we may conclude that $P(23\bar{2}\bar{3}1) = P(\text{cr}(P(23\bar{2}\bar{3}1))) = P(\bar{1}113\bar{3})$.

3.3. Robinson-Schensted type C correspondence, plactic and coplactic equivalence. Let $[\pm n]^*$ be the free monoid on the alphabet $[\pm n]$. The Robinson-Schensted type C correspondence [18, Theorem 5.2.2] is a combinatorial bijection between words $w \in [\pm n]^*$ and pairs (T, Q) where T is a KN tableau and Q is an oscillating tableau, a sequence of Young diagrams that record, by order, the shapes of the tableaux obtained while inserting w , whose final shape is the same as T . Every two consecutive shapes of the

oscillating tableau differ in exactly one cell and its length is the same of w . Since both the symplectic *jeu de taquin* and the Baker-Lecouvey insertion are reversible [3, 18], we have that every pair (T, Q) , with the same final shape, is originated by exactly one word. The Robinson-Schensted type C correspondence is the following map:

$$[\pm n]^* \rightarrow \bigsqcup_{\lambda} \mathcal{KN}(\lambda, n) \times \mathcal{O}(\lambda, n) : w \mapsto (P(w), Q(w))$$

where the union is over all partitions λ with at most n parts, and $\mathcal{O}(\lambda, n)$ is the set of all oscillating tableau with final shape λ and all shapes of the sequence have at most n rows.

Example 12. In Example 11, the word $w = 23\bar{2}31$ is associated to the pair

$$\left(\begin{array}{|c|c|c|} \hline 1 & 1 & \bar{1} \\ \hline 3 & & \\ \hline \bar{3} & & \\ \hline \end{array}, \begin{array}{|c|} \hline \\ \hline \end{array}, \begin{array}{|c|} \hline \\ \hline \end{array}, \begin{array}{|c|c|} \hline & \\ \hline & \\ \hline & \\ \hline & \\ \hline & \\ \hline & \\ \hline \end{array}, \begin{array}{|c|c|c|} \hline & & \\ \hline & & \\ \hline & & \\ \hline & & \\ \hline & & \\ \hline & & \\ \hline & & \\ \hline \end{array} \right).$$

Given $w_1, w_2 \in [\pm n]^*$, the relation $w_1 \sim w_2 \Leftrightarrow P(w_1) = P(w_2)$ defines an equivalence relation on $[\pm n]^*$ known as Knuth equivalence. The type C plactic monoid is the quotient $[\pm n]^* / \sim$ where each Knuth (plactic) class is uniquely identified with a KN tableau [16, 18]. The quotient $[\pm n]^* / \sim$ can also be described as the quotient of $[\pm n]^*$ by the elementary Knuth relations:

- K1: $\gamma\beta\alpha \sim \beta\gamma\alpha$, where $\gamma < \alpha \leq \beta$ and $(\beta, \gamma) \neq (\bar{x}, x)$ for all $x \in [n]$.
- K2: $\alpha\beta\gamma \sim \alpha\gamma\beta$, where $\gamma \leq \alpha < \beta$ and $(\beta, \gamma) \neq (\bar{x}, x)$ for all $x \in [n]$.
- K3: $y + 1\bar{y} + \bar{1}\beta \sim \bar{y}y\beta$, where $y < \beta < \bar{y}$ and $y \in [n - 1]$.
- K4: $\beta\bar{y}y \sim \beta y + 1\bar{y} + \bar{1}$, where $y < \beta < \bar{y}$ and $y \in [n - 1]$.
- K5: $w \sim w \setminus \{z, \bar{z}\}$, where $w \in [\pm n]^*$ and $z \in [n]$ are such that w is a non-admissible column that the $1CC$ breaks at z , and any proper factor of w is an admissible column.

Remark. It can be proved that given a word $w \in [\pm n]^*$, any proper factor is admissible if and only if any proper prefix of w is admissible. Thus, in order to be able to apply the Knuth relation K5 to a subword w' of w , we only need to check that all proper prefixes of w' are admissible, instead of all proper factors.

When Knuth relations are applied to subwords of a word, the weight is preserved while the length may not. Knuth relations can be seen as *jeu de taquin* moves on words or a diagonally shaped tableau, and each symplectic *jeu de taquin* slide preserves the Knuth class of the reading word of a tableau

[18, Theorem 6.3.8]. In Example 11 the words $23\bar{2}\bar{3}1$ and $\bar{1}113\bar{3}$ are Knuth related: $\bar{1}113\bar{3} \stackrel{K2}{\approx} \bar{1}131\bar{3} \stackrel{K2}{\approx} \bar{1}13\bar{3}1 \stackrel{K3}{\approx} 2\bar{2}3\bar{3}1 \stackrel{K1}{\approx} 23\bar{2}\bar{3}1$.

3.4. Crystal graphs in type C and coplactic monoid. Crystals were originally defined in quantum groups. Here we define them axiomatically associated to a root system Φ and a weight lattice Λ [5]. Let V be an Euclidean space with inner product $\langle \cdot, \cdot \rangle$. Fix a root system Φ with simple roots $\{\alpha_i \mid i \in I\}$ where I is an index set and a weight lattice $\Lambda \supseteq \mathbb{Z}\text{-span}\{\alpha_i \mid i \in I\}$. A Kashiwara crystal of type Φ is a nonempty set \mathfrak{B} together with maps [5]:

$$e_i, f_i : \mathfrak{B} \rightarrow \mathfrak{B} \sqcup \{0\} \quad \varepsilon_i, \varphi_i : \mathfrak{B} \rightarrow \mathbb{Z} \sqcup \{-\infty\} \quad wt : \mathfrak{B} \rightarrow \Lambda$$

where $i \in I$ and $0 \notin \mathfrak{B}$ is an auxiliary element, satisfying the following conditions:

- (1) if $a, b \in \mathfrak{B}$ then $e_i(a) = b \Leftrightarrow f_i(b) = a$. In this case, we also have $wt(b) = wt(a) + \alpha_i$, $\varepsilon_i(b) = \varepsilon_i(a) - 1$ and $\varphi_i(b) = \varphi_i(a) + 1$;
- (2) for all $a \in \mathfrak{B}$, we have $\varphi_i(a) = \langle wt(a), \frac{2\alpha_i}{\langle \alpha_i, \alpha_i \rangle} \rangle + \varepsilon_i(a)$.

The crystals we deal with are the ones of a $U_q(sp_{2n})$ -module. They are seminormal [5], and satisfy $\varphi_i(a) = \max\{k \in \mathbb{Z} \geq 0 \mid f_i^k(a) \neq 0\}$ and $\varepsilon_i(a) = \max\{k \in \mathbb{Z} \geq 0 \mid e_i^k(a) \neq 0\}$. An element $u \in \mathfrak{B}$ such that $e_i(u) = 0$ for all $i \in I$ is called a *highest weight element*. A *lowest weight element* is an element $u \in \mathfrak{B}$ such that $f_i(u) = 0$ for all $i \in I$. We associate with \mathfrak{B} a coloured oriented graph with vertices in \mathfrak{B} and edges labeled by $i \in I$: $b \xrightarrow{i} b'$ iff $b' = f_i(b)$, $i \in I$, $b, b' \in \mathfrak{B}$. This is the *crystal graph* of \mathfrak{B} .

If \mathfrak{B} and \mathfrak{C} are two seminormal crystals associated to the same root system, the *tensor product* $\mathfrak{B} \otimes \mathfrak{C}$ is also a seminormal crystal. As a set, we will have the Cartesian product $\mathfrak{B} \times \mathfrak{C}$, where its elements are denoted by $b \otimes c$, $b \in \mathfrak{B}$ and $c \in \mathfrak{C}$, with $wt(b \otimes c) = wt(b) + wt(c)$, $f_i(b \otimes c) = \begin{cases} f_i(x) \otimes y & \text{if } \varphi_i(c) \leq \varepsilon_i(b) \\ x \otimes f_i(y) & \text{if } \varphi_i(c) > \varepsilon_i(b) \end{cases}$, $e_i(b \otimes c) = \begin{cases} e_i(x) \otimes y & \text{if } \varphi_i(c) < \varepsilon_i(b) \\ x \otimes e_i(y) & \text{if } \varphi_i(c) \geq \varepsilon_i(b) \end{cases}$. If \mathfrak{B} and \mathfrak{C} are finite, $\varphi_i(b \otimes c) = \varphi_i(x) + \max(0, \varphi_i(c) - \varepsilon_i(x))$ and $\varepsilon_i(b \otimes c) = \varepsilon_i(x) + \max(0, \varepsilon_i(x) - \varphi_i(c))$.

In type C_n , we consider $\{\mathbf{e}_i\}_{i=1}^n$ the canonical basis of \mathbb{R}^n . The root system is $\Phi_C = \{\pm \mathbf{e}_i \pm \mathbf{e}_j \mid i < j\} \cup \{\pm 2\mathbf{e}_i\}$ and the simple roots are $\alpha_i = \mathbf{e}_i - \mathbf{e}_{i+1}$, if $i \in [n-1]$, $\alpha_n = 2\mathbf{e}_n$. The weight lattice \mathbb{Z}^n has dominant weights $\lambda = (\lambda_1 \geq \dots \geq \lambda_n \geq 0)$. In type C_n , the standard crystal is seminormal

and has the following crystal graph: $1 \xrightarrow{1} 2 \xrightarrow{2} \dots \xrightarrow{n-1} n \xrightarrow{n} \bar{n} \xrightarrow{n-1} \dots \xrightarrow{1} \bar{1}$ with set $\mathfrak{B} = [\pm n]$, $wt(\boxed{i}) = \mathbf{e}_i$, $wt(\overline{\boxed{i}}) = -\mathbf{e}_i$. The highest weight element is the word 1, and the highest weight \mathbf{e}_1 . We denote the crystal by $\mathfrak{B}^{\mathbf{e}_1}$.

For later convenience, (this becomes clear in the next section) the crystal operators will act, from now on, on the right.

The crystal $\mathfrak{B}^{\mathbf{e}_1}$ is the crystal on the words of $[\pm n]^*$ of a sole letter. The tensor product of crystals allows us to define the crystal $G_n = \bigoplus_{k \geq 0} (\mathfrak{B}^{\mathbf{e}_1})^{\otimes k}$ of all words in $[\pm n]^*$, where the vertex $w_1 \otimes \dots \otimes w_k$ is identified with the word $w_1 \dots w_k \in [\pm n]^*$. The action of the operators e_i and f_i is easily given by the signature rule [13, 18, 5]. We substitute each letter w_j by $+$ if $w_j \in \{i, \overline{i+1}\}$ or by $-$ if $w_j \in \{i+1, \overline{i}\}$, and erase it in any other case. Then successively erase any pair $+-$ until all the remaining letters form a word that looks like $-^a +^b$. Then $\varphi_i(w) = b$ and $\varepsilon_i(w) = a$, e_i acts on the letter associated to the rightmost unbracketed $-$ (i.e., not erased), whereas f_i acts on the letter w_j as-

sociated to the leftmost unbracketed $+$, $(w_j)f_i = \begin{cases} i+1 & \text{if } w_j = i \wedge i \neq n \\ \overline{i} & \text{if } w_j = \overline{i+1} \\ \bar{n} & \text{if } w_j = i \wedge i = n \end{cases}$,

and the other letters of w are unchanged, and e_i is the inverse map. If $b = 0$ then $(w)f_i = 0$ and if $a = 0$ then $(w)e_i = 0$.

The crystal G_n , as a graph, is the union of connected components where each component has a unique highest weight word. Two connected components are isomorphic if and only if they have the same highest weight [12]. Two words in $[\pm n]^*$ are said to be crystal connected or coplactic equivalent if and only if they belong to the same connected component of G_n . This means that both words are obtained from the same highest weight word, through a sequence of crystal operators f_i , or one is obtained from another by some sequence of crystal operators f_i and e_j , $i, j \in [n]$. The connected components of G_n are the coplactic classes in the Robinson-Schensted correspondence that identify words with the same oscillating tableau [18, Proposition 5.2.1]. Also, two words $w_1, w_2 \in [\pm n]^*$ are Knuth equivalent if and only if they occur in the same place in two isomorphic connected components of G_n , that is, they are obtained from two highest words with the same weight through a same sequence of crystal operators [18]. Crystal operators are coplactic and commute with the *jeu de taquin*. The next proposition identifies all highest weight words of G_n .

Proposition 9. *Let w be a word in the alphabet $[\pm n]$. Then w is a highest weight word if and only if the weight of all its prefixes (including itself) is a partition. In this case, one has that $P(w) = K(\lambda)$ the Yamanouchi tableau of shape λ , the weight of w*

Proof: Part "if": We will prove the contrapositive of the statement. There is a i such that $(w)e_i \neq 0$. Let k be the position of the leftmost $-$ of the signature rule of w , and consider the prefix w_k with the first k letters. Since the k -th letter of w had an unbracketed $-$ in the signature rule then the last letter of w_k will also be an unbracketed $-$. Hence there are more $-$ than $+$ in the signature rule of w_k . Call t_α to the number of α in w_k . We have that $t_i + t_{\overline{i+1}} < t_{i+1} + t_{\overline{i}} \Leftrightarrow t_i - t_{\overline{i}} < t_{i+1} - t_{\overline{i+1}}$, hence the weight of w_k is not a partition.

Part "only if": We will once again prove the contrapositive of the statement. Let w_k be a prefix such that its weight is not a partition. Hence there is $i \in [n]$ such that $t_i - t_{\overline{i}} < t_{i+1} - t_{\overline{i+1}} \Leftrightarrow t_i + t_{\overline{i+1}} < t_{i+1} + t_{\overline{i}}$, hence for this i there will be more $-$ than $+$ in the signature rule of w_k . So in the first k letters of w there will be more $-$ than $+$, so there is an unbracketed $-$ in w , hence $(w)e_i \neq 0$. Note that the argument works even if $i = n$. In this case we need to assume $t_{n+1} = t_{\overline{n+1}} = 0$.

It follows from [18, Proposition 3.2.6] that the insertion of the highest word w of weight λ is $K(\lambda)$. ■

Choose a word $w \in [\pm n]^*$ such that the shape of $P(w)$ is λ . If we replace every word of its coplactic class with its insertion tableau we obtain the crystal of tableaux \mathfrak{B}^λ that has all KN tableaux of shape λ on the alphabet $[\pm n]$. The crystal \mathfrak{B}^λ does not depend on the initial choice of word w , as long as $P(w)$ has shape λ . [18, Theorem 6.3.8].

4. Right and Left Keys and Demazure atoms in type C

In this section, we define type C frank words on the alphabet $[\pm n]$ and use them to create the right and left key maps, that send KN tableaux to key tableaux in type C. The main result of this section is the type C version [17, Theorem 3.8], due to Lascoux and Schützenberger, which, using the right key map, gives a combinatorial description of Demazure atoms in type C.

4.1. Frank words in type C. Frank words were introduced in type A by Lascoux and Schützenberger in [17]. We start by defining frank words in the alphabet $[\pm n]$.

Definition 9. Let w be word on the alphabet $[\pm n]$. We say that w is a type C frank word if the length of its columns form a multiset equal to the multiset formed by the length of the columns of the tableau $P(w)$.

Example 13. In Example 11 we have that $P(23\bar{2}\bar{3}1) = P(\bar{1}113\bar{3}) = \begin{array}{|c|c|c|} \hline 1 & 1 & \bar{1} \\ \hline 3 & & \\ \hline 3 & & \\ \hline \end{array}$.

Since $23\bar{2}\bar{3}1$ and $\bar{1}113\bar{3}$ have one column of length 3 and two columns of length 1, they are frank words.

Given a frank word w , the number of letters of w is the same as the number of cells of $P(w)$, hence the case 3 of the Baker-Lecouvey insertion doesn't happen.

Proposition 10. Let w be frank word on the alphabet $[\pm n]$. All columns of w are admissible.

Proof: Suppose that the statement is false. So there is a factor of w that is a non-admissible column with all of its proper factors admissible. Hence we can apply the Knuth relation $K5$, meaning that w is Knuth related to a smaller word w' . But in this case, the number of letters of w' is less than the number of cells of $P(w) = P(w')$, which is a contradiction. ■

The following proposition is an extension of [8, Proposition 7] on SSYT to KN tableaux.

Proposition 11. Let T be a KN tableau of shape λ . Let μ/ν be a skew diagram with same number of columns of each length as T . Then there is a unique KN skew tableau S with shape μ/ν that rectifies to T and $cr(S)$ is a frank word.

Proof: If T is a Yamanouchi tableau $K(\lambda)$ and $S \in \mathcal{KN}(\mu/\nu, n)$ rectifies to $K(\lambda)$, then, since S and $K(\lambda)$ have the same number of cells, all entries of S are unbarred, hence S is a semistandard skew tableau. So, it follows from [8, Proposition 7] that S exists and is unique. If T is not a Yamanouchi tableau, note that T is crystal connected to $K(\lambda)$ and from [18, Theorem 6.3.8] we have that the symplectic *jeu de taquin* slides commutes with the action of the crystal operators. Consider Y'_λ the only tableau on the skew-shape μ/ν that rectifies to Y_λ , which exists due to [8, Proposition 7]. Since S rectifies to T , which is crystal connected to $K(\lambda)$, and Y'_λ rectifies to $K(\lambda)$, S is crystal connected to Y'_λ and the path has same sequence of colours as the one from T to $K(\lambda)$. Hence S exists and is uniquely defined. ■

Corollary 12. *Let S be as in the previous proposition. The last column of S depends only on the length of that column.*

Proof: All other skew tableaux with given last column length can be found from a given one by playing the symplectic *jeu de taquin* or its reverse in all columns except the last one. Note that S has the same number of cells of the tableau obtained after rectifying, hence we can't lose cells when applying the symplectic *jeu de taquin* or its reverse. ■

Fixed a KN tableau T , consider the set of all possible last columns taken from skew-tableaux with same number of columns of each length as T . Corollary 12 implies that this set has one element for each distinct column length of T . For every column C in this set, consider the columns rC , its right column. The next proposition implies that this set of right columns is nested, if we see each column as the set formed by its elements.

Proposition 13. *Consider T a two-column KN skew tableau C_1C_2 with an empty cell in the first column. Slide that cell once via symplectic *jeu de taquin*, obtaining a two-column KN skew tableau $C'_1C'_2$ with an empty cell. Then $rC'_2 \subseteq rC_2$.*

Proof: If the sliding was vertical then $C'_2 = C_2$, hence $rC'_2 = rC_2$. If the sliding was horizontal, call β to the number on the cell right of the empty cell on $spl(T)$. Call Φ to the function that takes an admissible column to the associated coadmissible column.

If $\beta = b$ is unbarred then $C'_2 = \Phi^{-1}(\Phi(C_2) \setminus \{b\})$. In this case $\Phi(C'_2) = \Phi(C_2) \setminus \{b\}$, hence rC_2 and rC'_2 have the same barred part. Consider $z_1 < \dots < z_\ell$ the unbarred letters that appear on C_2 and not on $\Phi(C_2)$. When we take b from $\Phi(C_2)$, if $\bar{b} \in \Phi(C_2)$ our set of letters $z_1 < \dots < z_\ell$ will lose an element, giving the inclusion of the unbarred part of C'_2 in C_2 ; if $\bar{b} \notin \Phi(C_2)$, then $b \in C_2$ and in C'_2 the least $z_i > b$ may reduce to b , and subsequent z_j may reduce to z_{j-1} . Hence we have the inclusion of the unbarred part of C'_2 in C_2 .

If $\beta = \bar{b}$ is barred then $C'_2 = C_2 \setminus \{\bar{b}\}$. In this case rC_2 and rC'_2 have the same unbarred part. Consider $\bar{t}_1 > \dots > \bar{t}_\ell$ the barred letters that appear on $\Phi(C_2)$ and not on C_2 . When we take \bar{b} from C_2 , if $b \in C_2$ our set of $\bar{t}_1 > \dots > \bar{t}_\ell$ letters will lose an element, giving the inclusion of the barred part of rC'_2 in rC_2 ; if $b \notin C_2$, then $\bar{b} \in \Phi(C_2)$ and in C'_2 the least $\bar{z}_i > \bar{b}$ may reduce to \bar{b} , and subsequent bigger \bar{z}_j 's may reduce to \bar{z}_{j+1} . Hence we have the inclusion of the barred part of $\Phi(C'_2)$ in $\Phi(C_2)$. ■

This proposition defines a map that sends a KN tableau to key tableau in type C , identified as the (symplectic) right key of a given KN tableau.

Theorem 14 (Right key map). *Given a KN tableau T , we can replace each column with a column of the same size taken from the right columns of the last columns of all skew tableaux associated to it. We call this tableau the right key tableau of T and denote it by $K_+(T)$.*


Proof: The previous proposition implies that the columns of $K_+(T)$ are nested and do not have symmetric entries. So, it is indeed a KN key tableau. ■

Remark. *Recall the set up of Proposition 11. If the shape of S , μ/ν , is such that every two consecutive columns have at least one cell in the same row, then each column of S is a column of the word $cr(S)$, hence $cr(S)$ is a frank word. Moreover, the columns of S appear in reverse order in $cr(S)$. Therefore, given a KN tableau T , the columns of $K_+(T)$ can be also found as the right columns of the first columns of frank words associated to T .*

If T is a SSYT then this right key map coincides with the one defined by Lascoux and Schützenberger in [17].

Example 14. *The tableau $T = \begin{array}{|c|c|c|} \hline 1 & 3 & \bar{1} \\ \hline 3 & 3 & \\ \hline 3 & & \\ \hline \end{array}$ gives rise to six KN skew tableaux*

with same number of columns of each length as T , each one corresponding to a permutation of its column lengths, and each one is associated to its column reading, which is a frank word.



The right key tableau associated to T has as columns $r \begin{array}{|c|} \hline 3 \\ \hline 3 \\ \hline 1 \\ \hline \end{array}$, $r \begin{array}{|c|} \hline 3 \\ \hline 1 \\ \hline \end{array}$ and $r \begin{array}{|c|} \hline 1 \\ \hline \end{array}$.

Hence $K_+(T) = \begin{array}{|c|c|c|} \hline 3 & 3 & \bar{1} \\ \hline 2 & 1 & \\ \hline 1 & & \\ \hline \end{array}$.

In the same spirit of the right key, we define the left key of a KN tableau. Just like in Proposition 13, we can prove that the slides of the symplectic *jeu de taquin* are effectively adding an entry to ℓC_1 , i.e. $\ell C_1 \subseteq \ell C'_1$, hence the left columns of the first columns of all skew tableaux with the same number of columns of each length as T will be nested.

So, if we replace each column of T with a column of the same size taken from the left columns of the first columns of all skew tableaux associated to it we obtain the left key $K_-(T)$.

Example 15. In Example 14 we have that the left key of $T = \begin{array}{|c|c|c|} \hline 1 & 3 & \bar{1} \\ \hline 3 & 3 & \\ \hline 3 & & \\ \hline \end{array}$ has

as columns

$$\ell \begin{array}{|c|} \hline 1 \\ \hline 3 \\ \hline 3 \\ \hline \end{array}, \ell \begin{array}{|c|} \hline 1 \\ \hline 2 \\ \hline \end{array} \text{ and } \ell \begin{array}{|c|} \hline 2 \\ \hline \end{array}. \text{ Hence } K_-(T) = \begin{array}{|c|c|c|} \hline 1 & 1 & 2 \\ \hline 2 & 2 & \\ \hline 3 & & \\ \hline \end{array}.$$

4.2. Demazure crystals and right key tableaux. Let $\lambda \in \mathbb{Z}^n$ be a partition and $v \in \lambda B_n$. We define $\mathfrak{U}(v) = \{T \in \mathcal{KN}(\lambda, n) \mid K_+(T) = K(v)\}$ the set of KN tableaux of B^λ with right key $K(v)$.

Given a subset X of \mathfrak{B}^λ , consider the operator \mathfrak{D}_i on X , with $i \in [n]$ defined by $X\mathfrak{D}_i = \{x \in \mathfrak{B}^\lambda \mid (x)e_i^k \in X \text{ for some } k \geq 0\}$ [5]. If $v = \lambda\sigma$ where $\sigma = s_{i_1} \dots s_{i_{\ell(\sigma)}} \in B_n$ is a reduced word, we define the Demazure crystal to be

$$\mathfrak{B}_v = \{K(\lambda)\}\mathfrak{D}_{i_1} \dots \mathfrak{D}_{i_{\ell(\sigma)}}. \quad (1)$$

This definition is independent of the reduced word for σ [5, Theorem 13.5]. In particular, when σ is the longest element of B_n we recover \mathfrak{B}^λ . Also this definition is independent of the coset representative of $W_\lambda\sigma$, that is, $\mathfrak{B}_{\lambda\sigma} = \mathfrak{B}_{\lambda\sigma_v}$. From [4, Proposition 2.4.4], σ uniquely factorizes as $\sigma'\sigma_v$ where $\sigma' \in W_\lambda$ and $\ell(\sigma) = \ell(\sigma') + \ell(\sigma_v)$. From the signature rule, Subsection 3.4, if $\sigma' = s_{j_1} \dots s_{j_{\ell(\sigma')}} \in W_\lambda$ is a reduced word, $\mathfrak{B}_{\lambda\sigma'} = \mathfrak{B}_\lambda = \{K(\lambda)\}\mathfrak{D}_{i_1} \dots \mathfrak{D}_{i_{\ell(\sigma')}} = \{K(\lambda)\}$ and we may write in (1) $\mathfrak{B}_{\lambda\sigma} = \mathfrak{B}_v$.

From [4, Proposition 2.5.1], if $\rho \leq \sigma$ in B_n then $\rho_u \leq \sigma_v$ where $u = \lambda\rho$. Since $(x)e_i^0 = x$, if $\rho \leq \sigma$ then $\mathfrak{B}_{\lambda\rho} = \mathfrak{B}_{\lambda\rho_u} \subseteq \mathfrak{B}_{\lambda\sigma_v} = \mathfrak{B}_v$. Thus we define the Demazure atom crystal $\hat{\mathfrak{B}}_v$ to be

$$\hat{\mathfrak{B}}_v = \hat{\mathfrak{B}}_{\lambda\sigma} := \mathfrak{B}_{\lambda\sigma_v} \setminus \bigcup_{\rho_u < \sigma_v} \mathfrak{B}_{\lambda\rho_u} = \mathfrak{B}_v \setminus \bigcup_{u < v} \mathfrak{B}_u = \mathfrak{B}_v \setminus \bigcup_{K(u) < K(v)} \mathfrak{B}_u, \quad (2)$$

where the two rightmost identities follow from Theorem 7.

Lemma 1. *Let $\sigma = s_i$ be a generator of B_n and C an admissible column such that $(C)f_i \neq 0$. Then $wt(rC) = wt(r((C)f_i))$ or $wt(rC) = wt(r((C)f_i))\sigma$.*

Proof: Let $i = n$. We can apply f_i to C if and only if $n \in C$ and $\bar{n} \notin C$. In this case $n \in rC$ and after applying f_i we have $n \notin C$ and $\bar{n} \in C$, hence $\bar{n} \in rC$. So $wt(rC) = wt(r((C)f_n))s_n$.

Let $i < n$. We can apply f_i to C , so we have 6 cases to study:

- (1) $i \in C, i+1, \overline{i+1}, \bar{i} \notin C$: In this case we have that $i+1 \in (C)f_i, i, \overline{i+1}, \bar{i} \notin (C)f_i$. Note that $\bar{i} \notin rC$ and $\overline{i+1} \notin r((C)f_i)$. If $\overline{i+1} \notin rC$ then $\bar{i} \notin r((C)f_i)$, hence f_i swaps the weight of i and $i+1$ from $(1, 0)$ to $(0, 1)$, respectively. If $\overline{i+1} \in rC$ then $\bar{i} \in r((C)f_i)$, hence f_i swaps the weight of i and $i+1$ from $(1, -1)$ to $(-1, 1)$.
- (2) $i, \overline{i+1} \in C, i+1, \bar{i} \notin C$: In this case we have that $i+1, \overline{i+1} \in (C)f_i, i, \bar{i} \notin (C)f_i$. Note that $i, \overline{i+1} \in rC, i+1, \bar{i} \notin rC$ and that $i+1, \bar{i} \in r((C)f_i), i, \overline{i+1} \notin r((C)f_i)$, and all other appearances in rC are intact. Hence f_i swaps the weight of i and $i+1$ from $(1, -1)$ to $(-1, 1)$.
- (3) $i+1, \overline{i+1} \in C, i, \bar{i} \notin C$: In this case we have that $i+1, \bar{i} \in (C)f_i, i, \overline{i+1} \notin (C)f_i$. Note that $i+1, \bar{i} \in rC, i, \overline{i+1} \notin rC$ and that $i+1, \bar{i} \in r((C)f_i), i, \overline{i+1} \notin r((C)f_i)$, and all other appearances in rC are intact. Hence f_i did nothing to weight of rC .
- (4) $i, i+1, \overline{i+1} \in C, \bar{i} \notin C$: In this case we have that $i, i+1, \bar{i} \in (C)f_i, \overline{i+1} \notin (C)f_i$. Note that $i, i+1 \in rC, \overline{i+1}, \bar{i} \notin rC$ and that $i, i+1 \in r((C)f_i), \overline{i+1}, \bar{i} \notin r((C)f_i)$, and all other appearances in rC are intact. Hence f_i did nothing to weight of rC .
- (5) $i, \overline{i+1}, \bar{i} \in C, i+1 \notin C$: In this case we have that $i+1, \overline{i+1}, \bar{i} \in (C)f_i, i \notin (C)f_i$. Note that $i, \overline{i+1} \in rC, i+1, \bar{i} \notin rC$ and that $i+1, \bar{i} \in r((C)f_i), i, \overline{i+1} \notin r((C)f_i)$, and all other appearances in rC are intact. Hence f_i swaps the weight of i and $i+1$ from $(1, -1)$ to $(-1, 1)$.
- (6) $\overline{i+1} \in C, i, i+1, \bar{i} \notin C$: In this case we have that $\bar{i} \in (C)f_i, i, i+1, \overline{i+1} \notin (C)f_i$. Note that $i, i+1 \notin rC$ and $\overline{i+1} \in rC$. If $\bar{i} \in rC$ then we have $i, i+1 \notin r((C)f_i)$ and $\overline{i+1}, \bar{i} \in r((C)f_i)$, so f_i did nothing to weight of rC . If $\bar{i} \notin rC$ then $\overline{i+1} \notin r((C)f_i)$ and $\bar{i} \in r((C)f_i)$, hence f_i swaps the weight of i and $i+1$ from $(0, -1)$ to $(-1, 0)$.

■

Remark. *All the cases where the weight is preserved happen to have equal weight for i or $i + 1$ in rC or we are in a column C in which we can also apply e_i .*

Hence we have the following corollaries:

Corollary 15. *Let T be a KN tableau and $i \in [n]$. If $K_+(T) = K(v)$, for some $v = (v_1, \dots, v_n) \in \mathbb{Z}^n$, then $K_+((T)f_i) = K(v)$ or $K_+((T)f_i) = K(vs_i)$. Moreover, $K_+((T)f_i) = K(vs_i)$ only if $v_i > v_{i+1}$ (in the usual ordering of real numbers) and $1 \leq i < n$, or, $v_i > 0$ and $i = n$.*

Proof: Consider a multiset of frank words F such that the multiset of length of their first columns is the same of the multiset of lengths of columns of T .

If $K_+((T)f_i) = K_+(T)$ then we are done. Else there are two cases: $1 \leq i < n$ and $i = n$.

Consider $1 \leq i < n$. There is a column of T whose weight of i is bigger than its weight for $i + 1$. Since T is a key tableau, this implies that in all columns of T weight of i is bigger or equal than the weight of $i + 1$.

Let A be the subset of F such that the weight of i and $i + 1$ in the right column of its first column is different and does not swap when we apply f_i to the frank word.

Consider (a, b) the sum of weights of i and $i + 1$, respectively, of all right columns of first columns of words in A , and (c, d) defined analogously to $F \setminus A$.

The weights of i and $i + 1$ in $K_+(T)$ is $(a, b) + (c, d) = (a + c, b + d)$ and the weights of i and $i + 1$ in $K_+((T)f_i)$ is $(a, b) + (d, c) = (a + d, b + c)$, and note that $(a + c, b + d) \in (a + d, b + c)B_2$, because f_i doesn't change any other weight (Lemma 1).

Since in all columns of T weight of i is bigger or equal than the weight of $i + 1$, $a \geq 0$ and $b \leq 0$, and they are equal when $A = \emptyset$, so $(a + c, b + d) \in (a + d, b + c)s_1$, hence $\text{wt}(K_+((T)f_i)) = vs_i$. Hence we assume $a \neq b$. If $c = d$ we have $\text{wt}(K_+((T)f_i)) = v$, hence $K_+((T)f_i) = K(v) = K_+(T)$, which is a contradiction.

This implies that $(a + c, b + d) = (a + d, b + c)\sigma$ where $\sigma = \overline{12}$ or $\sigma = \overline{21}$. The first case implies that $a = \frac{-c-d}{2} = b$ and the second case implies $c = \frac{-a-b}{2} = d$, hence there are not more possibilities for the weight of $K_+((T)f_i)$.

The case $i = n$ is a simpler version of this one. ■

Corollary 16. *Let $\sigma = s_i$ be a generator of B_n and C an admissible column. Then $\text{wt}(rC) = \text{wt}(r((C)e_i))$ or $\text{wt}(rC) = \text{wt}(r((C)e_i))\sigma$.*

Proof: Call C' to $(C)e_i$. By Lemma 1 we have that $wt(rC') = wt(r((C')f_i))$ or $wt(C') = wt(r((C')f_i))\sigma$, so we have that $wt(r((C)e_i)) = wt(rC)$ or $wt((C)e_i) = wt(rC)\sigma \Leftrightarrow wt((C)e_i)\sigma = wt(rC)$. ■

Lemma 2. *Let $i \in [n]$ and C be an admissible column such that one of the following happens*

- (1) $i < n$ and the weight of i in rC is less than the weight of $i + 1$ in rC ;
- (2) $i = n$ and weight of i is negative in rC ,

then we can apply e_i to C (in the sense $(C)e_i \neq 0$).

Proof: If $i = n$ then $-n$ appears on rC and n does not. Since n is the biggest unbarred letter of the alphabet we have that $-n$ also appears in C and n does not. Hence we can apply e_n to C .

If $i < n$ and the weight of i in rC is less than the weight of $i + 1$ in rC then the weight of both can be one of the following three options: $(0, 1)$, $(-1, 1)$, $(-1, 0)$. Note that rC does not have symmetric entries. So in the first two cases we have that $i + 1$ exists in rC and i does not, hence $i + 1$ exists in C and i does not, so we can apply e_i to C . In the last case, we have that \bar{i} exists in rC and $i + 1$ and $\overline{i + 1}$ does not. Hence we have that \bar{i} exists in C and i or $\overline{i + 1}$ does not, so we can apply e_i to C . ■

The next theorem is the main theorem of this paper. It gives a description of a Demazure crystal atom in type C using the right key map Theorem 14. Lascoux and Schützenberger, in [17, Theorem 3.8], proved the type A version of this theorem, which consists in considering the case when $v \in \mathbb{N}^n$ and, consequently, $\sigma_v \in \mathfrak{S}_n$. For inductive reasoning, used in what follows, we recall the chain property on the set of minimal length coset representatives modulo W_λ [4, Theorem 2.5.5].

Theorem 17. *Let $v \in \lambda B_n$. Then $\mathfrak{U}(v) = \hat{\mathfrak{B}}_v$.*

Proof: Let ρ be a minimal length coset representative modulo W_λ such that $v = \lambda\rho$. We will proceed by induction on $\ell(\rho)$. If $\ell(\rho) = 0$ then $\rho = id$ and $v = \lambda$. In this case we have that $\hat{\mathfrak{B}}_\lambda = \{K(\lambda)\} = \mathfrak{U}(\lambda)$.

Let $\rho \geq 0$. Consider $\sigma = s_i$ a generator of B_n such that $\rho\sigma > \rho$ and $\lambda\rho\sigma \neq \lambda\rho = v$, i.e., $\rho\sigma\rho^{-1} \notin W_\lambda$. Recall e_i , ε_i , f_i and ϕ_i from the definition of the crystal \mathfrak{B}^λ . If $T \in \mathfrak{B}_{\lambda\rho\sigma}$ then T is obtained after applying f_i (maybe more than once) to a tableau in $\hat{\mathfrak{B}}_{\lambda\rho}$, which by inductive hypothesis exists in

$\mathfrak{U}(v)$. By Corollary 15, if $(T)f_i \notin \mathfrak{U}(v)$ then $(T)f_i \in \mathfrak{U}(v\sigma)$. So it is enough to prove that given a tableau $T \in \mathfrak{U}(v) \cup \mathfrak{U}(v\sigma)$ then $(T)e_i^{\varepsilon_i(T)} \in \mathfrak{U}(v)$.

We have two different cases to consider: $i = n$ and $i < n$.

If $T \in \mathfrak{U}(v\sigma)$ then, if $i < n$, there exists a frank word of T such that, if V_1 is its first column then rV_1 has less weight for i than for $i + 1$ (less in the usual ordering of real numbers); if $i = n$, there exists a frank word of T such that, if V_1 is its first column then rV_1 has negative weight for i . Since we are in the column rV_1 , if $i < n$, i and $i + 1$ can have weights $(0, 1)$, $(-1, 1)$ or $(-1, 0)$ and if $i = n$ then i has weight -1 . Note that these are the exact conditions of Lemma 2. In either case, due to Lemma 2, we can applying e_i enough times to the frank word associated until this no longer happens. This is true because we only need to look to V_1 to see if it changes after applying e_i enough times to the frank word. In the signature rule we have that successive applications of e_i changes the letters of a word from the end to the beginning, so, from the remark after Lemma 1, the number of times that we need to apply e_i , in order to conditions of Lemma 2 do not hold for the first column, is $\varepsilon_i(T)$. So $K_+ \left((T)e_i^{\varepsilon_i(T)} \right) \neq K(v\sigma)$, hence, from Corollary 16, we have that $(T)e_i^{\varepsilon_i(T)} \in \mathfrak{U}(v)$.

If $T \in \mathfrak{U}(v)$ then $(T)e_i^{\varepsilon_i(T)} \in \mathfrak{U}(v)$ because if not, $(T)e_i^{\varepsilon_i(T)}$ will be in a Demazure crystal associated to $\rho' \in B_n$, with $\rho' < \rho$ such that $\rho'\sigma = \rho$. This cannot happen because in this case $\rho' = \rho\sigma < \rho$, which is a contradiction. ■

4.3. Combinatorial description of type C Demazure characters and atoms.

Given $v \in \lambda B_n$ define the Demazure character (or key polynomial), κ_v , and the Demazure atom in type C , $\hat{\kappa}_v$, as the generating functions of the KN tableaux weights in \mathfrak{B}_v and $\hat{\mathfrak{B}}_v$, respectively: $\kappa_v = \sum_{T \in \mathfrak{B}_{\lambda\sigma_v}} x^{\text{wt} T}$, $\hat{\kappa}_v =$

$\sum_{T \in \hat{\mathfrak{B}}_{\lambda\sigma_v}} x^{\text{wt} T}$. Theorem 17 detects the KN tableaux in \mathfrak{B}^λ contributing to the

Demazure atom $\hat{\kappa}_v$, $\hat{\kappa}_v = \sum_{\substack{K_+(T)=K(v) \\ T \in \mathfrak{B}^\lambda}} x^{\text{wt} T}$.

Proposition 18. *Given $v \in \lambda B_n$, one has $\kappa_v = \sum_{u \leq v} \hat{\kappa}_u$.*

Proof: It is enough to prove that $\mathfrak{B}_v = \bigcup_{u \leq v} \hat{\mathfrak{B}}_u$, because κ_v and $\hat{\kappa}_u$ are the generating functions of the tableaux weights in \mathfrak{B}_v and $\hat{\mathfrak{B}}_u$, respectively. Since $v = \lambda\sigma$, where $\sigma := \sigma_v$, we can rewrite the identity as $\mathfrak{B}_{\lambda\sigma} = \bigcup_{\rho \leq \sigma} \hat{\mathfrak{B}}_{\lambda\rho}$.

We will proceed by induction on $\ell(\sigma)$. If $\ell(\sigma) = 0$ then the result follows because $\mathfrak{B}_\lambda = \hat{\mathfrak{B}}_\lambda = \{K(\lambda)\}$. From 2, $\hat{\mathfrak{B}}_{\lambda\sigma} = \mathfrak{B}_{\lambda\sigma} \setminus \bigcup_{\rho < \sigma} \mathfrak{B}_{\lambda\rho}$, and by inductive hypothesis, we have that $\mathfrak{B}_{\lambda\rho} = \bigcup_{\rho' \leq \rho} \hat{\mathfrak{B}}_{\lambda\rho'}$. Hence:


$$\hat{\mathfrak{B}}_{\lambda\sigma} = \mathfrak{B}_{\lambda\sigma} \setminus \bigcup_{\rho < \sigma} \mathfrak{B}_{\lambda\rho} = \mathfrak{B}_{\lambda\sigma} \setminus \bigcup_{\rho < \sigma} \bigcup_{\rho' \leq \rho} \hat{\mathfrak{B}}_{\lambda\rho'} = \mathfrak{B}_{\lambda\sigma} \setminus \bigcup_{\rho' < \sigma} \hat{\mathfrak{B}}_{\lambda\rho'}$$

■

Proposition 18, the equivalence $u \leq v \Leftrightarrow K(u) \leq K(v)$, and Theorem 17, allow to detect the KN tableaux contributing to a key polynomial in type C :

$$\kappa_v = \sum_{u \leq v} \hat{\kappa}_u = \sum_{\substack{u \leq v \\ T \in \mathfrak{U}(u)}} x^{wtT} = \sum_{\substack{K(u) \leq K(v) \\ T \in \mathfrak{U}(u)}} x^{wtT} = \sum_{K(T) \leq K(v)} x^{wtT}.$$

Example 16. We start by looking to the crystal graph associated to the partition $\lambda = (2, 1)$:



The crystal is split into several parts. Each one of those parts is a Demazure atom and contains exactly one symplectic key tableau, so we can identify each part with the weight of that key tableau, which is a vector in the B_2 -orbit of $(2, 1)$. From the previous theorem we have that all tableaux in the same part have the same right key.

One can check that $\mathfrak{U}((1, \bar{2})) = \left\{ \begin{smallmatrix} 1 & \bar{2} \\ 2 \end{smallmatrix}, \begin{smallmatrix} 1 & \bar{2} \\ 2 \end{smallmatrix} \right\} = \hat{\mathfrak{B}}_{\lambda_{s_1 s_2}}$, for example.

Also,
 $\mathfrak{B}_{(1, \bar{2})} = \{T \in \mathfrak{B}^\lambda \mid K_+(T) \leq K((1, \bar{2}))\} = \left\{ \begin{smallmatrix} 1 & 1 \\ 2 \end{smallmatrix}, \begin{smallmatrix} 1 & 2 \\ 2 \end{smallmatrix}, \begin{smallmatrix} 1 & 1 \\ 2 \end{smallmatrix}, \begin{smallmatrix} 1 & \bar{2} \\ 2 \end{smallmatrix}, \begin{smallmatrix} 1 & \bar{2} \\ 2 \end{smallmatrix} \right\}.$

5. Realization of the Lusztig involution in types A and C

Let \mathfrak{B}^λ be the crystal with set $\mathcal{KN}(\lambda, n)$ (respectively $SSYT(\lambda, n)$).

Definition 10. The Lusztig involution $L : \mathfrak{B}^\lambda \rightarrow \mathfrak{B}^\lambda$ is the only involution such that for all $i \in I$ ($I = [n - 1]$ in type A_{n-1} and $I = [n]$ in type C_n):

- (1) $wt(L(x)) = (wt(x))\omega_0$, where ω_0 is the longest element of the Weyl group;
- (2) $(Lx)e_i = L((x)f_{i'})$ and $(Lx)f_i = L((x)e_{i'})$ where i' is such that $(\alpha_i)\omega_0 = -\alpha_{i'}$;
- (3) $\varepsilon_i(Lx) = \varphi_{i'}(x)$ and $\varphi_i(Lx) = \varepsilon_{i'}(x)$.

For type A we have that ω_0 is the reverse permutation and $i' = n - i$, and for type C_n we have $\omega_0 = -\text{Id}$ and $i' = i$, where Id is the identity map. In type C_n the involution can be seen as flipping the crystal upside down.

Definition 11. [5] Let \mathfrak{C} be a connected component in the type C_n crystal G_n . The dual crystal \mathfrak{C}^\vee is the crystal obtained from \mathfrak{C} after reversing the direction of all arrows. Also, if $x \in \mathfrak{C}$, then for its correspondent in \mathfrak{C}^\vee , x^\vee , we have $wt(x) = -wt(x^\vee)$.

In type C , since $i' = i$ and $\omega_0 = -\text{Id}$, it follows from the definition that \mathfrak{C} and \mathfrak{C}^\vee , as crystals in G_n , have the same highest weight. Therefore, they are isomorphic. In the case of \mathfrak{B}^λ , with set $\mathcal{KN}(\lambda, n)$, the Lusztig involution is a realization of the dual crystal. Hence the crystal \mathfrak{B}^λ with set $\mathcal{KN}(\lambda, n)$ is self-dual. We shall see other realizations of the dual.

5.1. Evacuation algorithms. In type A_{n-1} , the Lusztig involution on the crystal with set $SSYT(\lambda, n)$ is known as Schützenberger involution or evacuation, and takes $T \in SSYT(\lambda, n)$ to $T^{\text{Ev}} \in SSYT(\lambda, n)$, whose weight is $(wt T)\omega_0$, where ω_0 is the longest permutation of \mathfrak{S}_n , in the Bruhat order. Note that $(wt T)\omega_0$ is the vector $wt T$ in reverse order, i.e., $(v_1, \dots, v_n)\omega_0 = (v_n, \dots, v_1)$. In type C_n we will work with KN tableaux instead of SSYTs. Consider $T \in \mathcal{KN}(\lambda, n)$. In this case, $T^{\text{Ev}} \in \mathcal{KN}(\lambda, n)$ and $wt T = -wt T^{\text{Ev}} = (wt T^{\text{Ev}})\omega_0^C$, where ω_0^C is the longest permutation of B_n . The complement of a tableau or a word in types A_{n-1} or C_n consists in applying ω_0 or ω_0^C , respectively, to all its entries. In type A_{n-1} , it sends i to $n + 1 - i$ for all $i \in [n]$, i.e., $(i)\omega_0 = n + 1 - i$ and in type C_n we have $(i)\omega_0 = -i$. Given a SSYT, there are several algorithms, due to Schützenberger, to obtain a SSYT

with the same shape whose weight is its reverse. We recall some versions of them for which one is able to find analogues for KN tableaux.

Algorithm 1.

- (1) Define $w = cr(T)$.
- (2) Define w^* the word obtained by complementing its letters and writing it backwards.
- (3) $T^{Ev} := P(w^*)$.

Example 17. In type A , the tableau $T = \begin{array}{|c|c|c|c|} \hline 1 & 1 & 2 & 3 \\ \hline 2 & 3 & 3 & \\ \hline & & & 4 \\ \hline \end{array}$ has reading $w = 32313124$.

Then $w^* = 13424232$, and the column insertion of this word is $T^{Ev} = \begin{array}{|c|c|c|c|} \hline 1 & 2 & 2 & 3 \\ \hline 2 & 4 & 4 & \\ \hline & & & 3 \\ \hline \end{array}$.

In type C , consider the KN tableau $T = \begin{array}{|c|c|c|} \hline 1 & 3 & \bar{1} \\ \hline 3 & 3 & \\ \hline & & 3 \\ \hline \end{array}$. Then, $w = cr(T) = \bar{1}3\bar{3}1\bar{3}\bar{3}$ and $w^* = 3\bar{3}\bar{1}3\bar{3}\bar{3}1$. So now we insert w^* , obtaining the following sequence of tableaux: $\begin{array}{|c|} \hline 3 \\ \hline \end{array} \quad \begin{array}{|c|} \hline 3 \\ \hline 3 \\ \hline \end{array} \quad \begin{array}{|c|} \hline 3 \\ \hline 3 \\ \hline 1 \\ \hline \end{array} \quad \begin{array}{|c|} \hline 2 & \bar{2} \\ \hline 3 \\ \hline 1 \\ \hline \end{array} \quad \begin{array}{|c|} \hline 2 & \bar{2} \\ \hline 3 & \bar{1} \\ \hline 3 \\ \hline \end{array} \quad \begin{array}{|c|} \hline 1 & 2 & \bar{2} \\ \hline 3 & \bar{1} \\ \hline 3 \\ \hline \end{array} = P(w^*)$.

Algorithm 2.

- (1) Define $T^0 := \text{complement}(\pi\text{-rotate}(T))$.
- (2) $T^{Ev} := \text{rectification of } T^0$.

Example 18. In type A , consider the tableau $T = \begin{array}{|c|c|c|c|} \hline 1 & 1 & 2 & 3 \\ \hline 2 & 3 & 3 & \\ \hline & & & 4 \\ \hline \end{array}$. After π -

rotation and complement we have the skew tableau $T^0 = \begin{array}{|c|} \hline 1 \\ \hline 2 & 2 & 3 \\ \hline 2 & 3 & 4 & 4 \\ \hline \end{array}$ which,

after rectification, gives the tableau $T^{Ev} = \begin{array}{|c|c|c|c|} \hline 1 & 2 & 2 & 3 \\ \hline 2 & 4 & 4 & \\ \hline & & & 3 \\ \hline \end{array}$.

In type C , consider the KN tableau $T = \begin{array}{|c|c|c|} \hline 1 & 3 & \bar{1} \\ \hline 3 & 3 & \\ \hline & & 3 \\ \hline \end{array}$. Then, $T_0 = \begin{array}{|c|} \hline 3 \\ \hline 3 & 3 \\ \hline 1 & 3 & \bar{1} \\ \hline \end{array}$.

So now we have to rectify this skew tableau obtaining $T^{Ev} = \begin{array}{|c|} \hline 1 & 2 & \bar{2} \\ \hline 3 & \bar{1} \\ \hline 3 \\ \hline \end{array}$.

Given a KN (SSYT) tableau T , the algorithm characterize T^{Ev} as the unique KN tableau Knuth equivalent to $wt(T)^*$ and coplactic equivalent do T .

In both Cartan types we have that algorithms 1 and 2 produce the same tableau since the column reading of T^0 is w^* , $P(w^*) = rect(T^0) = rect(w^*)$, assuming that, in type C_n , T^0 is admissible. This can be concluded using the following lemma.

Lemma 3. *For type C_n , the split of a column C , $(\ell C, rC)$ is the rotation and complement of the split of the column $C^0 = Complement(\pi\text{-rotate}(C))$, $(\ell C^0, rC^0)$.*

Proof: Let's say that $(\ell C, rC) = \begin{array}{|c|c|} \hline A & A \\ \hline B & B \\ \hline \end{array}$ where $C = \begin{array}{|c|} \hline A \\ \hline B \\ \hline \end{array}$, $\ell C = \begin{array}{|c|} \hline A \\ \hline B \\ \hline \end{array}$ and $rC = \begin{array}{|c|} \hline A \\ \hline B \\ \hline \end{array}$, where A and A' are the unbarred letters of the columns C and ℓC , respectively, and B and rB are the barred letters of C and rC , respectively. Note that ℓC and C share the barred part and C and rC share the unbarred part.

We have that $C^0 = \begin{array}{|c|} \hline B^0 \\ \hline A^0 \\ \hline \end{array}$ and its split $(\ell C^0, rC^0) = \begin{array}{|c|c|} \hline B^0 & B^0 \\ \hline A^0 & A^0 \\ \hline \end{array}$. Ignoring bars and counting multiplicities, the letters that appear in C and C^0 are the same. Hence $B^{0'}$ has the same letters as B' , but they appear unbarred, hence $B^{0'} = B^0$. The same happens with $A^{0'}$ and A^0 . Now it is easy to see that $(\ell C^0, rC^0)$ is obtained from $(\ell C, rC)$ rotating and complementing. In particular $(rC)^0 = \ell C^0$ and $(\ell C)^0 = rC^0$. ■

We now set the Cartan type to be C . Given a word $w \in [\pm n]^*$, we define the w^* like in the Algorithm 1 and show that the map $*$ preserves Knuth equivalence.

Theorem 19. *Let $v, w \in [\pm n]^*$. Then $v \sim w$ if and only if $v^* \sim w^*$.*

Proof: It is enough to consider v and w only one Knuth relation apart, because all other cases are obtained by composing multiple Knuth relations. It is enough to consider each transformation applied in one direction, since the other direction is the same case, after swapping the roles of v and w .

K1 Consider $v = v_p \gamma \beta \alpha v_s$, with $\gamma < \alpha \leq \beta$ and $(\beta, \gamma) \neq (\bar{x}, x)$, where v_p is a prefix of v , v_s is a suffix of v , and $\gamma \beta \alpha$ are three consecutive letters of v . Then, $v \stackrel{K1}{\sim} w = v_p \beta \gamma \alpha v_s$. Note that $v^* = v_s^* \bar{\alpha} \bar{\beta} \bar{\gamma} v_p^*$ and

$w^\star = v_s^\star \bar{\alpha} \bar{\gamma} \bar{\beta} v_p^\star$, with $(\bar{\gamma}, \bar{\beta}) \neq (\bar{x}, x)$ and $\bar{\beta} \leq \bar{\alpha} < \bar{\gamma}$. Hence $v^\star \stackrel{K2}{\approx} w^\star$, so they are Knuth related.

K2 Consider $v = v_p \alpha \beta \gamma v_s$, with $\gamma \leq \alpha < \beta$ and $(\beta, \gamma) \neq (\bar{x}, x)$, where v_p is a prefix of v , v_s is a suffix of v , and $\alpha \beta \gamma$ are three consecutive letters of v . Then, $v \stackrel{K2}{\approx} w = v_p \alpha \gamma \beta v_s$. Note that $v^\star = v_s^\star \bar{\gamma} \bar{\beta} \bar{\alpha} v_p^\star$ and $w^\star = v_s^\star \bar{\beta} \bar{\gamma} \bar{\alpha} v_p^\star$, with $(\bar{\gamma}, \bar{\beta}) \neq (\bar{x}, x)$ and $\bar{\beta} < \bar{\alpha} \leq \bar{\gamma}$. Hence $v^\star \stackrel{K1}{\approx} w^\star$, so they are Knuth related.

K3 Consider $v = v_p (y+1) \bar{y} + \bar{1} \beta v_s$, with $y < \beta < \bar{y}$, where v_p is a prefix of v , v_s is a suffix of v , and $(y+1) \bar{y} + \bar{1} \beta$ are three consecutive letters of v . Then, $v \stackrel{K3}{\approx} w = v_p \bar{y} y \beta v_s$. Note that $v^\star = v_s^\star \bar{\beta} (y+1) \bar{y} + \bar{1} v_p^\star$ and $w^\star = v_s^\star \bar{\beta} \bar{y} y v_p^\star$, with $y < \beta < \bar{y}$. Hence $v^\star \stackrel{K4}{\approx} w^\star$, so they are Knuth related.

K4 Consider $v = v_p \alpha \bar{x} x v_s$, with $x < \alpha < \bar{x}$, where v_p is a prefix of v , v_s is a suffix of v , and $\alpha \bar{x} x$ are three consecutive letters of v . Then, $v \stackrel{K4}{\approx} w = v_p \alpha (x+1) \bar{x} + \bar{1} v_s$. Note that $v^\star = v_s^\star \bar{x} x \bar{\alpha} v_p^\star$ and $w^\star = v_s^\star (x+1) \bar{x} + \bar{1} \bar{\alpha} v_p^\star$, with $x < \alpha < \bar{x}$. Hence $v^\star \stackrel{K3}{\approx} w^\star$, so they are Knuth related.

K5 Consider w and $\{z, \bar{z}\} \in w$ such that $w \stackrel{K5}{\approx} w \setminus \{z, \bar{z}\}$. It is clear to see that a word v breaks the 1CC at z if and only if v^\star breaks the 1CC at z . So, if w is non admissible and all its factors are admissible then the same will happen to w^\star , because all of its factors are obtained after applying * to a factor of w . So we have that $w^\star \stackrel{K5}{\approx} w^\star \setminus \{z, \bar{z}\}$.

Hence the word operator * preserves Knuth equivalence. ■

Consider a KN tableau T with column reading w . The column reading of the tableau obtained after applying Algorithm 1 to T is Knuth-related to w^\star , because both give the same tableau if inserted. Since * is an involution ($(w^\star)^\star = w$), if we apply the algorithm again we will get a tableau whose column reading, by the last theorem, is Knuth equivalent to $(w^\star)^\star = w$, hence we will have T again. So Algorithm 1 is an involution. Next we conclude that algorithms 1 and 2 is a realization of the Lusztig involution for type C .

Theorem 20. *Let $w \in [\pm n]^\star$. The connected component of the crystal G_n that contains the word w is isomorphic to the one that contains the word w^\star . Therefore $P(w)$ and $P(w^\star)$ have the same shape and weights of opposite*

sign. Moreover, the two crystals are dual of each other and the \star map is a realization of the dual crystal.

Proof: Remember the crystal operators e_i and f_i from the definition of crystal. Note that $((w)f_i)^\star = (w^\star)e_i$, because in the signature rule applied to w and w^\star , the distance of the leftmost unbracketed $+$ of w to the beginning of the word is equal to the distance of the rightmost unbracketed $-$ of w^\star to the end of this word. Hence, the letter that changes when applying f_i to w is the complement of the letter that changes when applying e_i to w^\star , and the letter obtained on their position after applying the crystal operators are also complement of each other. Hence the crystal that contains the word w^\star is the dual to the one that contains w . But the crystal that contains w is self-dual, hence the crystals that contains any of the words are isomorphic. From [18, Theorem 3.2.8] $P(w)$ and $P(w^\star)$ have the same shape. ■

5.2. Right and left keys and Lusztig involution. The next result shows that the right and left key maps defined for KN tableaux anticommutes with the Lusztig involution. The evacuation of the right key of a tableau is the left key of the evacuation of the same tableau.

Proposition 21. *Let T be a KN tableau and Ev the type C Lusztig involution. Then*

$$K_+(T)^{Ev} = K_-(T^{Ev}).$$

Proof: Since the tableaux $K_+(T)$ and $K_-(T^{Ev})$ are a key tableaux, they are completely determined by their weights. Then we just need to prove that their weights are symmetric.

Fix a column C of $K_+(T)$. There is a frank word w , Knuth related to $cr(T)$, such that C is the right column of the first column of w . Let's say the w_k is the first column of w . From Proposition 19, w^\star is Knuth related to $cr(T)^\star$, hence $P(w^\star) = T^{Ev}$. Also note that the w^\star has the same number of columns of each length as w , hence it is a frank word, and its last column is w_k^\star . Note that Lemma 3 implies that if v is an admissible column, then $l(v^\star) = (rv)^\star$. So we have that $l(w_k^\star) = (rw_k)^\star$ is a column of $K_-(T^{Ev})$. Therefore, for each column C of $K_+(T)$ there is a column of $K_-(T^{Ev})$ whose weight is $(C)\omega_0$, hence $K_+(T)$ and $K_-(T^{Ev})$ have symmetric weights. ■

6. Final Remarks

In [26], Mason showed that Demazure atoms are specializations of non-symmetric Macdonald polynomials of type A with $q = t = 0$. This allowed to use the shapes of semi-skyline augmented fillings, in the combinatorial formula of non-symmetric Macdonald polynomials [9], which are in bijection with semi standard Young tableaux, to detect the right keys. It would be interesting to obtain a similar object for a KN tableau in type C . For example, semi-skyline augmented fillings have been instrumental to obtain a RSK type bijective proof [1] for the Lascoux non-symmetric Cauchy identity in type A [15].

7. Acknowledgments

This work was partially supported by the Center for Mathematics of the University of Coimbra - UID/MAT/00324/2019, funded by the Portuguese Government through FCT/MEC and co-funded by the European Regional Development Fund through the Partnership Agreement PT2020. It was also supported by FCT, through the grant PD/BD/142954/2018, under POCH funds, co-financed by the European Social Fund and Portuguese National Funds from MEC.

We are grateful to C. Lenart for his comments on the alcove path model.

References

1. O. Azenhas, A. Emami, *An analogue of the Robinson-Schensted-Knuth correspondence and non-symmetric Cauchy kernels for truncated staircases*, Eur. J. Combin. 46 16–44, 2015.
2. O. Azenhas, R. Mamede, *Key polynomials of type C* , slides from the presentation in Séminaire Lotharingien de Combinatoire 69, Strobl, 2012, <https://www.mat.univie.ac.at/slsc/>
3. T. H. Baker, *An insertion scheme for C_n crystals*, in M. Kashiwara and T. Miwa, eds., Physical Combinatorics, Birkhäuser, Boston, Vol. 191, 1–48, 2000.
4. A. Bjorner, F. Brenti, *Combinatorics of Coxeter Groups*, Springer (2005).
5. D. Bump, A. Schilling, *Crystal Bases. Representations and Combinatorics*, World Scientific Publishing Co. Pte. Ltd., 2017.
6. C. De Concini, *Symplectic standard tableaux*, Advances in Math. 34, 1–27, 1979
7. M. Demazure, *Une formule des caractères*, Bull.Sc.M. 98, 163–172, 1974.
8. W. Fulton, *Young Tableaux: With Applications to Representation Theory and Geometry*, Cambridge: Cambridge University Press, 1996.
9. J. Haglund, M. Haiman, N. Loehr, *A combinatorial formula for non-symmetric Macdonald polynomials*, Amer. J. Math. 13: 359–383, 2008.
10. N. Jacon, C. Lecouvey, *Keys and Demazure crystals for Kac-Moody algebras*, In: ArXiv e-prints. arXiv:1909.09520 [math.CO], 2019.

11. M. Kashiwara, *The crystal base and Littelmann's refined Demazure character formula*, Duke Math. J. 71 (3), 839–858, 1993.
12. M. Kashiwara, *On crystal bases*, Representations of groups (Banff, AB, 1994), 155–197, CMS Conf. Proc., 16, Amer. Math. Soc., Providence, RI, 1995.
13. M. Kashiwara, T. Nakashima, *Crystal graphs for representations of the q -analogue of classical Lie algebras*, J. Algebra 165, 295–345, 1994.
14. R. C. King, *Weight multiplicities for the classical groups*, Lecture Notes in Physics 50, 490–499, New York, Springer, 1975.
15. A. Lascoux, *Double crystal graphs*, Studies in Memory of Issai Schur, in: Progr. Math. Birkhäuser, 210: 95–114, 2003.
16. A. Lascoux, B. Leclerc, J. Y. Thibon. *Crystal graphs and q -analogs of weight multiplicities for the root system A_n* , Lett. Math. Phys. 35, 359–374, 1995.
17. A. Lascoux, M.-P. Schützenberger, *Keys & standard bases*, IMA Vol. Math. Appl., 19, Springer, New York, 125–144, 1990.
18. C. Lecouvey, *Schensted-type correspondence, plactic monoid, and jeu de taquin for type C_n* , J. Algebra 247, no. 2, 295–331, 2002.
19. C. Lecouvey, *Combinatorics of crystal graphs for the root systems of types A_n , B_n , C_n , D_n , G_2* , in Combinatorial Aspects of Integrable Systems, MSJ Memoirs vol 17, 11–41, 2007.
20. C. Lenart, *From Macdonald polynomials to a charge statistic beyond type A*, J. Combin. Theory Ser. A 119, 683–712, 2012.
21. C. Lenart and A. Lubovsky, *A generalization of the alcove model and its applications*, J. Algebraic Combin. 41, 751–783, 2015.
22. S. J. Lee, *Crystal structure on King tableaux and semistandard oscillating tableaux*, In: ArXiv e-prints. arXiv:1910.04459 [math.CO], 2019.
23. C. Lenart, *On the combinatorics of crystal graphs. I. Lusztig's involution*, Adv. Math. 211, no. 1, 204–243, 2007.
24. P. Littelmann, *Crystal graphs and Young tableaux*, J. Algebra, 175(1):65–87, 1995.
25. P. Littelmann, *A plactic algebra for semisimple Lie algebras*, Adv. in Math., 124, 312–331, 1996.
26. S. Mason, *An explicit construction of type A Demazure atoms*, J. Algebraic Combin. 29, no. 3, 295–313, 2009.
27. R. A. Proctor, *Interactions between combinatorics, Lie theory and algebraic geometry via the Bruhat orders*, Thesis (Ph.D.), MIT, 1981.
28. M.-P. Schützenberger, *La correspondance de Robinson*, in Combinatoire et représentation du groupe symétrique, Actes Table Ronde CNRS, Univ. Louis-Pasteur Strasbourg, Strasbourg, 1976
29. J. T. Sheats, *A symplectic jeu de taquin bijection between the tableaux of King and of De Concini*, Transactions of the AMS, Vol. 351, No. 9, 3569–3607, 1999.
30. A. Young, *On quantitative substitutional analysis*, IX. Proc. London Math. Soc. (2) 54, 219–253 1952.

JOÃO MIGUEL SANTOS

CMUC, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF COIMBRA, APARTADO 3008, 3001–454
COIMBRA, PORTUGAL

E-mail address: jmsantos@mat.uc.pt