

CONTINUOUS EXTENSIONS OF REAL FUNCTIONS ON ARBITRARY SUBLOCALES AND C -, C^* - AND z -EMBEDDINGS

ANA BELÉN AVILEZ AND JORGE PICADO

ABSTRACT: This paper extends the extension theorem for localic real functions of [J. Gutiérrez García and T. Kubiak, General insertion and extension theorems for localic real functions, *J. Pure Appl. Algebra* 215 (2011) 1198-1204] from complemented sublocales to arbitrary sublocales. As an application, the theory of point-free C -, C^* - and z -embeddings is revisited.

KEYWORDS: Frame, locale, sublocale, cozero sublocale, completely separated sublocales, C - and C^* -embedded sublocales, z -embedded sublocale, localic real function, localic extension theorem.

2020 MATHEMATICS SUBJECT CLASSIFICATION: 06D22, 18F70, 54C20, 54C30, 54C45.

Introduction

One of the main differences between classical topology and point-free topology is that every subspace of a space has a complement (i.e. the lattice of subspaces of a space is a Boolean algebra) whereas most sublocales of a locale are not complemented (making the lattice of sublocales of a locale more complicated than its classical counterpart [15]).

The following point-free counterpart of the extension theorem of Mrówka ([13, 5]) was proved in [8] for a *complemented* sublocale S of a locale L :

Theorem. *The following statements about a bounded continuous real-valued function f in S are equivalent.*

- (i) *There exists a continuous extension of f to L .*
- (ii) *For every pair $r < s$ in \mathbb{Q} , the sublocales $f(r, -)$ and $f(-, s)$ are completely separated in L .*

Received November 1, 2020.

The authors gratefully acknowledge the financial support from the Centre for Mathematics of the University of Coimbra (UIDB/00324/2020, funded by the Portuguese Government through FCT/MCTES). The first author also acknowledges the support of a PhD grant from FCT/MCTES (PD/BD/150353/2019).

The main purpose of this paper is to present a proof of this theorem for *arbitrary* sublocales S . We take the opportunity to revisit the theory of C - and C^* -quotients of [2] from the point of view of sublocale embeddings (in the vein of [9, 12]). In addition, we also treat the parallel class of z -embedded sublocales [12] (*coz*-onto frame quotients in [2, 6]). Our purpose is to illustrate what the sublocale formulation adds to the theory of C -, C^* - and z -quotients.

The outline of the paper is as follows. The first two sections concern notation, terminology and background on real-valued functions in locales. Section 3 discusses the main concept of the paper: complete separation of sublocales. The main result, the general extension theorem, is proved at the fourth section. In Section 5 we present, in the language of sublocales, some needed facts about the product of two functions and the existence of the corresponding multiplicative inverse. Section 6 contains characterizations of C - and C^* -embedded sublocales and Section 7 discusses z -embedded sublocales.

1. Preliminaries and notation

1.1. The categories of frames and locales. In point-free topology, topological spaces are replaced by locales, seen as generalised spaces where points are not explicitly mentioned. The relevant categories are the category **Frm** of frames and frame homomorphisms and its dual category **Loc** of locales and localic maps. Our notation and terminology for frames and locales is that of [14] (we refer in particular to Appendix 1 for our notation on posets and lattices). We recall here some of the basic notions involved.

A *frame* (or *locale*) L is a complete lattice in which

$$a \wedge \bigvee S = \bigvee \{a \wedge b \mid b \in S\} \quad \text{for any } a \in L \text{ and } S \subseteq L. \quad (1.1.1)$$

A *frame homomorphism* preserves all joins (in particular, the bottom element 0 of the lattice) and all finite meets (in particular, the top element 1).

In a frame L the mappings $(x \mapsto (a \wedge x)): L \rightarrow L$ preserve suprema and hence they have right Galois adjoints $(y \mapsto (a \rightarrow y)): L \rightarrow L$, satisfying

$$a \wedge x \leq y \quad \text{iff} \quad x \leq a \rightarrow y$$

and making L a complete Heyting algebra. The *pseudocomplement* of $a \in L$ is the element $a^* = a \rightarrow 0 = \bigvee \{x \mid x \wedge a = 0\}$.

Let L be a frame, $a \in L$ and $r_a: L \rightarrow \downarrow a$ the surjective frame homomorphism defined by $x \mapsto x \wedge a$. We shall need the following basic property of r_a .

Lemma. *Let $h: L \rightarrow M$ be a frame homomorphism. If $h(a) = 1$ then there is a (unique) frame homomorphism $\bar{h}: \downarrow a \rightarrow M$ such that the diagram*

$$\begin{array}{ccc} L & \xrightarrow{h} & M \\ r_a \downarrow & \nearrow \bar{h} & \\ \downarrow a & & \end{array}$$

commutes.

The *rather below relation* \prec_L in a frame L (briefly \prec when there is no danger of confusion) is defined by $b \prec a$ iff $b^* \vee a = 1$. The *completely below relation* $\prec\prec_L$ (or just $\prec\prec$) is the interpolative modification of the rather below relation. Elements $a, b \in L$ satisfy $b \prec\prec a$ if and only if there exists a subset $\{a_q \mid q \in [0, 1] \cap \mathbb{Q}\} \subseteq L$ with $a_0 = b$ and $a_1 = a$ such that $a_p \prec a_q$ whenever $p < q$ in $[0, 1] \cap \mathbb{Q}$. A frame L is *completely regular* if $a = \bigvee \{b \in L \mid b \prec\prec a\}$ for any $a \in L$.

A frame is *normal* if for any $a, b \in L$ such that $a \vee b = 1$ there are $u, v \in L$ such that $a \vee u = 1 = b \vee v$ and $u \wedge v = 0$. In a normal frame, \prec interpolates hence it coincides with $\prec\prec$.

It is of advantage to represent the category of locales as a concrete category as follows. Since frame homomorphisms $h: M \rightarrow L$ preserve all joins they have uniquely defined right adjoints $f = h_*: L \rightarrow M$. We will represent **Loc** as the category with frames for objects (in this context we often speak of frames as of locales) and meet preserving maps $f: L \rightarrow M$ such that f^* are frame homomorphisms (the *localic maps*) for morphisms. They are characterized by the following conditions:

$$\begin{aligned} & \text{a meet preserving } f: L \rightarrow M \text{ is a localic map iff } f(a) = 1 \Rightarrow \\ & a = 1 \text{ and } f(f^*(a) \rightarrow b) = a \rightarrow f(b). \end{aligned}$$

1.2. The coframe of sublocales. A *sublocale* of a locale L is a subset $S \subseteq L$ closed under arbitrary meets such that

$$\forall x \in L \quad \forall s \in S \quad (x \rightarrow s \in S).$$

These are precisely the subsets of L for which the embedding $S \hookrightarrow L$ is a localic map. For alternative representations of sublocales in the literature (namely, *frame quotients* or *nuclei*) see [14, III.5].

The system $\mathbf{S}(L)$ of all sublocales of L , partially ordered by inclusion, is a *coframe*, that is, its dual lattice is a frame (see [14, Theorem III.3.2.1] for a proof). Infima and suprema are given by

$$\bigwedge_{i \in J} S_i = \bigcap_{i \in J} S_i, \quad \bigvee_{i \in J} S_i = \{\bigwedge M \mid M \subseteq \bigcup_{i \in J} S_i\}.$$

The least element is the *void sublocale* $\mathbf{O} = \{1\}$ and the greatest element is the entire locale L . Being a co-Heyting algebra, $\mathbf{S}(L)$ has co-pseudocomplements (usually called *supplements*), that we denote by $S^\#$.

We shall need the following property (that is valid in any coframe):

$$S \cap T = \mathbf{O} \Rightarrow S \subseteq T^\# \tag{1.2.1}$$

(see [7] for more information on supplements in $\mathbf{S}(L)$).

For any $a \in L$, the sublocales

$$\mathbf{c}_L(a) = \uparrow a = \{x \in L \mid x \geq a\} \quad \text{and} \quad \mathbf{o}_L(a) = \{a \rightarrow b \mid b \in L\}$$

are the *closed* and *open* sublocales of L , respectively (that we shall denote simply by $\mathbf{c}(a)$ and $\mathbf{o}(a)$ when there is no danger of confusion). For each $a \in L$, $\mathbf{c}(a)$ and $\mathbf{o}(a)$ are complements of each other in $\mathbf{S}(L)$ and satisfy the identities

$$\bigcap_i \mathbf{c}(a_i) = \mathbf{c}(\bigvee_i a_i), \quad \mathbf{c}(a) \vee \mathbf{c}(b) = \mathbf{c}(a \wedge b), \tag{1.2.2}$$

$$\bigvee_i \mathbf{o}(a_i) = \mathbf{o}(\bigvee_i a_i) \quad \text{and} \quad \mathbf{o}(a) \cap \mathbf{o}(b) = \mathbf{o}(a \wedge b). \tag{1.2.3}$$

Let S be a sublocale of L and let j_S^* be the left adjoint of the localic embedding $j_S: S \hookrightarrow L$, which is given by $j_S^*(a) = \bigwedge \{s \in S \mid s \geq a\}$. The closed (resp. open) sublocales $\mathbf{c}_S(a)$ (resp. $\mathbf{o}_S(a)$) of S ($a \in S$) are precisely the intersections $\mathbf{c}(a) \cap S$ (resp. $\mathbf{o}(a) \cap S$) and we have, for any $a \in L$,

$$\mathbf{c}(a) \cap S = \mathbf{c}_S(j_S^*(a)) \quad \text{and} \quad \mathbf{o}(a) \cap S = \mathbf{o}_S(j_S^*(a)). \tag{1.2.4}$$

The *closure* \overline{S} of a sublocale S is the smallest closed sublocale containing S , and the *interior* $\text{int } S$ is the largest open sublocale contained in S . There is a particularly simple formula for the closure, namely $\overline{S} = \mathbf{c}(\bigwedge S)$. Hence $\overline{\mathbf{o}(a)} = \mathbf{c}(a^*)$ and, consequently, $\text{int } \mathbf{c}(a) = \mathbf{o}(a^*)$. Note that $a \prec b$ iff $\overline{\mathbf{o}(a)} \subseteq \mathbf{o}(b)$: indeed, $a^* \vee b = 1$ iff $\mathbf{c}(a^*) \cap \mathbf{c}(b) = \mathbf{O}$ iff $\overline{\mathbf{o}(a)} \cap \mathbf{c}(b) = \mathbf{O}$ iff $\overline{\mathbf{o}(a)} \subseteq \mathbf{o}(b)$.

1.3. Images and preimages. For any localic map $f: L \rightarrow M$ and any sublocale $S \subseteq L$ the standard set-theoretical image $f[S]$ is a sublocale of M . However the localic preimage $f_{-1}[T]$ of a sublocale $T \subseteq M$ does not coincide in general with the set-theoretical preimage $f^{-1}[T]$. It is given by

$$f_{-1}[T] = \bigvee \{S \mid S \in \mathbf{S}(L), S \subseteq f^{-1}[T]\}.$$

In particular, for a localic embedding $j: S \hookrightarrow L$, $j_{-1}[T] = T \cap S$.

One has the adjunction

$$\mathbf{S}(L) \begin{array}{c} \xrightarrow{f[-]} \\ \perp \\ \xleftarrow{f_{-1}[-]} \end{array} \mathbf{S}(M)$$

(since $f[S] \subseteq T$ iff $S \subseteq f_{-1}[T]$). The right adjoint $f_{-1}[-]$ is a coframe homomorphism (that is, $f_{-1}[-]: \mathbf{S}(M)^{op} \rightarrow \mathbf{S}(L)^{op}$ is a frame homomorphism) while $f[-]$ is a colocalic map.

(Localic) preimages of open resp. closed sublocales are open resp. closed and one has

$$f_{-1}[\mathbf{o}(a)] = \mathbf{o}(f^*(a)) \quad \text{and} \quad f_{-1}[\mathbf{c}(a)] = f^{-1}[\mathbf{c}(a)] = \mathbf{c}(f^*(a)). \quad (1.3.1)$$

(where f^* denotes the left adjoint of the localic map f).

1.4. The frame of reals. Recall the frame of reals $\mathfrak{L}(\mathbb{R})$ from [3]. Here we define it, equivalently, as the frame presented by generators $(r, -)$ and $(-, r)$ for all rationals r , and relations

$$(r1) \quad (p, -) \wedge (-, q) = 0 \quad \text{if } q \leq p,$$

$$(r2) \quad (p, -) \vee (-, q) = 1 \quad \text{if } p < q,$$

$$(r3) \quad (p, -) = \bigvee_{r > p} (r, -),$$

$$(r4) \quad (-, q) = \bigvee_{s < q} (-, s),$$

$$(r5) \quad \bigvee_{p \in \mathbb{Q}} (p, -) = 1,$$

$$(r6) \quad \bigvee_{q \in \mathbb{Q}} (-, q) = 1.$$

Note that $(-, q)^* = (q, -)$ and $(p, -)^* = (-, p)$.

For each $p < q$ in \mathbb{Q} , the element $(p, -) \wedge (-, q)$ in $\mathfrak{L}(\mathbb{R})$ is denoted by (p, q) . The *open interval frame* $\mathfrak{L}(p, q)$ is the frame

$$\mathfrak{L}(p, q) = \downarrow(p, q) = \{a \in \mathfrak{L}(\mathbb{R}) \mid a \leq (p, q)\}.$$

Remark. It should be remarked that

$$\mathfrak{L}(p, q) \cong \mathfrak{L}(\mathbb{R}).$$

For the proof, consider any order isomorphism ψ (with inverse φ) from $\langle p, q \rangle$ into \mathbb{Q} (where $\langle \cdot, \cdot \rangle$ stands for open interval in \mathbb{Q}) and define

$$\Phi: \mathfrak{L}(\mathbb{R}) \rightarrow \mathfrak{L}(p, q)$$

on generators by $\Phi(r, -) = (\varphi(r), q)$ and $\Phi(-, r) = (p, \varphi(r))$. It is straightforward to check that Φ turns defining relations (r1)–(r6) into identities in $\mathfrak{L}(p, q)$, hence it is a frame homomorphism, clearly surjective. Finally, define $\Psi_0: \mathfrak{L}(\mathbb{R}) \rightarrow \mathfrak{L}(\mathbb{R})$ by

$$\Psi_0(r, s) = \begin{cases} 1 & \text{if } r \leq p < q \leq s, \\ (-, \psi(s)) & \text{if } r \leq p < s < q, \\ (\psi(r), \psi(s)) & \text{if } p < r < s < q, \\ (\psi(r), -) & \text{if } p < r < q \leq s \\ 0 & \text{if } s \leq p \text{ or } q \leq r. \end{cases}$$

Again, it is straightforward to check that this is a frame homomorphism. Since $\Psi_0(p, q) = 1$, the restriction of Ψ_0 to $\mathfrak{L}(p, q)$ is, by Lemma 1.1, a frame homomorphism $\Psi: \mathfrak{L}(p, q) \rightarrow \mathfrak{L}(\mathbb{R})$, inverse to Φ .

1.5. Continuous real functions and scales. The ℓ -ring $\mathcal{R}(L)$ of *continuous real-valued functions* ([3]) on a frame L is the set of all frame homomorphism $f: \mathfrak{L}(\mathbb{R}) \rightarrow L$. Each element of $\mathcal{R}(L)$ is uniquely determined by a map defined on the generators of $\mathfrak{L}(\mathbb{R})$ that turns relations (r1)–(r6) into identities in L .

A *descending scale* (resp. *ascending scale*) in L is a family $(a_p)_{p \in \mathbb{Q}} \subseteq L$ such that

$$(S1) \quad p < q \Rightarrow a_q \prec a_p \text{ (resp. } a_p \prec a_q).$$

$$(S2) \quad \bigvee_{p \in \mathbb{Q}} a_p = 1 = \bigvee_{p \in \mathbb{Q}} a_p^*.$$

Remark. If all the a_p 's are complemented then $a_p \prec a_p$ for any p and thus condition (S1) amounts only to $p < q \Rightarrow a_q \leq a_p$ (resp. $a_p \leq a_q$).

Proposition 1.5.1. *Let $f: \mathfrak{L}(\mathbb{R}) \rightarrow L$ be a continuous real function on L . Then:*

- (1) *The family $(f(p, -))_{p \in \mathbb{Q}}$ is a descending scale in L .*
- (2) *The family $(f(-, q))_{q \in \mathbb{Q}}$ is an ascending scale in L .*

Proof: Let $p < q$. Then, by (r1), $f(q, -)^* \vee f(p, -) \geq f(-, q) \vee f(p, -) = f(1) = 1$. Clearly, $\bigvee_{p \in \mathbb{Q}} f(p, -) = f(1) = 1$ and

$$\bigvee_{p \in \mathbb{Q}} f(p, -)^* \geq \bigvee_{p \in \mathbb{Q}} f(-, p) = f(1) = 1. \quad \blacksquare$$

Conversely, we have:¹

Proposition 1.5.2. (1) *Let $(a_p)_{p \in \mathbb{Q}}$ be a descending scale in L . Then the formulas*

$$f(p, -) = \bigvee_{r > p} a_r \quad \text{and} \quad f(-, q) = \bigvee_{s < q} a_s^*$$

define a frame homomorphism $f: \mathfrak{L}(\mathbb{R}) \rightarrow L$.

(2) *Let $(a_p)_{p \in \mathbb{Q}}$ be an ascending scale in L . Then the formulas*

$$g(p, -) = \bigvee_{r > p} a_r^* \quad \text{and} \quad g(-, q) = \bigvee_{s < q} a_s$$

define a frame homomorphism $g: \mathfrak{L}(\mathbb{R}) \rightarrow L$.

1.6. Product of two functions. The *product* $f \cdot g$ of two continuous real functions $f, g: \mathfrak{L}(\mathbb{R}) \rightarrow L$ is given by the formula (see [3, Chapter 4])

$$(f \cdot g)(p, q) = \bigvee \{f(r, s) \wedge g(t, u) \mid \langle r, s \rangle \cdot \langle t, u \rangle \subseteq \langle p, q \rangle\} \quad (1.6.1)$$

where $\langle \cdot, \cdot \rangle$ denotes open intervals in \mathbb{Q} and the inclusion on the right means that $x \cdot y \in \langle p, q \rangle$ whenever $x \in \langle r, s \rangle$ and $y \in \langle t, u \rangle$.

1.7. Cozero and zero sublocales. The σ -frame² $\text{Coz } L \subseteq L$ of cozero elements plays an important role in the theory of continuous real functions ([3]). Recall that a *cozero element* of L is an element of the form $f((-, 0) \vee (0, -))$ for some frame homomorphism $f: \mathfrak{L}(\mathbb{R}) \rightarrow L$ (usually denoted as $\text{coz}(f)$). Cozero elements can be described without reference to the frame of reals as follows: $a \in L$ is a cozero element if and only if $a = \bigvee_{n=1}^{\infty} a_n$ for some $a_n \prec\prec a$, $n = 1, 2, \dots$

$\text{Coz } L$ is a *normal σ -frame*, that is, $a \vee b = 1$ ($a, b \in \text{Coz } L$) implies there exist c and d in $\text{Coz } L$ such that $a \vee c = 1 = b \vee d$ and $c \wedge d = 0$ ([4]).

¹It is easy to check that the given formulas turn the defining relations (r1)–(r6) of $\mathfrak{L}(\mathbb{R})$ into identities in L .

²A σ -frame is a lattice L in which all *countable* subsets have a join such that the distribution law (1.1.1) holds for any $a \in L$ and any countable $S \subseteq L$.

The *cozero sublocales* (resp. *zero sublocales* [12]) are the $\mathbf{c}(a)$ (resp. $\mathbf{o}(a)$) with $a \in \text{Coz } L$. We denote by

$$\text{CoZS}(L) \quad \text{and} \quad \text{ZS}(L)$$

the classes of cozero and zero sublocales respectively. The intersection

$$\text{CoZS}(L) \cap \text{ZS}(L)$$

is the class of clopen sublocales.

Any cozero sublocale is a G_δ -*sublocale*, that is, a countable intersection (in $\mathbf{S}(L)$) of open sublocales, while every zero sublocale is an F_σ -*sublocale*, that is, a countable join of closed sublocales (see [12, 5.3.1]).

2. Background on general localic real functions

2.1. General real functions and scales. Let L be a frame and consider its *assembly frame*, that is, the dual frame $\mathbf{S}(L)^{op}$ of the coframe of sublocales of L . Meets and joins in $\mathbf{S}(L)^{op}$ are given by respectively

$$\prod_{i \in J} S_i = \bigvee_{i \in J} S_i \quad \text{and} \quad \bigsqcup_{i \in J} S_i = \bigcap_{i \in J} S_i.$$

By the identities in (1.2.2), the set of all closed sublocales of L form a subframe of $\mathbf{S}(L)^{op}$ isomorphic to the given L . Hence the ℓ -ring $\mathcal{R}(\mathbf{S}(L)^{op})$ is an extension of $\mathcal{R}(L)$, regarded as the ring of general real functions on L and denoted simply by $\mathbf{F}(L)$ (see [9, 11] for motivation and more information). It is partially ordered by

$$f \leq g \quad \text{iff} \quad f(-, r) \subseteq g(-, r) \quad \text{iff} \quad g(r, -) \subseteq f(r, -)$$

for all $r \in \mathbb{Q}$.

Note that a descending scale in the frame $\mathbf{S}(L)^{op}$ is a family $(S_p)_{p \in \mathbb{Q}}$ of sublocales of L satisfying

(S1) $p < q \Rightarrow S_q \prec S_p$ (i.e. $S_q^\# \cap S_p = \mathbf{0}$), and

(S2) $\bigcap_{p \in \mathbb{Q}} S_p = \mathbf{0} = \bigcap_{p \in \mathbb{Q}} S_p^\#$.

We will need the following two facts from [9, 4.4]:

Proposition. *Let $f_1, f_2 \in \mathbf{F}(L)$ be generated by descending scales $(S_r)_{r \in \mathbb{Q}}$ and $(T_r)_{r \in \mathbb{Q}}$ respectively. Then:*

- (1) $f_1(-, r)^\# \subseteq S_r \subseteq f_1(r, -)$ for every $r \in \mathbb{Q}$.
- (2) $f_2 \leq f_1$ iff $S_r \subseteq T_s$ for every $r < s$.

2.2. Semicontinuous functions. The extension $F(L)$ of $\mathcal{R}(L)$ allows to deal with more general types of real functions. In particular, an $f \in F(L)$ is *lower* (resp. *upper*) *semicontinuous* if $f(r, -)$ (resp. $f(-, r)$) is a closed sublocale for any $r \in \mathbb{Q}$. It is *continuous* if it is both lower and upper semicontinuous (that is, if $f(p, q)$ is a closed sublocale for every p, q). Of course, the subring of all continuous members of $F(L)$, denoted by $C(L)$, is an isomorphic copy of $\mathcal{R}(L)$ inside $F(L)$. Along this paper, we will work always in $F(L)$ and regard $\mathcal{R}(L)$ as the subring $C(L)$ of $F(L)$.

For any $f \in C(L)$ and $r \in \mathbb{Q}$, both $f(-, r)$ and $f(r, -)$ are cozero sublocales ([12, 5.3.1]).

2.3. Constant functions. For each $r \in \mathbb{Q}$, $(S_p^r \mid p \in \mathbb{Q})$ defined by $S_p^r = \mathbf{0}$ if $p < r$ and $S_p^r = L$ if $p \geq r$ is a descending scale in $\mathbf{S}(L)^{op}$. The corresponding function in $C(L)$, the *constant function* \mathbf{r} , is given by

$$\mathbf{r}(p, -) = \begin{cases} \mathbf{0} & \text{if } p < r \\ L & \text{if } p \geq r \end{cases} \quad \text{and} \quad \mathbf{r}(-, q) = \begin{cases} L & \text{if } q \leq r \\ \mathbf{0} & \text{if } q > r. \end{cases}$$

2.4. Bounded functions. The *bounded part* $C^*(L)$ of $C(L)$ consists of all $f \in C(L)$ such that $\mathbf{p} \leq f \leq \mathbf{q}$, that is, $f(-, p) \cap f(q, -) = L$, for some pair $p < q$ in \mathbb{Q} .

By the isomorphism between $\mathcal{R}(L)$ and $C(L)$ every cozero sublocale is of the form $\mathbf{c}(a) = f(0, -) \cap f(-, 0)$ for some $f \in C(L)$ (which, furthermore, can always be considered to be bounded); so we can always assume that a cozero sublocale is of the form $f(0, -)$ for some continuous f satisfying $\mathbf{0} \leq f \leq \mathbf{1}$.

2.5. More examples. (1) By Remark 1.5, a family $(\mathbf{c}(a_p))_{p \in \mathbb{Q}}$ of closed sublocales is a descending scale iff

- (C1) $p < q \Rightarrow a_q \leq a_p$, and
- (C2) $\bigcap_{p \in \mathbb{Q}} \mathbf{c}(a_p) = \mathbf{0} = \bigcap_{p \in \mathbb{Q}} \mathbf{o}(a_p)$.

In this case the formulas

$$f(p, -) = \bigcap_{r > p} \mathbf{c}(a_r) \quad \text{and} \quad f(-, q) = \bigcap_{s < q} \mathbf{o}(a_s)$$

given by Proposition 1.5.2 induce a lower semicontinuous function $f \in F(L)$.

(2) Similarly, a family $(\mathbf{o}(a_p))_{p \in \mathbb{Q}}$ of open sublocales is a descending scale iff

- (O1) $p < q \Rightarrow a_p \leq a_q$, and
- (O2) $\bigcap_{p \in \mathbb{Q}} \mathbf{o}(a_p) = \mathbf{0} = \bigcap_{p \in \mathbb{Q}} \mathbf{c}(a_p)$.

In this case the formulas

$$f(p, -) = \bigcap_{r>p} \mathfrak{o}(a_r) \quad \text{and} \quad f(-, q) = \bigcap_{s<q} \mathfrak{c}(a_s)$$

given by 1.5.2 induce an upper semicontinuous function $f \in \mathbf{F}(L)$.

(3) The condition $\bigcap_{p \in \mathbb{Q}} \mathfrak{c}(a_p) = \mathbf{O}$ means that $\bigvee_{p \in \mathbb{Q}} a_p = 1$ but the condition $\bigcap_{p \in \mathbb{Q}} \mathfrak{o}(a_p) = \mathbf{O}$ is generally weaker than $\bigvee_{p \in \mathbb{Q}} a_p^* = 1$. However, in case $(a_p)_{p \in \mathbb{Q}} \subseteq L$ satisfies 1.5(S1), then it is easy to see that

$$\bigcap_{p \in \mathbb{Q}} \mathfrak{o}(a_p) = \mathfrak{c}\left(\bigvee_{p \in \mathbb{Q}} a_p^*\right), \quad \bigcap_{s<q} \mathfrak{o}(a_s) = \mathfrak{c}\left(\bigvee_{s<q} a_s^*\right) \quad \text{and} \quad \bigcap_{r>p} \mathfrak{o}(a_r) = \mathfrak{c}\left(\bigvee_{r>p} a_r^*\right),$$

and, consequently, the conditions $\bigcap_{p \in \mathbb{Q}} \mathfrak{o}(a_p) = \mathbf{O}$ and $\bigvee_{p \in \mathbb{Q}} a_p^* = 1$ do coincide. Hence we have:

Proposition. (1) *Let $(a_p)_{p \in \mathbb{Q}}$ be a descending scale in L . Then the family $(\mathfrak{c}(a_p))_{p \in \mathbb{Q}}$ is a descending scale in $\mathbf{S}(L)^{op}$ and the induced function $f \in \mathbf{F}(L)$ given by*

$$f(p, -) = \bigcap_{r>p} \mathfrak{c}(a_r) = \mathfrak{c}\left(\bigvee_{r>p} a_r\right) \quad \text{and} \quad f(-, q) = \bigcap_{s<q} \mathfrak{o}(a_s) = \mathfrak{c}\left(\bigvee_{s<q} a_s^*\right)$$

is continuous.

(2) *Let $(a_p)_{p \in \mathbb{Q}}$ be an ascending scale in L . Then the family $(\mathfrak{o}(a_p))_{p \in \mathbb{Q}}$ is a descending scale in $\mathbf{S}(L)^{op}$ and the induced function $g \in \mathbf{F}(L)$ given by*

$$g(p, -) = \bigcap_{r>p} \mathfrak{o}(a_r) = \mathfrak{c}\left(\bigvee_{r>p} a_r^*\right) \quad \text{and} \quad g(-, q) = \bigcap_{s<q} \mathfrak{c}(a_s) = \mathfrak{c}\left(\bigvee_{s<q} a_s\right)$$

is continuous. ■

These are precisely (up to the isomorphism between L and the subframe of $\mathbf{S}(L)^{op}$ of all closed sublocales) the functions f and g of 1.5.2.

2.6. Continuous extensions. Let S be a sublocale of L with localic embedding $j: S \hookrightarrow L$. A function $f \in \mathbf{C}(S)$ is said to have a *continuous extension* to L if there is an $\bar{f} \in \mathbf{C}(L)$ such that the diagram

$$\begin{array}{ccc} \mathfrak{L}(\mathbb{R}) & \xrightarrow{\bar{f}} & \mathbf{S}(L)^{op} \\ & \searrow f & \downarrow j_{-1}[-]: T \mapsto T \cap S \\ & & \mathbf{S}(S)^{op} \end{array}$$

commutes (that is, $\bar{f}(a) \cap S = f(a)$ for every $a \in \mathfrak{L}(\mathbb{R})$).

3. Completely separated sublocales

Two sublocales S and T of L are said to be *completely separated in L* if

$$S \subseteq f(0, -) \quad \text{and} \quad T \subseteq f(-, 1)$$

for some $f \in \mathbf{C}(L)$ such that $\mathbf{0} \leq f \leq \mathbf{1}$.

This notion was first studied in [2] in terms of quotient maps and cozero elements and equivalently reformulated in [8] in terms of sublocales and continuous real functions. We refer to [12] for several results about completely separated sublocales. E.g., sublocales $\mathbf{c}(a)$ and $\mathbf{o}(b)$ are completely separated iff $b \prec\prec a$ ([12, Lemma 5.4.2]).

Proposition 3.1. *Two sublocales S and T of L are completely separated in L if and only if they are contained in disjoint cozero sublocales of L .*

Proof: The implication ‘ \Rightarrow ’ is obvious. Conversely, let $\mathbf{c}(a), \mathbf{c}(b) \in \mathbf{CozS}(L)$ such that $S \subseteq \mathbf{c}(a)$, $T \subseteq \mathbf{c}(b)$, and $\mathbf{c}(a) \cap \mathbf{c}(b) = \mathbf{O}$. The last identity means that $a \vee b = 1$. Since a and b are cozero elements of L and $\mathbf{Coz}(L)$ is a normal sub- σ -frame of L (recall 1.7), there exist $u, v \in \mathbf{Coz}(L)$ such that $a \vee u = 1 = b \vee v$ and $u \wedge v = 0$. This implies that $u \prec b$ and, again by normality, \prec interpolates and thus $u \prec\prec b$. Then (see e.g. [12, 5.4.3]) there is an $f \in \mathbf{C}^*(L)$ such that $\mathbf{o}(u) \subseteq f(0, -)$ and $\mathbf{c}(b) \subseteq f(-, 1)$. Since $\mathbf{c}(a) \subseteq \mathbf{o}(u)$ (because $a \vee u = 1$) we may then conclude that $S \subseteq \mathbf{c}(a) \subseteq f(0, -)$ and $T \subseteq \mathbf{c}(b) \subseteq f(-, 1)$. \blacksquare

Remark 3.2. By the well-known Urysohn’s separation lemma for locales ([3, Proposition 5]), in a normal locale any two disjoint closed sublocales are completely separated.

Let U and V be sublocales of a sublocale S of L (hence, sublocales of L). If U and V are completely separated in L with $f \in \mathbf{C}(L)$ satisfying $U \subseteq f(0, -)$ and $V \subseteq f(-, 1)$, consider the composite

$$\mathfrak{L}(\mathbb{R}) \xrightarrow{f} \mathbf{S}(L)^{op} \xrightarrow{j_{-1}[-]} \mathbf{S}(S)^{op}$$

where $j: S \hookrightarrow L$. Since $j_{-1}[-]$ is given by intersection with S , it is easy to see that $j_{-1}[-]f \in \mathbf{C}(S)$. Further, $U \subseteq f(0, -) \cap S = j_{-1}[f(0, -)]$ and $V \subseteq f(-, 1) \cap S = j_{-1}[f(-, 1)]$. Hence U and V are completely separated in S .

Hence, if two sublocales of S are completely separated in L , then they are completely separated in S . Of course, the converse does not hold in general:

two sublocales that are completely separated in S may not be completely separated in L as some easy examples show.

More specifically, we have:

Proposition 3.3. *The following statements are equivalent for a sublocale S of L .*

- (i) *Any two completely separated sublocales of S are completely separated in L .*
- (ii) *For every $f \in \mathbf{C}^*(S)$ and every pair $r < s$ in \mathbb{Q} , $f(r, -)$ and $f(-, s)$ are completely separated in L .*

Proof: (i) \Rightarrow (ii): For any $f \in \mathbf{C}^*(S)$ and $r < s$, $f(r, -)$ and $f(-, s)$ are disjoint cozero sublocales of S :

$$f(r, -) \cap f(-, s) = f((r, -) \vee (-, s)) = f(1) = \mathbf{O}.$$

Therefore, by the previous proposition, they are completely separated in S hence in L (by assumption).

(ii) \Rightarrow (i): Let U, V be sublocales of S , completely separated in S . Then $U \subseteq f(0, -)$ and $V \subseteq f(-, 1)$ for some $f \in \mathbf{C}^*(S)$ and by assumption, $f(0, -)$ and $f(-, 1)$ are completely separated in L . Hence U and V are also completely separated in L . ■

Remark 3.4. It is an easy exercise to check that each one of the following conditions is also equivalent to the assertions in Proposition 3.3:

- (iii) *Any two closed (resp. open) sublocales of S that are completely separated in S , are completely separated in L .*
- (iv) *Any two cozero (resp. zero) sublocales of S that are completely separated in S , are completely separated in L .*
- (v) *Any open and any closed (resp. any zero and any cozero) sublocales of S that are completely separated in S , are completely separated in L .*

They are the localic formulations of some of the equivalent conditions in [2, Theorem 7.1.1] presented in terms of frame quotients. We believe that the language of sublocales helps to clarify the notions and statements. For example, it is now clear that the terminology “ m -completely separated” (with respect to a quotient m of a frame L , that is, a sublocale M of L) introduced in [2, pp. 123] is superfluous. In fact, it is precisely “complete separation”: for any sublocale M of L , two sublocales of M are M -completely separated if and only if they are completely separated in L .

Another equivalent condition in [2, Theorem 7.1.1] is formulated in terms of certain order relation

$$\prec\prec_m$$

in L . The condition $b \prec\prec_m a$ translates into the present language as the condition that the sublocales $\mathfrak{o}_S(b)$ and $\mathfrak{c}_S(a)$ of S are completely separated in L . As we observed earlier, this is stronger than saying that $\mathfrak{o}_S(b)$ and $\mathfrak{c}_S(a)$ are completely separated in S , that is, that $b \prec\prec_S a$ (by the result recalled above that in any locale L , $\mathfrak{o}_L(b)$ and $\mathfrak{c}_L(a)$ are completely separated in L iff $b \prec\prec_L a$).

4. Extension theorem for arbitrary sublocales

In this section we generalize the general extension theorem of [8] to arbitrary sublocales. To prove it we will need the following insertion theorem from [8]:

Theorem 4.1. ([8, Theorem 4.2]) *Let L be a frame and $f_1, f_2 \in \mathbf{F}(L)$. The following statements are equivalent.*

- (i) *There exists $h \in \mathbf{C}(L)$ such that $f_2 \leq h \leq f_1$.*
- (ii) *The sublocales $f_2(-, s)$ and $f_1(r, -)$ are completely separated in L for every $r < s$ in \mathbb{Q} .*

Theorem 4.2. *Let S be a sublocale of L . The following statements about an $f \in \mathbf{C}^*(S)$ are equivalent.*

- (i) *There exists a bounded continuous extension of f to L .*
- (ii) *The sublocales $f(r, -)$ and $f(-, s)$ are completely separated in L for every $r < s$ in \mathbb{Q} .*

Proof: (i) \Rightarrow (ii): If f has a bounded continuous extension to L , say \bar{f} , then, for every $r < s$,

$$f(r, -) = \bar{f}(r, -) \cap S \subseteq \bar{f}(r, -) \quad \text{and} \quad f(-, s) = \bar{f}(-, s) \cap S \subseteq \bar{f}(-, s).$$

Moreover,

$$\bar{f}(r, -) \cap \bar{f}(-, s) = \bar{f}((r, -) \vee (-, s)) = \bar{f}(1) = \mathbf{0}.$$

Hence $f(r, -)$ and $f(-, s)$ are contained in the disjoint cozero sublocales $\bar{f}(r, -)$ and $\bar{f}(-, s)$, and thus they are completely separated.

(ii) \Rightarrow (i): Let $f \in \mathbf{C}^*(S)$. We may assume that $\mathbf{0} \leq f \leq \mathbf{1}$ with no loss of generality. For each $r \in \mathbb{Q}$ set

$$S_r = \begin{cases} \mathbf{0} & \text{if } r < 0 \\ \bigcap \{C \in \mathbf{CoZS}(L) \mid f(r, -) \subseteq C\} & \text{if } 0 \leq r < 1 \\ L & \text{if } r \geq 1 \end{cases}$$

and

$$T_r = \begin{cases} \mathbf{0} & \text{if } r \leq 0 \\ \bigvee \{C \in \mathbf{ZS}(L) \mid f(-, r) \subseteq C^\#\} & \text{if } 0 < r \leq 1 \\ L & \text{if } r > 1. \end{cases}$$

Each S_r is a closed sublocale of L whereas T_r is open. For any $r < s$, we have $S_r \subseteq S_s$, since $f(r, -) \subseteq f(s, -)$, and $T_r \subseteq T_s$, since $f(-, s) \subseteq f(-, r)$, hence $T_s^\# \subseteq T_r^\#$. Note that for any $0 < r \leq 1$,

$$T_r^\# = \bigcap \{C \in \mathbf{CoZS}(L) \mid f(-, r) \subseteq C\}$$

because T_r is a join of open sublocales, that is, an open sublocale, hence its complement is simply the intersection of the complements of those open sublocales.

Further $\bigcap_{r \in \mathbb{Q}} S_r = \mathbf{0} = \bigcap_{r \in \mathbb{Q}} T_r$, hence $(S_r)_{r \in \mathbb{Q}}$ and $(T_r)_{r \in \mathbb{Q}}$ are descending scales, with corresponding functions $f_1, f_2 \in F(L)$ defined by (recall 1.5.2)

$$\begin{aligned} f_1: \mathfrak{L}(\mathbb{R}) &\rightarrow \mathbf{S}(L)^{op} & f_2: \mathfrak{L}(\mathbb{R}) &\rightarrow \mathbf{S}(L)^{op} \\ f_1(r, -) &= \bigcap_{p > r} S_p & f_2(r, -) &= \bigcap_{p > r} T_p \\ f_1(-, s) &= \bigcap_{q < s} S_q^\# & f_2(-, s) &= \bigcap_{q < s} T_q^\#. \end{aligned}$$

In particular,

$$f_1(r, -) = \begin{cases} \mathbf{0} & \text{if } r < 0 \\ \bigcap \{C \in \mathbf{CoZS}(L) \mid f(p, -) \subseteq C \text{ for some } p > r\} & \text{if } 0 \leq r < 1 \\ L & \text{if } r \geq 1 \end{cases}$$

and

$$f_2(-, s) = \begin{cases} L & \text{if } s \leq 0 \\ \bigcap \{C \in \mathbf{CoZS}(L) \mid f(-, q) \subseteq C \text{ for some } q < s\} & \text{if } 0 < s \leq 1 \\ \mathbf{0} & \text{if } s > 1. \end{cases}$$

Claim 1: $f_2 \leq f_1$.

We will show this using Proposition 2.1, by proving that $S_r \subseteq T_s$ for every $r < s$. If $r < 0$ then $S_r = \mathbf{0} \subseteq T_s$. If $s > 1$, then $S_r \subseteq L = T_s$. Finally, if $0 \leq r < s \leq 1$, then $f(r, -)$ and $f(-, s)$ are completely separated in L thus there exist disjoint $C_1, C_2 \in \text{CoZS}(L)$ such that $f(r, -) \subseteq C_1$ and $f(-, s) \subseteq C_2$ and then, by (1.2.1),

$$S_r \subseteq C_1 \subseteq C_2^\# \subseteq T_s.$$

Claim 2: *There exists $h \in \mathbf{C}^*(L)$ such that $f_2 \leq h \leq f_1$.*

By Theorem 4.1 it suffices to show that $f_1(r, -)$ and $f_2(-, s)$ are completely separated for any $r < s$. Again, the cases $r < 0$ and $s > 1$ are trivial. If $0 \leq r < s \leq 1$ consider $p, q \in \mathbb{Q}$ such that $0 \leq r < p < q < s \leq 1$. By the assumption, there are disjoint $C_1, C_2 \in \text{CoZS}(L)$ such that $f(p, -) \subseteq C_1$ and $f(-, q) \subseteq C_2$. Then

$$f_1(r, -) \subseteq S_p \subseteq C_1 \quad \text{and} \quad f_2(-, s) \subseteq T_q^\# \subseteq C_2.$$

Hence $f_1(r, -)$ and $f_2(-, s)$ are completely separated.

Claim 3: *h is a continuous bounded extension of f to L .*

(1) We need to show that $h(r, -) \cap S = f(r, -)$ for every $r \in \mathbb{Q}$. We have the following three cases:

- $r < 0$: We have $h(r, -) \subseteq f_2(r, -) = \bigcap_{p>r} T_p = \mathbf{0}$ and $f(r, -) = \mathbf{0}$ (because $\mathbf{0} \leq f$). Hence $h(r, -) \cap S = \mathbf{0} = f(r, -)$.
- $r \geq 1$: In this case, $L = f_1(r, -) \subseteq h(r, -)$ and $f(r, -) = S$ since $f \leq \mathbf{1}$. Hence $h(r, -) \cap S = L \cap S = S = f(r, -)$.
- $0 \leq r < 1$: For every $p > r$, $f(r, -) \subseteq f(p, -) \subseteq S_p$, hence $f(r, -) \subseteq f_1(r, -) \subseteq h(r, -)$ (since $h \leq f_1$). On the other hand, since $f_2 \leq h$, then

$$S \cap h(r, -) \subseteq S \cap f_2(r, -) = S \cap \bigcap_{p>r} T_p = \bigcap_{p>r} (S \cap T_p).$$

Moreover, since $f(-, p) \subseteq T_p^\#$, we have $f(-, p) \cap T_p = \mathbf{0}$. In particular, $f(-, p) \cap S \cap T_p = \mathbf{0}$. Hence, by (1.2.1), $S \cap T_p \subseteq f(-, p)^{\#s} \subseteq f(p, -)$

(where $(-)^{\#s}$ denotes supplementation in $\mathbf{S}(S)$). Note that $S \cap T_p \subseteq f(p, -)$ also holds for $p > 1$ since $T_p = L$ and $f(p, -) = S$. Hence,

$$S \cap h(r, -) \subseteq \bigcap_{p>r} (S \cap T_p) \subseteq \bigcap_{p>r} f(p, -) = f(r, -).$$

(2) Finally, we need to show that $h(-, s) \cap S = f(-, s)$ for every $s \in \mathbb{Q}$. Again it suffices to analyse the following three cases:

- $s \leq 0$: In this case, $L = f_2(-, s) \subseteq h(-, s)$ and $f(-, s) = S$ (because $\mathbf{0} \leq f$). Then $h(-, s) \cap S = L \cap S = S = f(-, s)$.
- $1 < s$: Then $h(-, s) \subseteq f_1(-, s) = \bigcap_{q<s} S_q^{\#} = \mathbf{0}$ and $f(-, s) = \mathbf{0}$ since $f \leq \mathbf{1}$. Then $h(-, s) \cap S = \mathbf{0} = f(-, s) \cap S$.
- $0 < s \leq 1$: First, we have (since $\mathbf{0} \leq f$ and $f_2 \leq h$):

$$\begin{aligned} f(-, s) &= \bigcap_{0<t<s} f(-, t) \subseteq \bigcap \{C \in \mathbf{CoZS}(L) \mid f(-, t) \subseteq C \text{ for some } 0 < t < s\} \\ &= \bigcap_{0<t<s} T_t^{\#} = f_2(-, s) \subseteq h(-, s). \end{aligned}$$

On the other hand, since $f(t, -) \subseteq S_t$ for every $t \geq 0$, we have $f(t, -) \cap S \cap S_t^{\#} = \mathbf{0}$, that is (by (1.2.1)), $S \cap S_t^{\#} \subseteq f(t, -)^{\#s} \subseteq f(-, t)$. Finally, from $h \leq f_1$ it follows that

$$S \cap h(-, s) \subseteq S \cap f_1(-, s) \subseteq \bigcap_{t<s} (S \cap S_t^{\#}) \subseteq \bigcap_{t<s} f(-, t) = f(-, s).$$

In conclusion, h is a continuous extension of f to L . ■

This theorem is the extension to arbitrary sublocales of the main theorem of [8] (proved only for complemented sublocales). It was originally stated as part of [2, Theorem 7.1.1], but the proof there requires some background results on the localic Yosida representation, complete separation in archimedean f -rings and uniformities. The proof above uses only basic facts about localic real functions and sublocale lattices.

5. The multiplicative inverse of a function

Let $f \cdot g$ denote the product of two real functions $f, g \in \mathbf{F}(L)$. It may be computed with formula (1.6.1) applied on frame $\mathbf{S}(L)^{op}$. Alternative formulas for the computation of $f \cdot g$ may be consulted in [12, 4.4] or [11, 4.3]. Here we only need to recall the particular case $f, g \geq \mathbf{0}$.

Proposition 5.1. *Let $\mathbf{0} \leq f, g \in \mathbf{F}(L)$. Then:*

$$(1) (f \cdot g)(p, -) = \begin{cases} \bigcap_{r>0} (f(r, -) \vee g(\frac{p}{r}, -)) & \text{if } p \geq 0 \\ \mathbf{0} & \text{if } p < 0. \end{cases}$$

$$(2) (f \cdot g)(-, q) = \begin{cases} \bigcap_{s>0} (f(-, s) \vee g(-, \frac{q}{s})) & \text{if } q > 0 \\ L & \text{if } q \leq 0. \end{cases}$$

We will also need the familiar fact ([12, 5.3]) that

$$\text{coz}(f \cdot g) = \text{coz}(f) \wedge \text{coz}(g). \quad (5.1.1)$$

The following proposition can be found in [2].

Proposition 5.2. *A frame homomorphism $f: \mathfrak{L}(\mathbb{R}) \rightarrow L$ has a multiplicative inverse if and only if $\text{coz}(f) = 1$.*

This result can be also proved with the point-free description of the reals as a frame presented by generators and relations. The idea for the proof is to mimick the classical proof that constructs the multiplicative inverse of a function $f: X \rightarrow \mathbb{R}$ by composing it with $g: \mathbb{R} \setminus \{0\} \rightarrow \mathbb{R}$ ($x \mapsto \frac{1}{x}$) provided the image of f is contained in $\mathbb{R} \setminus \{0\}$.

Indeed, if $\text{coz}(f) = 1$ there is by Lemma 1.1 a frame homomorphism \bar{f} such that the diagram

$$\begin{array}{ccc} \mathfrak{L}(\mathbb{R}) & \xrightarrow{f} & L \\ \downarrow & \searrow \bar{f} & \uparrow \\ \downarrow((0, -) \vee (-, 0)) & & \end{array}$$

commutes. We can compose \bar{f} with

$$g: \mathfrak{L}(\mathbb{R}) \rightarrow \mathfrak{L}(\mathbb{R} \setminus \{0\}) = \downarrow((0, -) \vee (-, 0)),$$

the point-free version of the mapping $x \mapsto \frac{1}{x}$ above, given by

$$g(p, -) = \begin{cases} (0, \frac{1}{p}) & \text{if } p > 0 \\ (0, -) & \text{if } p = 0 \\ (-, \frac{1}{p}) \vee (0, -) & \text{if } p < 0 \end{cases}$$

and

$$g(-, q) = \begin{cases} (\frac{1}{q}, 0) & \text{if } q < 0 \\ (-, 0) & \text{if } q = 0 \\ (-, 0) \vee (\frac{1}{q}, -) & \text{if } q > 0. \end{cases}$$

The composite $\bar{f}g$ is the multiplicative inverse of f . The verification details are left to the reader.

Note. Classically, when a function does not have a multiplicative inverse, one restricts it to its cozero set in order to compose it with $x \mapsto \frac{1}{x}$. Similarly, if $f: \mathfrak{L}(\mathbb{R}) \rightarrow L$ is a frame homomorphism, by 1.1 there exists \bar{f} such that the diagram

$$\begin{array}{ccc} \mathfrak{L}(\mathbb{R}) & \xrightarrow{f} & L \xrightarrow{p} \Downarrow \text{coz}(f) \\ \downarrow & \nearrow \bar{f} & \\ \mathfrak{L}(\mathbb{R} \setminus \{0\}) & & \end{array}$$

commutes. Then $\mathfrak{L}(\mathbb{R}) \xrightarrow{g} \mathfrak{L}(\mathbb{R} \setminus \{0\}) \xrightarrow{\bar{f}} L$ is the multiplicative inverse of pf .

6. C - and C^* -embedded sublocales

Recall 2.6. A sublocale S of L is said to be C -embedded (resp. C^* -embedded) if every f in $\mathbf{C}(S)$ (resp. in $\mathbf{C}^*(S)$) has a continuous extension (resp. bounded continuous extension) to L .

As an immediate consequence of the Extension Theorem 4.2 and Proposition 3.3, we have:³

Theorem 6.1. *The following statements about a sublocale S of L are equivalent.*

- (i) S is C^* -embedded in L .
- (ii) For every $f \in \mathbf{C}^*(S)$ and every pair $r < s$ in \mathbb{Q} , $f(r, -)$ and $f(-, s)$ are completely separated in L .
- (iii) Any two completely separated sublocales of S are completely separated in L . ■

³The counterpart to equivalence (i) \Leftrightarrow (iii) in the classical setting is known as the Uryshon's Extension Theorem (cf. [1, 6.6]).

Next result identifies C -embedded sublocales in among C^* -embedded sublocales.

Theorem 6.2. *The following statements about a sublocale S of L are equivalent.*

- (i) S is C -embedded in L .
- (ii) S is C^* -embedded and every cozero sublocale of L disjoint from S is completely separated from S .

Proof: Let S be a C -embedded sublocale of L . Of course, S is C^* -embedded. Consider a cozero sublocale C of L disjoint from S , say $C = f(0, -)$, for some f such that $\mathbf{0} \leq f \leq \mathbf{1}$ (recall 2.4). Then consider the composite

$$h: \mathfrak{L}(\mathbb{R}) \xrightarrow{f} \mathfrak{S}(L)^{op} \xrightarrow{j_{-1}[-]} \mathfrak{S}(S)^{op}$$

where j is the localic embedding $S \hookrightarrow L$. The cozero sublocale defined by h is

$$\text{coz}(h) = h(0, -) \cap h(-, 0) = j_{-1}[C] \cap j_{-1}[L] = \mathbf{0} \cap S = \mathbf{0}.$$

By Proposition 5.2, h has a multiplicative inverse $g: \mathfrak{L}(\mathbb{R}) \rightarrow \mathfrak{S}(S)^{op}$ and since S is C -embedded, there is an extension

$$\bar{g}: \mathfrak{L}(\mathbb{R}) \rightarrow \mathfrak{S}(L)^{op}$$

such that $j_{-1}[-]\bar{g} = g$. We may assume, without loss of generality, that $\bar{g} \geq \mathbf{0}$ (because $\bar{g} \vee \mathbf{0}$ will be also an extension of $g \geq \mathbf{0}$). Then we can use the formula for the product in 5.1 and conclude that $\bar{g} \cdot f$ completely separates the sublocales S and C . Indeed,

$$(\bar{g} \cdot f)(0, -) = \bigcap_{s>0} \bar{g}(s, -) \vee f(0, -) = C \vee \bigcap_{s>0} \bar{g}(s, -) = C \vee \bar{g}(0, -) \supseteq C$$

and

$$(\bar{g} \cdot f)(-, 1) = \bigcap_{s>0} \bar{g}(-, s) \vee f(-, \frac{1}{s})$$

from which it follows that

$$\begin{aligned} j_{-1}[(\bar{g} \cdot f)(-, 1)] &= \bigcap_{s>0} j_{-1}[\bar{g}(-, s)] \vee j_{-1}[f(-, \frac{1}{s})] = \\ &= \bigcap_{s>0} g(-, s) \vee h(-, \frac{1}{s}) = (h \cdot g)(-, 1) = \mathbf{1}(-, 1) = S \end{aligned}$$

and hence $S \cap (\bar{g} \cdot f)(-, 1) = S$, that is, $S \subseteq (\bar{g} \cdot f)(-, 1)$.

Conversely, assume that S is a C^* -embedded sublocale of L , with localic embedding $j: S \hookrightarrow L$ such that every cozero sublocale disjoint from S is

completely separated from S . In order to show that S is also C -embedded consider an $f \in \mathbf{C}(S)$. Now recall Remark 1.4 and consider an order isomorphism ψ from $\langle -1, 1 \rangle$ into \mathbb{Q} . Using the notation from 1.4, we have the commutative diagram

$$\begin{array}{ccc}
 \mathfrak{L}(\mathbb{R}) & \xrightarrow{\Psi_0} & \mathfrak{L}(\mathbb{R}) \\
 & \searrow^{r_{(-1,1)}} & \nearrow^{\Phi} \\
 & & \mathfrak{L}(-1, 1) \\
 & & \nearrow^{\Psi = \Psi_0|_{\mathfrak{L}(-1,1)}}
 \end{array}$$

The composite $f\Psi_0$ is a bounded frame homomorphism (since $\Psi_0(-1, 1) = 1$). Hence, since S is C^* -embedded, there is an $\bar{f} \in \mathbf{C}^*(L)$ (with $\mathbf{p} < \bar{f} < \mathbf{q}$) such that the diagram

$$\begin{array}{ccc}
 \mathfrak{L}(\mathbb{R}) & \xrightarrow{\exists \bar{f}} & \mathbf{S}(L)^{op} \\
 \downarrow^{r_{(-1,1)}} & \searrow^{\Psi_0} & \downarrow^{j_{-1}[-]} \\
 \mathfrak{L}(-1, 1) & \xleftarrow{\Phi} & \mathfrak{L}(\mathbb{R}) \xrightarrow{f} \mathbf{S}(S)^{op} \\
 & \searrow^{\Psi} & \\
 & & \mathbf{S}(S)^{op}
 \end{array}$$

commutes. To show that S is C -embedded it suffices to find a

$$\bar{g}: \mathfrak{L}(-1, 1) \rightarrow \mathbf{S}(L)^{op}$$

such that $\bar{g}r_{(-1,1)} = \bar{f}$, because then $\bar{g}\Phi$ will be a continuous extension of f to L :

$$f\Psi r_{(-1,1)} = j_{-1}[-]\bar{f} = j_{-1}[-]\bar{g}r_{(-1,1)} \Rightarrow f\Psi = j_{-1}[-]\bar{g} \Leftrightarrow f = j_{-1}[-]\bar{g}\Phi.$$

For that, by Lemma 1.1, it suffices to find a $g: \mathfrak{L}(\mathbb{R}) \rightarrow \mathbf{S}(L)^{op}$ such that $g(-1, 1) = \mathbf{O}$ and $j_{-1}[-]g = f\Psi_0$:

$$\begin{array}{ccc}
 \mathfrak{L}(\mathbb{R}) & \xrightarrow{\exists g} & \mathbf{S}(L)^{op} \\
 \downarrow^{r_{(-1,1)}} & \searrow^{\Psi_0} & \downarrow^{j_{-1}[-]} \\
 \mathfrak{L}(-1, 1) & \xleftarrow{\Phi} & \mathfrak{L}(\mathbb{R}) \xrightarrow{f} \mathbf{S}(S)^{op} \\
 & \searrow^{\Psi} & \\
 & & \mathbf{S}(S)^{op}
 \end{array}$$

We conclude the proof by showing how to get such map g .

Let $\bar{f}(-1, 1) = \mathbf{c}_L(a) \in \mathbf{CozS}(L)$. Then $j_{-1}[\mathbf{c}_L(a)] = f\Psi_0(-1, 1) = f(1) = \mathbf{O}$. On the other hand (recall 1.3 and (1.2.4)), $j_{-1}[\mathbf{c}_L(a)] = \mathbf{c}_S(j^*(a)) = \mathbf{c}_L(a) \cap S$. Hence, $S \cap \mathbf{c}_L(a) = \mathbf{O}$. Then, by assumption, S and $\mathbf{c}_L(a)$ are completely separated, that is, there is an

$$h: \mathfrak{L}(\mathbb{R}) \rightarrow \mathbf{S}(L)^{op} \quad (\mathbf{0} \leq h \leq \mathbf{1})$$

such that $\mathbf{c}_L(a) \subseteq h(0, -)$ and $S \subseteq h(-, 1)$. We claim that $h \cdot \bar{f}$ is the function g we are searching for. We only need to check that $(h \cdot \bar{f})(-1, 1) = \mathbf{O}$ and $j_{-1}[-](h \cdot \bar{f}) = f\Psi_0$.

By 1.6.1 we have

$$\begin{aligned} (h \cdot \bar{f})(-1, 1) &= \bigcap \{h(r, s) \vee \bar{f}(t, u) \mid \langle r, s \rangle \cdot \langle t, u \rangle \subseteq \langle -1, 1 \rangle\} \\ &\subseteq \bigcap \{h(-y, y) \vee \bar{f}(-\frac{1}{y}, \frac{1}{y}) \mid 1 < y\} \\ &\stackrel{(*)}{=} \bigcap \{\bar{f}(-\frac{1}{y}, \frac{1}{y}) \mid 1 < y\} = \bar{f}(-1, 1) \end{aligned}$$

(the equality $(*)$ follows from the fact that $\mathbf{0} \leq h \leq \mathbf{1}$). Consequently,

$$\begin{aligned} (h \cdot \bar{f})(-1, 1) &= (h \cdot \bar{f})(-1, 1) \cap \bar{f}(-1, 1) \\ &= \bigcap \{h(r, s) \vee \bar{f}(u, v) \mid \langle r, s \rangle \cdot \langle u, v \rangle \subseteq \langle -1, 1 \rangle\} \cap \bar{f}(-1, 1) \\ &\subseteq \bigcap \{h(-\frac{1}{y}, \frac{1}{y}) \vee \bar{f}(-y, y) \mid y > \max\{|p|, |q|, 1\}\} \cap \bar{f}(-1, 1) \\ &\stackrel{(*)}{=} \bigcap \{h(-\frac{1}{y}, \frac{1}{y}) \mid y > \max\{|p|, |q|, 1\}\} \cap \bar{f}(-1, 1) \\ &\subseteq \bigcap \{h(-\frac{1}{y}, \frac{1}{y}) \mid y > \max\{|p|, |q|, 1\}\} \cap h(0, -) \\ &= \bigcap \{h(-\frac{1}{y}, \frac{1}{y}) \cap h(0, -) \mid y > \max\{|p|, |q|, 1\}\} \\ &\stackrel{(**)}{=} \bigcap \{h(-\frac{1}{y}, -) \mid y > \max\{|p|, |q|, 1\}\} = \mathbf{O} \end{aligned}$$

(where $(*)$ follows from $\mathbf{p} < \bar{f} < \mathbf{q}$ and $(**)$ from $h \geq \mathbf{0}$).

Finally, in order to show that $j_{-1}[-](h \cdot \bar{f}) = f\Psi_0$ note first that, for any $(u, v) \in \mathfrak{L}(\mathbb{R})$, if $1 \notin \langle u, v \rangle$, then

$$\begin{aligned} j_{-1}[-]h(u, v) &\supseteq j_{-1}[-]h((1, -) \vee (-, 1)) = j_{-1}[-](h(1, -) \vee h(-, 1)) \\ &\supseteq j_{-1}[-](h(1, -) \cap S) = j_{-1}[-](S) = S, \end{aligned}$$

otherwise,

$$\begin{aligned} \mathbf{0} &= j_{-1}[-]h(1) = j_{-1}[-]h((- , 1) \vee (1, -) \vee (u, v)) \\ &= j_{-1}[-]h(u, v) \cap S = j_{-1}[-]h(u, v). \end{aligned}$$

Hence,

$$\begin{aligned} j_{-1}[-](h \cdot \bar{f})(r, s) &= \\ &= \bigcap \{j_{-1}[h(u, v)] \vee f\Psi_0(z, w) \mid 1 \in \langle u, v \rangle, \langle u, v \rangle \cdot \langle z, w \rangle \subseteq \langle r, s \rangle\} \\ &= \bigcap \{f\Psi_0(z, w) \mid 1 \in \langle u, v \rangle, \langle u, v \rangle \cdot \langle z, w \rangle \subseteq \langle r, s \rangle\} \\ &= \bigcap \{f\Psi_0(z, w) \mid r < z < w < s\} = f\Psi_0(r, s). \quad \blacksquare \end{aligned}$$

7. z -embedded sublocales

As is well known (see e.g. [2]), normality can be characterized in terms of C -embedded and C^* -embedded sublocales:

Theorem 7.1. *The following are equivalent for a locale L :*

- (i) L is normal.
- (ii) Any two disjoint closed sublocales are completely separated.
- (iii) Every closed sublocale is C -embedded.
- (iv) Every closed sublocale is C^* -embedded.

Our purpose in this final section is to study another class of sublocales, the z -embedded sublocales, inspired by the classical results in [1, Section 7]. In particular, we will present a characterization of normality, similar to the one above, in terms of z -embeddings.

Recall from [12] that a sublocale S of L is z -embedded if for every zero sublocale Z of S there is a zero sublocale W of L such that $W \cap S = Z$; in other words, S is z -embedded if for every cozero sublocale C of S there is a cozero sublocale D of L such that $D \cap S = C$.

Remark 7.2. The motivation for this notion is the following. Let $f: L \rightarrow M$ be a localic map and recall the adjunction $f[-] \dashv f_{-1}[-]$ from 1.3. Recall also (1.3.1). Since frame homomorphisms preserve cozero elements, $f_{-1}[-]$ restricts to maps

$$f_{-1}^z[-]: \mathbf{ZS}(M) \rightarrow \mathbf{ZS}(L) \quad \text{and} \quad f_{-1}^{\text{coz}}[-]: \mathbf{CozS}(M) \rightarrow \mathbf{CozS}(L).$$

The former (the *zero map*) is a σ -frame homomorphism and the latter (the *cozero map*) is a σ -coframe homomorphism. Clearly, $f_{-1}^z[-]$ is surjective if

and only if $f_{-1}^{\text{coz}}[-]$ is surjective. In this case, we say that f is a z -map. For the particular case of a localic embedding $j: S \hookrightarrow L$, j is a z -map iff S is z -embedded in L .

We do not pursue this approach in the present paper. The study of general z -maps is left to a subsequent article.

We start with a result that can be found in [6, Proposition 3.5] formulated in terms of frame quotients.

Proposition 7.3. *The following statements about a sublocale S of L are equivalent.*

- (i) S is z -embedded in L .
- (ii) For any two completely separated sublocales U, V of S there is a $C \in \text{CozS}(L)$, such that $U \subseteq C$ and $V \subseteq C^\#$.
- (iii) If U and V are completely separated sublocales of S , then they are S -separated; i.e., there exist cozero sublocales of L , say C_1 and C_2 , such that

$$U \subseteq C_1 \quad V \subseteq C_2 \quad \text{and} \quad C_1 \cap C_2 \cap S = \mathbf{O}.$$

Proof: (i) \Rightarrow (ii): Let U and V be completely separated in S . There exist cozero sublocales C_1 and C_2 in S such that $U \subseteq C_1$, $V \subseteq C_2$ and $C_1 \cap C_2 = \mathbf{O}$. Since S is z -embedded, there exists also a cozero sublocale C of L such that $C_1 = S \cap C$. Clearly, $U \subseteq C_1 \subseteq C$. Since $C_1 \cap V \subseteq C_1 \cap C_2 = \mathbf{O}$ we have

$$V \subseteq C_1^\# = (S \cap C)^\# = \mathbf{O} \vee C^\# = C^\#.$$

(ii) \Rightarrow (iii): By 3.1 it suffices to show statement (iii) for disjoint cozero sublocales of S . Take $C_1, C_2 \in \text{CozS}(S)$ such that $C_1 \cap C_2 = \mathbf{O}$. From the assumption there is a cozero sublocale C of L such that $C_1 \subseteq C$ and $C_2 \subseteq C^\#$. Now, $S \cap C$ is a cozero sublocale in S disjoint from C_2 . Applying the assumption again we get a cozero sublocale D of L such that $C_2 \subseteq D$ and $S \cap C \subseteq D^\#$. Hence, $C \cap S \cap D = \mathbf{O}$ as required.

(iii) \Rightarrow (i): Let C be a cozero sublocale of S . Then $C = f(0, -)$ for some $f \in \mathbf{C}(S)$ with $\mathbf{0} \leq f \leq \mathbf{1}$. Therefore, C is a closed sublocale $\mathbf{c}_S(a)$ with $a \in \text{Coz}(S)$. Furthermore, consider the cozero sublocales

$$B_n = f(-, \frac{1}{n}) = \mathbf{c}_S(b_n) \quad (b_n \in \text{Coz}(S), n = 1, 2, \dots).$$

Note that B_n and C are completely separated in S . Indeed,

$$B_n \cap C = f(-, \frac{1}{n}) \cap f(0, -) = f((-, \frac{1}{n}) \vee (0, -)) = f(\mathbf{1}) = \mathbf{O}.$$

By assumption, there are $\mathbf{c}_L(a_n)$ and $\mathbf{c}_L(d_n)$ with $a_n, d_n \in \text{Coz}(L)$ such that

$$\mathbf{c}_S(a) \subseteq \mathbf{c}_L(a_n), \quad \mathbf{c}_S(b_n) \subseteq \mathbf{c}_L(d_n) \quad \text{and} \quad \mathbf{c}_L(a_n) \cap \mathbf{c}_L(d_n) \cap S = \mathbf{0}$$

for every $n \in \mathbb{N}$. Consider now the cozero sublocale $\bigcap_{n=1}^{\infty} \mathbf{c}_L(a_n) = \mathbf{c}_L(v)$ where $v \in \text{Coz}(L)$ and recall (1.2.4). Then

$$\mathbf{c}_L(v) \cap S = \mathbf{c}_S(j_S^*(v)) \quad \text{and} \quad \mathbf{c}_L(d_n) \cap S = \mathbf{c}_S(j_S^*(d_n)).$$

Clearly, $\mathbf{c}_S(a) \subseteq \mathbf{c}_S(j_S^*(v))$. In order to show that this is indeed an equality, note first that $\mathbf{c}_S(j_S^*(v)) \cap \mathbf{c}_S(j_S^*(d_n)) = \mathbf{0}$. Then

$$\mathbf{c}_S(j_S^*(v)) \subseteq \mathbf{c}_S(j_S^*(d_n))^{\#s} = \mathbf{o}_S(j_S^*(d_n))$$

and $f(-, \frac{1}{n}) = \mathbf{c}_S(b_n) \subseteq \mathbf{c}_S(j_S^*(d_n))$. Therefore, taking supplements in S ,

$$\mathbf{c}_S(j_S^*(v)) \subseteq \mathbf{o}_S(b_n) \subseteq f(-, \frac{1}{n})^{\#s} \subseteq f((-, \frac{1}{n})^*) \subseteq f(\frac{1}{n}, -)$$

for $n = 1, 2, \dots$. Hence,

$$\mathbf{c}_L(v) \cap S = \mathbf{c}_S(j_S^*(v)) \subseteq \bigcap_{n=1}^{\infty} f(\frac{1}{n}, -) = f\left(\bigvee_{n=1}^{\infty} (\frac{1}{n}, -)\right) = f(0, -) = \mathbf{c}_S(a). \quad \blacksquare$$

Clearly, if T is a sublocale of S , z -embedded in L , then it is z -embedded in S . It is also easy to see that any C^* -embedded sublocale in L is z -embedded in L . Indeed, if $C \in \text{CozS}(S)$ then $C = f(0, -)$ for some $f \in \mathbf{C}^*(S)$ and $\mathbf{0} \leq f \leq \mathbf{1}$; then there exists an $\bar{f} \in \mathbf{C}^*(L)$ such that $j_{-1}[-]\bar{f} = f$, which means that

$$S \cap \bar{f}(0, -) \cap \bar{f}(-, 0) = j_{-1}[\bar{f}((0, -) \vee (-, 0))] = f(0, -) = C$$

(and $\bar{f}(0, -) \cap \bar{f}(-, 0) = \bar{f}((-, 0) \vee (0, -)) \in \text{Coz} L$).

In conclusion,

$$C\text{-embedded} \Rightarrow C^*\text{-embedded} \Rightarrow z\text{-embedded}.$$

Next result is a consequence of our extension theorem (via Theorem 6.1).

Proposition 7.4. *A sublocale S of a locale L is C^* -embedded in L if and only if it is z -embedded in L and for any sublocale T of S and $C \in \text{CozS}(L)$, if T and $S \cap C$ are S -separated in L , then T and $S \cap C$ are completely separated.*

Proof: If S is a C^* -embedded sublocale of L then it is z -embedded. Consider a sublocale T of S and $C \in \text{CozS}(L)$ such that T and $S \cap C$ are S -separated in L . This means there are $C_1, C_2 \in \text{CozS}(L)$ such that

$$T \subseteq C_1, \quad S \cap C \subseteq C_2 \quad \text{and} \quad C_1 \cap C_2 \cap S = \mathbf{0}.$$

In particular, $T \subseteq C_1 \cap S$ and $S \cap C \subseteq C_2 \cap S$. Thus, T and $S \cap C$ are completely separated in S because $C_1 \cap S, C_2 \cap S \in \mathbf{CozS}(S)$. Finally, by Theorem 6.1, T and $S \cap C$ are completely separated in L .

Conversely, we will prove that S is C^* -embedded using the characterization in 6.1. Let T and V be completely separated in S . There exist $C_1, C_2 \in \mathbf{CozSub}(S)$ such that $T \subseteq C_1$, $V \subseteq C_2$ and $C_1 \cap C_2 = \mathbf{O}$. Since S is z -embedded, then $C_1 = S \cap U_1$ and $C_2 = S \cap U_2$ for some $U_1, U_2 \in \mathbf{CozS}(L)$. Thus, $S \cap U_1$ and $S \cap U_2$ are S -separated, and by assumption they must be completely separated in L . Hence, T and V are also completely separated in L . ■

Remark 7.5. This result is another good example of the advantages of sublocale language in terms of conciseness and clarity. Indeed, the result is stated in [6, Proposition 4.3] in terms of frame quotients; a closer inspection to assertions (2) and (3) reveals, when formulated in terms of sublocales, that they express precisely the same fact.

A further consequence of the extension theorem is the following:

Proposition 7.6. *A sublocale S of a locale L is C -embedded in L if and only if it is z -embedded in L and it is completely separated from every cozero sublocale disjoint from it.*

Proof: If S is C -embedded then it is z -embedded. The other conclusion follows from Theorem 6.2. Conversely, assume that S is z -embedded in L and it is completely separated from every cozero sublocale disjoint from it. By 6.2, it suffices to show that S is C^* -embedded. We will do this using Theorem 6.1. Consider completely separated sublocales T and M in S . They are S -separated by 7.3, meaning that there are $C_1, C_2 \in \mathbf{CozS}(L)$ such that

$$T \subseteq C_1, M \subseteq C_2 \quad \text{and} \quad C_1 \cap C_2 \cap S = \mathbf{O}.$$

Take $C = C_1 \cap C_2 \in \mathbf{CozS}(L)$. Then $C \cap S = \mathbf{O}$ and by assumption, there is a $D \in \mathbf{CozS}(L)$ such that $S \subseteq D$ and $C \cap D = \mathbf{O}$. Hence,

$$T \subseteq D \cap C_1, \quad M \subseteq D \cap C_2 \quad \text{and} \quad D \cap C_1 \cap C_2 = D \cap C = \mathbf{O},$$

which means that S and T are completely separated in L . ■

Recall that a sublocale S of L is G_δ -dense in L if $S \cap A \neq \mathbf{O}$ for every G_δ -sublocale A of L . The preceding result characterizes C -embedded sublocales in among z -embedded sublocales. Next proposition shows that the class of z -embedded G_δ -dense sublocales is an example of such C -embedded sublocales.

Proposition 7.7. *Any z -embedded G_δ -dense sublocale of L is C -embedded in L .*

Proof: Let S be a z -embedded G_δ -dense sublocale of L . Recall from [12, Corollary 5.6.1] that every cozero sublocale is a G_δ -sublocale. Hence there is no cozero sublocale disjoint from S and S is C -embedded by the previous proposition. \blacksquare

Lemma 7.8. *In a normal locale, every F_σ -sublocale is z -embedded.*

Proof: Let S be an F_σ -sublocale of L , say $S = \bigvee_{n=1}^{\infty} \mathbf{c}(a_n)$. Consider a cozero sublocale $C = g(0, -)$ in S for some $g \in \mathbf{C}^*(S)$ with $\mathbf{0} \leq g \leq \mathbf{1}$. Let $\mathbf{c}(b)$ be the closure of C in L . Furthermore, consider for each $n = 1, 2, \dots$

$$T_n = \mathbf{c}(a_n) \vee \mathbf{c}(b) = \mathbf{c}(a_n \wedge b)$$

and $g_n: \mathfrak{L}(\mathbb{R}) \rightarrow \mathbf{S}(T_n)^{op}$ defined by

$$g_n(p, -) = \begin{cases} \mathbf{0} & \text{if } p < 0 \\ \mathbf{c}(b) \vee (g(p, -) \cap \mathbf{c}(a_n)) & \text{if } 0 \leq p < 1 \\ \mathbf{c}(b) \vee \mathbf{c}(a_n) & \text{if } p \geq 1 \end{cases}$$

and

$$g_n(-, q) = \begin{cases} \mathbf{c}(b) \vee \mathbf{c}(a_n) & \text{if } q \leq 0 \\ g(-, q) \cap \mathbf{c}(a_n) & \text{if } 0 < q \leq 1 \\ \mathbf{0} & \text{if } q > 1. \end{cases}$$

Let us confirm that this defines indeed a frame homomorphism by checking that it turns relations (r1)-(r6) into identities in the frame $\mathbf{S}(T_n)^{op}$:

(r1) $g_n(p, -) \vee g_n(-, q) = \mathbf{c}(b) \vee \mathbf{c}(a_n)$ whenever $p \geq q$:

The only nontrivial case is when $0 \leq p < 1$ and $0 < q \leq 1$ where we have

$$\begin{aligned} g_n(p, -) \vee g_n(-, q) &= (\mathbf{c}(b) \vee (g(p, -) \cap \mathbf{c}(a_n))) \vee (g(-, q) \cap \mathbf{c}(a_n)) \\ &= \mathbf{c}(b) \vee (\mathbf{c}(a_n) \cap (g(p, -) \vee g(-, q))) = \mathbf{c}(b) \vee (\mathbf{c}(a_n) \cap S) \\ &= \mathbf{c}(b) \vee \mathbf{c}(a_n). \end{aligned}$$

(r2) $g_n(p, -) \cap g_n(-, q) = \mathbf{0}$ whenever $p < q$:

The only nontrivial case is when $0 \leq p < 1$ and $0 < q \leq 1$ and we have

$$\begin{aligned} g_n(p, -) \cap g_n(-, q) &= (\mathbf{c}(b) \vee (g(p, -) \cap \mathbf{c}(a_n))) \cap (g(-, q) \cap \mathbf{c}(a_n)) \\ &= (\mathbf{c}(b) \cap g(-, q) \cap \mathbf{c}(a_n)) \vee (g(p, -) \cap \mathbf{c}(a_n) \cap g(-, q)) \\ &= (\mathbf{c}(b) \cap g(-, q) \cap \mathbf{c}(a_n) \cap S) \vee \mathbf{0} \\ &= g(0, -) \cap g(-, q) \cap \mathbf{c}(a_n) = \mathbf{0}. \end{aligned}$$

(Note that we are using the fact that $\mathbf{c}(b)$ is the closure in L of $g(0, -)$; hence, $\mathbf{c}(b) \cap S = g(0, -)$.)

(r3) $\bigcap_{r>p} g_n(r, -) = g_n(p, -)$:

For the only nontrivial case, when $0 \leq p < 1$, we have

$$\begin{aligned} \bigcap_{r>p} g_n(r, -) &= \mathbf{c}(b) \vee \bigcap_{1>r>p} (g(r, -) \cap \mathbf{c}(a_n)) \\ &= \mathbf{c}(b) \vee (g(p, -) \cap \mathbf{c}(a_n)) = g_n(p, -). \end{aligned}$$

Note that the second equality holds since g is a frame homomorphism and $g \leq \mathbf{1}$.

(r4) $\bigcap_{s<q} g_n(-, s) = g_n(-, q)$:

For the only nontrivial case $0 < q \leq 1$ we have

$$\bigcap_{s<q} g_n(-, s) = \bigcap_{0<s<q} (g(-, s) \cap \mathbf{c}(a_n)) = g(-, q) \cap \mathbf{c}(a_n) = g_n(-, q)$$

(the second equality holds since g is a frame homomorphism and $\mathbf{0} \leq g$).

(r5) $\bigcap_{p \in \mathbb{Q}} g_n(p, -) = \mathbf{0}$ is clear.

(r6) $\bigcap_{q \in \mathbb{Q}} g_n(-, q) = \mathbf{0}$ is also obvious.

In order to see that g_n is continuous for every n it suffices to check that $g(p, -) \cap \mathbf{c}(a_n)$ and $g(-, q) \cap \mathbf{c}(a_n)$ are closed sublocales in T_n for every $0 \leq p < 1$ and $0 < q \leq 1$. Regarding the former, since g is continuous, $g(p, -)$ is closed in S and thus there is a $d \in L$ such that $\mathbf{c}(d) \cap S = g(p, -)$. Hence

$$\begin{aligned} (\mathbf{c}(d) \cap \mathbf{c}(a_n)) \cap (\mathbf{c}(a_n) \vee \mathbf{c}(b)) &= (\mathbf{c}(d) \cap \mathbf{c}(a_n)) \vee (\mathbf{c}(d) \cap \mathbf{c}(a_n) \cap \mathbf{c}(b)) \\ &= (g(p, -) \cap \mathbf{c}(a_n)) \vee (g(p, -) \cap \mathbf{c}(a_n) \cap g(0, -)) \\ &= (g(p, -) \cap \mathbf{c}(a_n)) \vee (\mathbf{c}(a_n) \cap g(0, -)) \\ &= \mathbf{c}(a_n) \cap (g(p, -) \vee g(0, -)) = g(p, -) \cap \mathbf{c}(a_n). \end{aligned}$$

Similarly, if $\mathbf{c}(d) \cap S = g(-, q)$ we have

$$\begin{aligned} (\mathbf{c}(d) \cap \mathbf{c}(a_n)) \cap (\mathbf{c}(a_n) \vee \mathbf{c}(b)) &= (\mathbf{c}(d) \cap \mathbf{c}(a_n)) \vee (\mathbf{c}(d) \cap \mathbf{c}(a_n) \cap \mathbf{c}(b)) \\ &= (g(-, q) \cap \mathbf{c}(a_n)) \vee (g(-, q) \cap \mathbf{c}(a_n) \cap g(0, -)) \\ &= (g(-, q) \cap \mathbf{c}(a_n)) \vee \mathbf{O} = g(-, q) \cap \mathbf{c}(a_n). \end{aligned}$$

By Theorem 7.1 we know that T_n is C -embedded in L . Consequently, there are $f_n \in \mathbf{C}(L)$ ($n = 1, 2, \dots$) such that $(j_n)_{-1}[-]f_n = g_n$ where j_n is the localic embedding of T_n in L . Take

$$F = \bigcap_{n=1}^{\infty} (f_n(0, -) \cap f_n(-, 0)) \in \mathbf{CozS}(L).$$

We claim that $F \cap S = C$. First note that

$$\begin{aligned} g_n(0, -) \cap g_n(-, 0) &= (\mathbf{c}(b) \vee (g(0, -) \cap \mathbf{c}(a_n))) \cap (\mathbf{c}(b) \vee \mathbf{c}(a_n)) \\ &= \mathbf{c}(b) \vee ((\mathbf{c}(b) \vee (g(0, -) \cap \mathbf{c}(a_n))) \cap \mathbf{c}(a_n)) \\ &= \mathbf{c}(b) \vee (\mathbf{c}(b) \cap \mathbf{c}(a_n)) \vee (\mathbf{c}(a_n) \cap g(0, -)) \\ &= \mathbf{c}(b) \vee (\mathbf{c}(a_n) \wedge g(0, -)) = (\mathbf{c}(b) \vee \mathbf{c}(a_n)) \cap (\mathbf{c}(b) \vee g(0, -)) \\ &= (\mathbf{c}(b) \vee \mathbf{c}(a_n)) \cap \mathbf{c}(b) = \mathbf{c}(b), \end{aligned}$$

hence $g(0, -) \subseteq \mathbf{c}(b) = g_n(0, -) \cap g_n(-, 0) \subseteq f_n(0, -) \cap f_n(-, 0)$ for every n . Therefore, $g(0, -) \subseteq F \cap S$. For the converse inclusion we have

$$\begin{aligned} F \cap S &= \bigvee F \cap \mathbf{c}(a_n) \subseteq \bigvee f_n(0, -) \cap f_n(-, 0) \cap \mathbf{c}(a_n) \\ &= \bigvee f_n(0, -) \cap f_n(-, 0) \cap \mathbf{c}(a_n) \cap T_n = \bigvee g_n(0, -) \cap g_n(-, 0) \cap \mathbf{c}(a_n) \\ &= \bigvee \mathbf{c}(b) \cap \mathbf{c}(a_n) = \mathbf{c}(b) \cap S = g(0, -) \end{aligned}$$

where the first and the last equalities hold because F and $\mathbf{c}(b)$ are closed sublocales in particular complemented sublocales ([14, VI.4.4.1]). We have shown that an arbitrary cozero sublocale of S is the intersection of S with a cozero sublocale in L . In conclusion, S is z -embedded in L . \blacksquare

Lemma 7.9. *If S is a sublocale of L with the property that whenever $S \subseteq \mathfrak{o}(a)$ there is a normal and z -embedded F such that $S \subseteq F \subseteq \mathfrak{o}(a)$, then S is z -embedded in L .*

Proof: Let $A = \mathbf{c}_S(b)$ (with $b \in \mathbf{Coz}(S)$) be a cozero sublocale of S . Then A is a G_δ -sublocale of S , that is, $A = \bigcap_{n=1}^{\infty} \mathfrak{o}_S(b_n)$ for some $b_n \in S$. Consider the open sublocales $\mathfrak{o}_L(b \vee b_n)$ for $n = 1, 2, \dots$. Since $\mathbf{c}_S(b) \subseteq \mathfrak{o}_S(b_n)$ we have

$$S \cap \mathbf{c}_L(b \vee b_n) = S \cap \mathbf{c}_L(b) \cap \mathbf{c}_L(b_n) = \mathbf{c}_S(b) \cap \mathbf{c}_S(b_n) = \mathbf{O}.$$

Hence $S \subseteq \mathfrak{o}_L(b \vee b_n)$. By assumption, there is for each n a normal and z -embedded sublocale T_n such that $S \subseteq T_n \subseteq \mathfrak{o}_L(b \vee b_n)$. Note that $\mathfrak{c}_L(b) \cap T_n$ and $\mathfrak{c}_L(b_n) \cap T_n$ are disjoint; indeed

$$\mathfrak{c}_L(b) \cap T_n \cap \mathfrak{c}_L(b_n) \subseteq \mathfrak{c}_L(b \vee b_n) \cap \mathfrak{o}_L(b \vee b_n) = \mathbf{O}.$$

Recall Remark 3.2. By the normality of T_n , $\mathfrak{c}_L(b) \cap T_n$ and $\mathfrak{c}_L(b_n) \cap T_n$ are then completely separated in T_n . Consequently, there is a $C_n \in \mathbf{CozS}(T_n)$ such that

$$T_n \cap \mathfrak{c}_L(b) \subseteq C_n \quad \text{and} \quad T_n \cap \mathfrak{c}_L(b_n) \cap C_n = \mathbf{O}.$$

On the other hand, by z -embeddedness of T_n there is a $C'_n \in \mathbf{CozS}(L)$ such that $T_n \cap C'_n = C_n$. Finally, consider the cozero sublocale $\bigcap_{n=1}^{\infty} C'_n$. We claim $A = S \cap \bigcap_{n=1}^{\infty} C'_n$. The inclusion ' \subseteq ' is clear because

$$A \subseteq S \quad \text{and} \quad A \subseteq \mathfrak{c}_L(b) \cap T_n \subseteq C_n \subseteq C'_n$$

for every n . Conversely,

$$S \cap \bigcap_{n=1}^{\infty} C'_n = S \cap \bigcap_{n=1}^{\infty} (C'_n \cap T_n) = S \cap \bigcap_{n=1}^{\infty} C_n \stackrel{(*)}{\subseteq} S \cap \bigcap_{n=1}^{\infty} \mathfrak{o}_L(b_n) = \bigcap_{n=1}^{\infty} \mathfrak{o}_S(b_n) = A.$$

where $(*)$ holds because $T_n \cap \mathfrak{c}_L(b_n) \cap C_n = \mathbf{O}$ hence $C_n = T_n \cap C_n \subseteq \mathfrak{o}_L(b_n)$. ■

Finally, we say that a sublocale S of L is F_σ -generalized if for every open sublocale T of L with $S \subseteq T$, there exists an F_σ -sublocale F such that $S \subseteq F \subseteq T$. Obviously, any F_σ -sublocale is F_σ -generalized.

Theorem 7.10. *The following statements about a locale L are equivalent.*

- (i) L is normal.
- (ii) Every closed sublocale of L is z -embedded in L .
- (iii) Every F_σ -sublocale of L is z -embedded in L .
- (iv) Every generalized F_σ -sublocale of L is z -embedded in L .
- (v) For any closed sublocale F of L and any cozero sublocale C of L , $F \vee C$ is z -embedded in L .
- (vi) For any closed sublocale F of L and any cozero sublocale C of L such that $F \cap C = \mathbf{O}$, $F \vee C$ is z -embedded in L .

Proof: (i) \Rightarrow (iv): Let S be a generalized F_σ -sublocale of L . If $S \subseteq \mathfrak{o}(a)$ then there is an F_σ -sublocale F such that $S \subseteq F \subseteq \mathfrak{o}(a)$. By Lemma 7.8, F is z -embedded. On the other hand, being an F_σ -sublocale of a normal locale, F is also normal (see [10, Proposition 6.4]). Hence, the conclusion follows from Lemma 7.9.

(iv) \Rightarrow (iii) \Rightarrow (ii) is obvious.

(ii) \Rightarrow (i): To prove that L is normal it suffices to show that every closed sublocale is C -embedded (by Theorem 7.1). Let $\mathfrak{c}(s)$ be a closed sublocale of L ; by assumption, it is z -embedded. We will use Proposition 7.6 to show that it is also C -embedded. Let $\mathfrak{c}(a) \in \mathbf{CozS}(L)$ such that $\mathfrak{c}(s) \cap \mathfrak{c}(a) = \mathbf{O}$ and consider the closed sublocale $T = \mathfrak{c}(a) \vee \mathfrak{c}(s)$. We claim $\mathfrak{c}(s)$ is a cozero sublocale of T . Indeed,

$$\mathfrak{o}(a) \cap T = \mathfrak{o}(a) \cap (\mathfrak{c}(a) \vee \mathfrak{c}(s)) = \mathfrak{c}(s) \quad \text{and} \quad \mathfrak{c}(s) \cap T = \mathfrak{c}(s) \cap (\mathfrak{c}(a) \vee \mathfrak{c}(s)) = \mathfrak{c}(s)$$

assert that $\mathfrak{c}(s)$ is a clopen sublocale of T ; therefore (recall 1.7) it is both a zero and a cozero sublocale of T . By assumption, T is z -embedded because it is closed; thus, $\mathfrak{c}(s) = T \cap C$ for some $C \in \mathbf{CozS}(L)$. Hence,

$$\mathfrak{c}(s) \subseteq C \quad \text{and} \quad C \cap \mathfrak{c}(a) = C \cap (\mathfrak{c}(a) \cap T) = \mathfrak{c}(s) \cap \mathfrak{c}(a) = \mathbf{O}$$

which means that $\mathfrak{c}(a)$ and $\mathfrak{c}(s)$ are completely separated, as required.

(iii) \Rightarrow (v) is clear (since $F \vee C$ is an F_σ -sublocale), (v) \Rightarrow (vi) is trivial and (vi) \Rightarrow (ii) follows by taking $C = \mathbf{O}$. \blacksquare

References

- [1] R. Alò and H. Shapiro, *Normal Topological Spaces*, Cambridge Tracts in Mathematics, vol. 65, Cambridge University Press, 2008 (Reprint of the 1974 original edition).
- [2] R. N. Ball and J. Walters-Wayland, C - and C^* -quotients in pointfree topology, *Dissertationes Mathematicae (Rozprawy Mat.)* 412 (2002) 1-62.
- [3] B. Banaschewski, *The real numbers in pointfree topology*, Textos de Matemática, vol. 12, Universidade de Coimbra, 1997.
- [4] B. Banaschewski and C. Gilmour, Cozero bases of frames, *J. Pure Appl. Algebra* 157 (2001) 1-22.
- [5] R. L. Blair, Extensions of Lebesgue sets and of real valued functions, *Czechoslovak Math. J.* 31 (1981) 63-74.
- [6] T. Dube and J. Walters-Wayland, Coz-onto frame maps and some applications, *Appl. Categ. Structures* 15 (2007) 119-133.
- [7] M. J. Ferreira, J. Picado and S. Pinto, Remainders in pointfree topology, *Topology Appl.* 245 (2018) 21-45.
- [8] J. Gutiérrez García and T. Kubiak, General insertion and extension theorems for localic real functions, *J. Pure Appl. Algebra* 215 (2011) 1198-1204.
- [9] J. Gutiérrez García, T. Kubiak and J. Picado, Localic real functions: a general setting, *J. Pure Appl. Algebra* 213 (2009) 1064-1074.
- [10] J. Gutiérrez García, I. Mozo Carollo, J. Picado and J. Walters-Wayland, Hedgehog frames and a cardinal extension of normality, *J. Pure Appl. Algebra* 223 (2019) 2345-2370.
- [11] J. Gutiérrez García and J. Picado, Rings of real functions in pointfree topology, *Topology Appl.* 158 (2011) 2264-2278.

- [12] J. Gutiérrez García, J. Picado and A. Pultr, Notes on point-free real functions and sublocales, in: *Categorical Methods in Algebra and Topology*, Textos de Matemática, DMUC, vol. 46, pp. 167-200, 2014.
- [13] S. Mrówka, On some approximation theorems, *Nieuw Archief voor Wiskunde* (3) 16 (1968) 94-111.
- [14] J. Picado and A. Pultr, *Frames and locales: Topology without points*, Frontiers in Mathematics, vol. 28, Springer, Basel, 2012.
- [15] T. Plewe, Sublocale lattices, *J. Pure Appl. Algebra* 168 (2002) 309-326.

ANA BELÉN AVILEZ

UNIVERSITY OF COIMBRA, CMUC, DEPARTMENT OF MATHEMATICS, 3001-501 COIMBRA, PORTUGAL
E-mail address: `anabelenag@ciencias.unam.mx`

JORGE PICADO

UNIVERSITY OF COIMBRA, CMUC, DEPARTMENT OF MATHEMATICS, 3001-501 COIMBRA, PORTUGAL
E-mail address: `picado@mat.uc.pt`