

ON EMBEDDINGS OF 2-STRING TANGLES INTO THE UNKNOT, THE UNLINK AND SPLIT LINKS

JOÃO M. NOGUEIRA AND ANTÓNIO SALGUEIRO

ABSTRACT: In this paper we study further when tangles embed into the unknot, the unlink or a split link. In particular, we study obstructions to these properties through geometric characterizations, tangle sums and colorings. As an application we determine when each prime 2-string tangle with up to seven crossings embeds into the unknot, the unlink or a split link.

KEYWORDS: Embeddings of tangles; unknot; unlink; split link; colorings.
AMS SUBJECT CLASSIFICATION (2010): 57M25, 57N10.

1. Introduction

A *tangle* \mathcal{T} is a pair (B, σ) formed by a ball B and a collection of properly embedded disjoint arcs σ in B . If σ has n components we say that \mathcal{T} is a n -string tangle. Two tangles $\mathcal{T}, \mathcal{T}'$ are *equivalent*, denoted by $\mathcal{T} \approx \mathcal{T}'$, if there is an isotopy of B sending \mathcal{T} to \mathcal{T}' , and *strongly equivalent*, denoted by $\mathcal{T} = \mathcal{T}'$, if this isotopy fixes ∂B .

Let K be a link in S^3 , and B a ball in S^3 with exterior B' . If $\mathcal{T} = (B, B \cap K)$ and $\mathcal{T}' = (B', B' \cap K)$ are tangles, then we say that $\mathcal{T} \cup \mathcal{T}'$ is a *tangle decomposition* of K and that K is a *closure* of \mathcal{T} (and of \mathcal{T}'). In case there is a tangle decomposition of K with \mathcal{T} one of the tangle components, we also say that \mathcal{T} embeds into the pair (S^3, K) , or, for abbreviation, that it embeds into K . Similarly, let $\mathcal{T}_1 = (B_1, \sigma_1)$ and $\mathcal{T}_2 = (B_2, \sigma_2)$ be tangles such that $B_1 \cap B_2$ is a disk and $\mathcal{T} = (B_1 \cup B_2, \sigma_1 \cup \sigma_2)$ is a tangle. Then we say that $\mathcal{T}_1 \cup \mathcal{T}_2$ is a *tangle decomposition* of \mathcal{T} .

A fundamental question in knot theory is determining whether a knot or link is actually the unknot (resp., an unlink or a split link). In this paper, we continue the study of when a tangle embeds or not into the unknot, the unlink or a split link. If a tangle \mathcal{T} embeds into the unknot, the unlink, or a split link, we say that \mathcal{T} is *unknottable*, *unlinkable* or *splittable*. We refer

Received December 11, 2020.

This work was partially supported by the Centre for Mathematics of the University of Coimbra - UIDB/00324/2020, funded by the Portuguese Government through FCT/MCTES.

to a tangle \mathcal{U} such that $\mathcal{T} \cup \mathcal{U}$ is a tangle decomposition of the unknot, the unlink or of a split link, respectively, as an *unknotting*, *unlinking* or *splitting closure tangle* of \mathcal{T} .

A *projection* of a tangle \mathcal{T} is the image $p(\mathcal{T})$ of the tangle by an orthogonal projection p to a plane such that the preimage of each point of $p(\mathcal{T})$ has at most two points, and there are a finite number of double points, which are called the *crossings* of the projection. (See Figure 2 a) for an example of a tangle diagram.) A projection always exists in the piecewise linear category. If these crossings are decorated with broken lines to show the overcrosses and undercrosses, then we get a *diagram* of \mathcal{T} . A tangle diagram \mathcal{D} is called *unknottable*, *unlinkable* or *splittable* if there is a diagram of the unknot, the unlink or a split link, respectively, that contains \mathcal{D} .

The study of embeddings of tangles into the unknot, the unlink or split links has been considered before in several papers. In [10], Krebes shows that the greatest common divisor of the determinant of the numerator and denominator closure of a 2-string tangle divides the determinant of any closure link of the tangle, and is able to show several tangles not to be unknottable. This work uses a combinatorial interpretation of the determinant in terms of link diagrams and Kauffman brackets. In [18], Silver and Williams gave a new shorter proof of this result. In [11], this approach is generalised to n -string tangles with certain bracket-derived invariants of link diagrams. In [17], Silver and Williams extend Krebes' result using Fox colorings. More recently, Kauffman and Lopes [9] use colorings of knot diagrams with involutory quandles to study how to obtain tangles that are not unknottable. In [15], Ruberman extends the work of Krebes following a homological interpretation of the determinant of a link as the order of the first homology of the 2-fold branched cover of S^3 over the link; and also introduces an obstruction for a 2-string tangle to be unlinkable, through an application of work of Cochran and Ruberman [3] on tangle invariants from higher order linking numbers. In [14], Przytycki, Silver and Williams extend the work of Krebes and of Ruberman to n -string tangles and use the Jones polynomial instead of just the determinant. With this work, they are able to show an example of a 2-string tangle from [10] that is not unknottable and unlinkable and which is not possible to prove using the determinant result of [10]. In the present paper, we are also able to show that this tangle is not unknottable

and unlinkable with a different approach.

For 2-string tangles in particular, more can be said from the literature. As observed in section 4, a 2-string tangle is unknottable (resp., unlinkable) if and only if the unknotting (resp., unlinking) closure tangle is a rational tangle, and it is splittable if and only if it has a rational tangle as a splitting closure tangle. Under these circumstances, with respect to strong equivalence, the unknotting closure tangle of an unknottable 2-string essential tangle is unique [1, 2], and the unlinking (resp. rational splitting) closure tangle of an unlinkable (resp., splittable) 2-string tangle is unique [5]. Furthermore, in case the 2-string tangle is essential, it cannot be unknottable and splittable (or unlinkable) simultaneously [16]. New proofs of these results were also obtained by Taylor in [19]. For the case of a rational tangle it is possible to determine exactly which rational tangles are its unknotting closure tangles, as seen in the work of Ernst and Sumners [4] and of Kauffman and Lambropoulou in [8].

This paper is organized as follows. In Section 2, we observe some fundamental basic properties on embeddings of tangles into the unknot, the unlink and a split link. We also observe that Conjecture 3.1 of [9] by Kauffman and Lopes is false. In Section 3, we give a geometric characterization for a 2-string tangle to be unknottable, unlinkable or splittable. In Section 4, we study the behavior of unknottability, unlinkability and unsplittability under the sum of tangles. In particular, we determine when a Montesinos tangle is unknottable, unlinkable or splittable. In Section 5, we study further when the coloring invariants are an obstruction for a tangle being unknottable. We also apply these invariants on the study of unlinkable tangles for the first time. In particular, we determine the unlinking closure tangle candidate for any 2-string tangle. In Section 6, we determine which tangles of the table classifying all 2-string tangles up to 7 crossings (table 1 in [6]) are unknottable, unlinkable or unsplittable. We refer to this table throughout the paper.

2. Basic properties

In this section, we observe basic properties of unknottable, unlinkable and splittable tangles.

Theorem 2.1. *Let \mathcal{T} be a tangle. The following are equivalent:*

- (a) for every tangle \mathcal{T}' (strongly) equivalent to \mathcal{T} , every diagram \mathcal{D}' of \mathcal{T}' is unknottable;
- (b) there is a tangle \mathcal{T}' (strongly) equivalent to \mathcal{T} that has a unknottable diagram;
- (c) \mathcal{T} is unknottable.

A similar result holds for unlinkable/splittable tangles.

Proof: The implication (a) \Rightarrow (b) is immediate, by considering any diagram of \mathcal{T} .

To prove (b) \Rightarrow (c), consider an unknottable diagram \mathcal{D}' of \mathcal{T}' . Then there is a diagram D of the unknot K that contains D' . Consider an isotopy λ of B sending \mathcal{T}' to \mathcal{T} . Extend λ to S^3 by defining it on the exterior B' of B as the conjugate of λ by a symmetry interchanging B and B' . Then $\lambda(K)$ is the unknot with a decomposition $\mathcal{T} \cup \mathcal{U}$. Therefore, \mathcal{T} is unknottable.

Finally, to prove (c) \Rightarrow (a), suppose that \mathcal{T} is unknottable. Then there is a tangle \mathcal{U} such that $\mathcal{T} \cup \mathcal{U}$ is a tangle decomposition of the unknot. Consider an isotopy λ of B sending \mathcal{T} to \mathcal{T}' (and its extension to S^3) and a diagram \mathcal{D}' of \mathcal{T}' . Then $\lambda(\mathcal{T}) \cup \lambda(\mathcal{U})$ is a tangle decomposition of the unknot that has a diagram containing \mathcal{D}' . Therefore \mathcal{D}' is unknottable. ■

We say that the tangle $\mathcal{T} = \mathcal{T}_1 \cup \mathcal{T}_2$ is the *union* of the tangles $\mathcal{T}_1 = (B_1, \sigma_1)$ and $\mathcal{T}_2 = (B_2, \sigma_2)$ if $B_1 \cap B_2$ is a disk disjoint from $\sigma_1 \cup \sigma_2$. If a tangle $\mathcal{T}_1 = (B_1, \sigma_1)$ is embedded in a tangle $\mathcal{T} = (B, \sigma)$, we say that \mathcal{T}_1 is a *subtangle* of \mathcal{T} .

Theorem 2.2. *Let \mathcal{T}_1 be a subtangle of \mathcal{T} .*

- (a) *If \mathcal{T} is unknottable, then \mathcal{T}_1 is unknottable.*
- (b) *If \mathcal{T} is unlinkable, then \mathcal{T}_1 is unlinkable if it intersects more than one string of \mathcal{T} ; otherwise \mathcal{T}_1 is unknottable.*
- (c) *If \mathcal{T} is splittable and \mathcal{T}_1 intersects two strings of \mathcal{T} separated by the splitting decomposition, then \mathcal{T}_1 is splittable.*

Proof: If $\mathcal{T} \cup \mathcal{T}'$ is a tangle decomposition of a link K , then $\mathcal{T}_1 \cup (\mathcal{T}' \cup (\mathcal{T} - \mathcal{T}_1))$ is also a tangle decomposition of K .

If K is an unknot, then \mathcal{T}_1 is unknottable; if K is an unlink, then \mathcal{T}_1 is unlinkable if it intersects more than one string of \mathcal{T} and unknottable otherwise; if K is a split link, then \mathcal{T}_1 is splittable if it intersects two strings of \mathcal{T} separated by the splitting decomposition. ■

As an immediate consequence of this theorem, we have the following corollary.

Corollary 2.2.1. *Let $\mathcal{T}_1 \cup \mathcal{T}_2$ be a tangle decomposition of \mathcal{T} .*

- (a) *If \mathcal{T} is unknottable, then \mathcal{T}_1 and \mathcal{T}_2 are unknottable.*
- (b) *If \mathcal{T} is unlinkable, then \mathcal{T}_1 and \mathcal{T}_2 are unlinkable or unknottable. If the tangle decomposition is not an union, then at least one of \mathcal{T}_1 or \mathcal{T}_2 is unlinkable.*
- (c) *If \mathcal{T} is splittable and the tangle decomposition is not an union, then at least one of \mathcal{T}_1 or \mathcal{T}_2 is splittable.*

For instance, since $7_{16} \approx \mathcal{T} * [-2]$, where $\mathcal{T} \approx 6_2$, and the tangle 6_2 is not unknottable nor unlinkable (see Section 6), then 7_{16} is also not unknottable nor unlinkable (as both strings are unknotted, from Proposition 4.1, it is also not splittable).

The reciprocal of this corollary is not true. For instance, 6_2 is not unknottable, unlinkable or splittable, as verified in Section 6, but it has a decomposition into two trivial 2-string tangles. However, in the case where the decomposition of tangles is an union, the reciprocal is also true, as stated in the following theorem.

Theorem 2.3. *Let \mathcal{T} be the union of the tangles \mathcal{T}_1 and \mathcal{T}_2 . Then*

- (a) *\mathcal{T} is unknottable if and only if \mathcal{T}_1 and \mathcal{T}_2 are unknottable.*
- (b) *\mathcal{T} is unlinkable if and only if \mathcal{T}_1 and \mathcal{T}_2 are unlinkable or unknottable.*
- (c) *\mathcal{T} is splittable.*

Proof: (a) Suppose that \mathcal{T} is unknottable. By Corollary 2.2.1a), \mathcal{T}_1 and \mathcal{T}_2 are unknottable. Now suppose that \mathcal{T}_1 and \mathcal{T}_2 are unknottable. Then there exist tangles $\mathcal{T}'_1, \mathcal{T}'_2$ such that $\mathcal{T}_1 \cup \mathcal{T}'_1$ and $\mathcal{T}_2 \cup \mathcal{T}'_2$ are tangle decompositions of the unknot. By adding a single crossing between \mathcal{T}'_1 and \mathcal{T}'_2 , as in Figure 1, we obtain an unknot K containing \mathcal{T} . Therefore \mathcal{T} is unknottable.

(b) Suppose that \mathcal{T} is unlinkable. By Corollary 2.2.1 (b), \mathcal{T}_1 and \mathcal{T}_2 are unlinkable or unknottable. Now suppose that \mathcal{T}_1 and \mathcal{T}_2 are unlinkable or unknottable. Then there exist tangles $\mathcal{T}'_1, \mathcal{T}'_2$ such that $\mathcal{T}_1 \cup \mathcal{T}'_1$ and $\mathcal{T}_2 \cup \mathcal{T}'_2$ are tangle decompositions of the unknot or an unlink. Hence, the union of the tangles $\mathcal{T}'_1, \mathcal{T}'_2$ and \mathcal{T} is an unlink. Therefore \mathcal{T} is unlinkable.




FIGURE 1. The union of unknottable tangles is unknottable.

(c) From the decomposition of \mathcal{T} defined by \mathcal{T}_1 and \mathcal{T}_2 we obtain a split link by closing \mathcal{T} appropriately on each side of the decomposition. ■

Corollary 2.3.1. *Any n -string tangle equivalent to the trivial n -string tangle is unknottable and unlinkable.*

By this Corollary, any rational tangle is unknottable. (See also [4, 8].) For 2-string tangles the reciprocal of this corollary is true, as a 2-string tangle that is unknottable and splittable is trivial [5].

But in general, the reciprocal of this corollary is not true. In fact, consider the 3-string tangle \mathcal{T} defined by adding a parallel string to one of the strings of the 2-string tangle 6_3 . Then \mathcal{T} is unlinkable (it embeds into the three components unlink) and unknottable, but it is not equivalent to a trivial tangle.

We also observe that Conjecture 3.1 of [9] is false. In fact, the tangle \mathcal{T} of Figure 2a) is essential [13], but it is unknottable, as can be seen in Figure 2b), which shows a diagram of a trivial knot containing a diagram of \mathcal{T} . Moreover, \mathcal{T} is irreducible in the sense of [9], because \mathcal{T} has minimum crossing number [13] and Figure 2c) shows that its numerator closure is the nontrivial knot 5_1 .

3. Geometric characterization for 2-string tangles

In this section we characterize geometrically unknottable, unlinkable and splittable 2-string tangles.

Definition 1. Let $\mathcal{T} = (B, s_1 \cup s_2)$ be a 2-string tangle.

We say that s_1 and s_2 are *quasi-parallel* if there is a (punctured) disk P embedded in B such that s_1 and s_2 are in the same component of ∂P and




FIGURE 2. The tangle \mathcal{T} is irreducible and unknottable.

$\partial P - (s_1 \cup s_2) \subset \partial B$.

We say that s_1 and s_2 are *quasi-trivial* if there are two disjoint (punctured) disks P_1 and P_2 embedded in B co-bounded, respectively, by s_1 and s_2 .

We say that \mathcal{T} is *quasi-inessential* if there is a (punctured) disk D properly embedded in B , disjoint from s_1 and s_2 , such that each boundary component of D separates the ends of s_1 from the ends of s_2 . (See Figure 3.)




FIGURE 3. Quasi-parallel, quasi-trivial and quasi-inessential strings.

Lemma 3.1. *Let $\mathcal{T} = (B, s_1 \cup s_2)$ be a 2-string tangle, with s_1 and s_2 quasi-parallel. If \mathcal{T} is inessential, then it is trivial.*

Proof: Since s_1 and s_2 quasi-parallel, there exists a (punctured) disk P embedded in B such that s_1 and s_2 are in the same component b of ∂P and $\partial P - (s_1 \cup s_2) \subset \partial B$. Assume that P has the minimal possible number of punctures.

Since \mathcal{T} is inessential, there exists a disk D properly embedded in B separating the strings s_1 and s_2 . Assume that the number of components of $D \cap P$, denoted by $|D \cap P|$, is minimal, among all possible disks D . Let a and a' be the arcs defined by $b - (s_1 \cup s_2)$ and α be an outermost arc of $D \cap P$ in D .

Suppose that α is non-separating in P . Then, by compressing P along α and the corresponding outermost disk in D , we reduce the number of

punctures of P , contradicting its minimality. Otherwise, suppose that α is separating in P . If the ends of α are in the same component of $\partial P - (s_1 \cup s_2)$, by compressing P along α and the corresponding outermost disk in D , we reduce the number of punctures of P or reduce $|D \cap P|$, contradicting their minimality. Hence, α is separating and has ends in distinct components of $\partial P - (s_1 \cup s_2)$. As α is separating, it has to have ends in the same component of ∂P . Hence, α has ends in b , more precisely, one end in a and the other end in a' . Then, α cuts P into two disjoint embedded possibly punctured disks, one co-bounded by s_1 and the other by s_2 . Therefore, s_1 and s_2 are unknotted, which, for being inessential, implies that \mathcal{T} is trivial. \blacksquare

Theorem 3.2. *A 2-string tangle is unknottable if and only if its strings are quasi-parallel.*

Proof: Let \mathcal{T} be a 2-string tangle. Suppose that \mathcal{T} is inessential. Then \mathcal{T} is the union of two 1-string tangles \mathcal{T}_1 and \mathcal{T}_2 . By Theorem 2.3, \mathcal{T} is unknottable if and only if the tangles \mathcal{T}_1 and \mathcal{T}_2 are unknottable, that is, both strings are unknotted. By Lemma 3.1, this is equivalent to the strings being quasi-parallel.

Now consider the case when $\mathcal{T} = (B, s_1 \cup s_2)$ is essential. Suppose first that \mathcal{T} is unknottable and let $\mathcal{U} = (B', u_1 \cup u_2)$ be its unknotting closure tangle, with $\mathcal{T} \cup \mathcal{U}$ a tangle decomposition of the unknot K . Denote the sphere $B \cap B'$ by S . Since K is trivial, it bounds an embedded disk D in S^3 . Consider D such that the number of components of $D \cap S$, denoted by $|D \cap S|$, is minimal.

Let α be one of those components. If α is a closed curve, then it is essential in $S - (s_1 \cup s_2)$, otherwise α bounds a disk in $S - (s_1 \cup s_2)$ and we can reduce $|D \cap S|$, contradicting its minimality. As \mathcal{T} is essential, the disk that α bounds in D is in B' . Therefore, by removing all these disks from D , we obtain a (punctured) disk D' embedded in B .

Now let α be an arc component of $D \cap S$. Since the only intersections of ∂D and S are the endpoints of s_1 and s_2 , the arc α connects these endpoints and there are only two such arcs α_1 and α_2 .

Suppose first that α_i connects the endpoints of s_i , for $i = 1, 2$. Let D_1, D_2 be disjoint disks cut from D by α_1, α_2 . If $D_i \cap S$ contains a closed curve α , as in Figure 4, then, as observed before, α doesn't bound a disk in $S - (s_1 \cup s_2)$, it bounds a disk in B' . As D_1 is disjoint from D_2 , the curve α separates ∂s_1 from ∂s_2 in S , and the disk α bounds in B' separates u_1 and u_2 . Then,

without loss of generality, ∂u_i is attached to ∂s_i , but this contradicts K being a knot. Therefore, D_i is disjoint from S , hence s_i is trivial in \mathcal{T} , which contradicts \mathcal{T} being essential.




FIGURE 4. If α_i connects the ends of s_i .

Suppose now that α_1 and α_2 both connect an endpoint of s_1 and an endpoint of s_2 . Let P be the (punctured) disk cut from D' by α_1 and α_2 , as in Figure 5. Then P is a disk embedded in B such that s_1 and s_2 are in the same component of ∂P and $\partial P - (s_1 \cup s_2) \subset \partial B$. Therefore, s_1 and s_2 are quasi-parallel.




FIGURE 5. If α_i connects one end of s_1 to an end of s_2 .

Reciprocally, suppose that s_1 and s_2 are quasi-parallel and P is a corresponding punctured disk. Let K be the component of ∂P containing $s_1 \cup s_2$ and u_1 and u_2 be the two arcs of $K - (s_1 \cup s_2)$. Isotope the interior of u_1 and u_2 into the exterior B' of B so that $\mathcal{T}' = (B', u_1 \cup u_2)$ is a tangle. Then $\mathcal{T} \cup \mathcal{T}'$ is a tangle decomposition of K . Since each component of $\partial P - K$ bounds a disk in $B' - (u_1 \cup u_2)$, then K bounds a disk in S^3 , hence it is trivial. Therefore \mathcal{T} is unknottable. ■

Theorem 3.3. *A 2-string tangle is unlinkable if and only if its strings are quasi-trivial.*

Proof: Let $\mathcal{T} = (B, s_1 \cup s_2)$ be a 2-string tangle.

Suppose the strings of \mathcal{T} are quasi-trivial and let P_1, P_2 be the corresponding disjoint punctured disks each co-bound. Let b_i be the boundary component of P_i that contains s_i , for $i = 1, 2$. Assuming B in S^3 , let B' be the complement of B and let s'_i be the arc obtained by pushing the interior of $b_i - s_i$ into the interior of B' , so that $\mathcal{T}' = (B', s'_1 \cup s'_2)$ is a 2-string tangle. Let L be the link in S^3 defined by $b_1 \cup b_2$. Then $\mathcal{T} \cup \mathcal{T}'$ is a tangle decomposition of L . Since each component of $\partial P_i - b_i$ bounds a disk in $B' - (s'_1 \cup s'_2)$, we have that b_1 and b_2 bound disjoint disks. That is, L is a trivial two component link and, therefore, \mathcal{T} is unlinkable.

Reciprocally, suppose that \mathcal{T} is unlinkable with unlinking closure tangle \mathcal{U} . Let b_1 and b_2 be the components of L and O_1, O_2 the corresponding disjoint disks they bound. We have that $b_i \cap B$ is s_i . Hence, $O_i \cap \partial B$ is a collection of simple closed curves and an arc α_i sharing the ends of s_i . Let D_i be the (punctured) disk cut by α_i from O_i that is co-bounded by s_i . Then, s_1 and s_2 are quasi-trivial strings. \blacksquare

Theorem 3.4. *A 2-string tangle is splittable if and only if it is quasi-inessential.*

Proof: Suppose that \mathcal{T} is quasi-inessential and let D be the punctured disk as in the definition. Let s'_i be an arc in ∂B connecting ∂s_i , disjoint from ∂D . Then the circle $s_1 \cup s'_1$ together with the circle $s_2 \cup s'_2$ define a 2-component link L . Let B' be the exterior of B in S^3 and \mathcal{T}' the 2-string tangle defined by $(B', s'_1 \cup s'_2)$ after pushing the interior of s'_i into the interior of B' . Each boundary component of ∂D bounds a disk in $B' - (s'_1 \cup s'_2)$ resulting on a sphere S that separates the components of L . Then, $\mathcal{T} \cup \mathcal{T}'$ is a tangle decomposition of a split link, that is, \mathcal{T} is splittable.

Reciprocally, suppose that \mathcal{T} is splittable with splitting closure tangle \mathcal{U} . In case there is a disk in B separating s_1 and s_2 , then this disk makes \mathcal{T} quasi-inessential, by definition. Hence, we can assume that \mathcal{T} is essential. Let S be the split sphere of L and consider the intersection of S with B . Suppose that $|S \cap \partial B|$ is minimal among all such spheres S . Note that it is necessarily non-empty as B intersects both sides of S . As \mathcal{T} is essential, the innermost curves of $S \cap \partial B$ in ∂B have the corresponding innermost disks in B' . Suppose that some curve of $S \cap \partial B$ is not innermost in S . Let c be innermost among these curves. Hence, c cuts from S a disk intersecting ∂B only in innermost curves. Let E be the corresponding punctured disk in B . The boundary components of E are parallel in $\partial B - \partial(s_1 \cup s_2)$ and




FIGURE 6. The ends of a tangle

bound disks in $B' - (s'_1 \cup s'_2)$. By capping off the boundary components of E with those disks, either the resulting sphere splits the link L , and in this case we have a contradiction with $|S \cap \partial B|$ being minimal, or it bounds a ball in the exterior of L , and by cutting S along c and pasting a disk it bounds in $B' - (s'_1 \cup s'_2)$ we obtain a split sphere of L that reduces $|S \cap \partial B|$, contradicting its minimality. Hence, all curves of $S \cap \partial B$ are innermost in S with corresponding innermost disks in B' separating s'_1 from s'_2 . That is, S intersects B at a single punctured disk D that separates s_1 from s_2 in B and whose boundary components separate ∂s_1 from ∂s_2 in ∂B . ■

4. Algebraic properties for 2-string tangles

In this section we consider 2-string tangles (B, σ) with the ends of σ being four fixed points in the boundary circle b of the diagram disk as in the intercardinal directions NW , SW , NE and SE , illustrated in Figure 6. We say that a disk in ∂B is a *west disk* (resp., *east disk*) if the disk intersects b in a single arc containing NW and SW (resp., NE and SE) but not any of the points NE and SE (resp., NW and SW). Similarly we define a *north disk* and a *south disk* in ∂B .

Definition 2. We define the *sum* of two 2-string tangles \mathcal{T} and \mathcal{T}' by identifying the west disk in ∂B to an east disk in $\partial B'$. The resulting tangle sum is denoted $\mathcal{T} + \mathcal{T}'$.

We define the *product* of \mathcal{T} and \mathcal{T}' by identifying a south disk of \mathcal{T} to a north disk in \mathcal{T}' . The resulting tangle product is denoted $\mathcal{T} * \mathcal{T}'$.

Throughout this section we study unknotability unlinkability and splittability of 2-string tangles up to strong equivalence and the behavior of these properties under tangle sum and product.

Definition 3. Let \mathcal{T} be a 2-string tangle. The *numerator closure* of \mathcal{T} is the link $N(\mathcal{T})$ obtained by the sum of \mathcal{T} and the $[0]$ tangle; the *denominator closure* of \mathcal{T} is the link $D(\mathcal{T})$ obtained by the sum of \mathcal{T} and the $[\infty]$ tangle.

Definition 4. Let \mathcal{T} be a 2-string tangle.

- (a) The tangle obtained from \mathcal{T} by a 180° rotation around an horizontal axis on the plane is denoted by $\bar{\mathcal{T}}$.
- (b) The tangle obtained from \mathcal{T} by a 90° rotation is denoted by \mathcal{T}^\perp .

Note that a unknotting (resp. unlinking or splitting) closure tangle of a 2-string tangle \mathcal{T} can be considered as a 2-string tangle \mathcal{U} such that $N(\mathcal{T} + \mathcal{U})$ is the unknot (resp., the unlink or a split link). We have that $N(\mathcal{T} + \mathcal{U})$ is equivalent to $D(\mathcal{T} * \bar{\mathcal{U}})$ and for \mathcal{U} a rational tangle, $\mathcal{U} = \bar{\mathcal{U}}$. Then, with \mathcal{U} a rational tangle, $D(\mathcal{T} * \mathcal{U})$ is also the unknot (resp., the unlink or a split link). From the next proposition, the unknotting or unlinking closure tangle of a 2-string tangle is a rational tangle, and we can assume that the splitting closure tangle of a 2-string tangle is a rational tangle.

Proposition 4.1. *Let \mathcal{T} be an essential 2-string tangle. If \mathcal{T} is unknottable (resp., unlinkable), then any unknotting (resp., unlinking) closure tangle is rational. If \mathcal{T} is splittable, then it has a rational splitting closure tangle.*

Proof: Suppose that \mathcal{T} is unknottable or unlinkable with corresponding closure tangle \mathcal{U} , that is, $N(\mathcal{T} + \mathcal{U})$ is the unknot or the unlink, respectively. Therefore, \mathcal{U} cannot have local knots and, as \mathcal{T} is essential, it cannot be essential. Hence, \mathcal{U} is trivial, that is, it is a rational tangle.

Suppose now that $\mathcal{T} = (B, \sigma)$ is splittable with splitting closure tangle \mathcal{U} , that is, $N(\mathcal{T} + \mathcal{U})$ is a split link. Let S be a split sphere for $N(\mathcal{T} + \mathcal{U})$. Consider an innermost curve c of $S \cap \partial B$. As S is a sphere, c bounds a disk D in S , disjoint from the other components of $S \cap \partial B$. As \mathcal{T} is essential, D cannot be in B . Hence D is in the exterior of B . That is, \mathcal{U} is inessential. Therefore, D separates the two strings of \mathcal{U} . Hence, in case both are unknotted, the tangle is trivial, and in case at least one is knotted, it can be replaced in the same ball separated by D in B' by an unknotted string. The resulting tangle \mathcal{V} is a rational tangle. As the strings of \mathcal{V} are disjoint from S , and the end points of the strings of \mathcal{U} and \mathcal{V} are the same, we have that $N(\mathcal{T} + \mathcal{V})$ is a split link. That is, \mathcal{V} is a rational splitting closure tangle of \mathcal{T} . ■

From Property P and double branched covers of unknottable 2-string tangles being knot exteriors in S^3 we have the following theorem, which is a consequence of work by Bleiler and Scharlemann in [1] and [2].

Theorem 4.2 (Bleiler and Scharlemann [1], [2]). *If an unknottable 2-string tangle is essential, then it has a unique unknotting closure tangle.*

Remark 1. If a 2-string tangle has two different unknotting closure tangles, then it is rational.

A similar result was obtained by Eudave-Muñoz in [5] for splittability, and hence unlinkability, of 2-string tangles.

Theorem 4.3 (Eudave-Muñoz [5]). *If a 2-string tangle is unlinkable (resp., splittable) then it has a unique unlinking (resp., rational splitting) closure tangle.*

From work of Scharlemann [16] we have that an essential 2-string tangle cannot be splittable and unknottable simultaneously.

Theorem 4.4 (Scharlemann [16]). *If a 2-string tangle is unknottable and splittable, then it is a rational tangle.*

In the next theorem, we determine when a Montesinos tangle is unknottable/unlinkable. This result extends, for instance, Theorem 5 of [8], which gives all rational unknotting closures tangles for a rational tangle. From Theorem 2.2 of [4] we also obtain, in particular, the rational unknotting/unlinking closure tangles of a rational tangle.

Theorem 4.5. *Let $\mathcal{T} = \left[\frac{p_1}{q_1} \right] + \left[\frac{p_2}{q_2} \right] + \cdots + \left[\frac{p_n}{q_n} \right]$, with $n \geq 2$ and $q_i > 1$.*

Then,

- (a) \mathcal{T} is unknottable if and only if $n = 2$ and $p_1q_2 + p_2q_1 \equiv \pm 1 \pmod{q_1q_2}$.
- (b) \mathcal{T} is unlinkable if and only if $n = 2$, $q_1 = q_2$ and $p_1 + p_2 \equiv 0 \pmod{q_1}$.
- (c) \mathcal{T} is splittable if and only if it is unlinkable.

Moreover, if \mathcal{T} is unknottable (resp. unlinkable, splittable), then its unknotting (resp. unlinking, rational splitting) closure tangle \mathcal{U} is integral and, if $-1 < \frac{p_1}{q_1}, \frac{p_2}{q_2} < 1$, then \mathcal{U} is either $[0]$, $[1]$ or $[-1]$.

Proof: Since \mathcal{T} is not rational, then, by Proposition 4.1, it has a rational unknotting/unlinking/splittable closure tangle $\mathcal{U} = \left[\frac{p}{q} \right]$. The double cover of

S^3 branched over the Montesinos knot $N(\mathcal{T} + \mathcal{U})$ is a Seifert fibred manifold M [12], with invariant

$$(0; p_1/q_1, \dots, p_n/q_n, p/q).$$

By the classification of Seifert fibred spaces, if there are at least 3 exceptional fibers, then M is irreducible and not S^3 . Therefore, if \mathcal{T} is unknottable/unlinkable/splittable, then $n = 2$ and $q = 1$.

(a) \mathcal{T} is unknottable if and only if M is S^3 , which is equivalent to

$$\frac{p_1}{q_1} + \frac{p_2}{q_2} + p = \pm \frac{1}{q_1 q_2},$$

or, similarly, $p_1 q_2 + p_2 q_1 + p q_1 q_2 = \pm 1$. Moreover, if $-1 < \frac{p_1}{q_1}, \frac{p_2}{q_2} < 1$, then clearly $-2 < p < 2$.

(b) \mathcal{T} is unlinkable if and only if M is $S^2 \times S^1$, which is equivalent to

$$\frac{p_1}{q_1} + \frac{p_2}{q_2} + p = 0,$$

or, similarly, $q_1 = q_2$ and $p_1 + p_2 + p q_1 = 0$. Moreover, if $-1 < \frac{p_1}{q_1}, \frac{p_2}{q_2} < 1$, then clearly $-2 < p < 2$.

(c) \mathcal{T} is splittable if and only if M is reducible, which happens exactly when M is $S^2 \times S^1$, since M is orientable and the fibration base is S^2 . ■

The following theorem describes the effect of adding a rational tangle to an unknottable/unlinkable/splittable 2-string tangle.

Theorem 4.6. *Let \mathcal{T} be a 2-string tangle with unknotting closure tangle $\left[\begin{smallmatrix} r \\ s \end{smallmatrix}\right]$. Then,*

- (a) $\mathcal{T} + \left[\begin{smallmatrix} p \\ q \end{smallmatrix}\right]$ is unknottable if and only if $q = 1$ or $q = s \wedge p \equiv r \pmod{q}$.
- (b) $\mathcal{T} * \left[\begin{smallmatrix} p \\ q \end{smallmatrix}\right]$ is unknottable if and only if $p = 1$ or $p = r \wedge q \equiv s \pmod{p}$.

A similar result holds for unlinkable/splittable tangles.

Proof: Since the tangle \mathcal{T} has unknotting closure tangle $\left[\begin{smallmatrix} r \\ s \end{smallmatrix}\right]$, then $N(\mathcal{T} + \left[\begin{smallmatrix} r \\ s \end{smallmatrix}\right]) = D(\mathcal{T} * \left[\begin{smallmatrix} r \\ s \end{smallmatrix}\right]) = \text{unknot}$.

- (a) Suppose that $\mathcal{T} + \left[\begin{smallmatrix} p \\ q \end{smallmatrix}\right]$ is unknottable, with unknotting closure tangle U .

Then $N\left(\mathcal{T} + \left[\frac{p}{q}\right] + U\right) = \text{unknot}$, hence $\left[\frac{p}{q}\right] + U = \left[\frac{r}{s}\right]$, by Theorem 4.2. Therefore, $U = [n]$ and $\left[\frac{p}{q}\right] = \left[\frac{r}{s}\right] + [-n] = \left[\frac{r-ns}{s}\right]$. Since $(p, q) = (r, s) = 1$, we conclude that $q = s$ and $p \equiv r \pmod{q}$.

(b) Since $D\left(\mathcal{T} * \left[\frac{r}{s}\right]\right) = \text{unknot}$, then $N\left(\mathcal{T}^\perp + \left[\frac{-s}{r}\right]\right) = \text{unknot}$.

The tangle $\mathcal{T} * \left[\frac{p}{q}\right]$ is unknottable if and only if $\mathcal{T}^\perp + \left[\frac{-q}{p}\right]$ is unknottable.

This is equivalent, by (a), to $p = r$ and $q \equiv s \pmod{p}$. ■

The following corollary characterizes unknottable, unlinkable and splittable tangles among algebraic tangles with three rational components.

Corollary 4.6.1. *The tangle $\left(\left[\frac{p_1}{q_1}\right] + \left[\frac{p_2}{q_2}\right]\right) * \left[\frac{p_3}{q_3}\right]$, with $q_1, q_2, p_3 \neq 1$, is unknottable if and only if $\left[\frac{p_1}{q_1}\right] + \left[\frac{p_2}{q_2}\right]$ is unknottable, with unknotting closure tangle $[p_3]$, and $q_3 \equiv 1 \pmod{p_3}$.*

A similar result holds for unlinkable/splittable tangles.

Proof: Suppose that $\left(\left[\frac{p_1}{q_1}\right] + \left[\frac{p_2}{q_2}\right]\right) * \left[\frac{p_3}{q_3}\right]$ is unknottable. By Theorem 2.2 Corollary 2.2.1, $\left[\frac{p_1}{q_1}\right] + \left[\frac{p_2}{q_2}\right]$ is also unknottable, and, by Theorem 4.5, its unknotting closure tangle is an integral tangle $[n]$. Then, by Theorem 4.6(b), $\left(\left[\frac{p_1}{q_1}\right] + \left[\frac{p_2}{q_2}\right]\right) * \left[\frac{p_3}{q_3}\right]$ is unknottable if and only if $n = p_3$ and $q_3 \equiv 1 \pmod{p_3}$. ■

Theorem 4.7. *Let $\mathcal{T} = (B, \sigma)$ be an essential 2-string tangle and D a disk intersecting σ and separating the ends of σ in two sets of two points. Suppose that the tangles separated from T by D are essential. Then \mathcal{T} is unknottable if and only if it has an unknotting closure which is a rational tangle with the same slope as ∂D .*

A similar result holds for unlinkable/splittable tangles.

Proof: If \mathcal{T} has any unknotting closure tangle, then \mathcal{T} is unknottable, by definition.

Suppose now that \mathcal{T} is unknottable, with unknotting closure tangle \mathcal{U} . Let $\mathcal{T}_i = (B_i, \sigma_i)$ denote the tangles separated by D from \mathcal{T} , for $i = 1, 2$. Consider the tangle $\mathcal{T}_2 + \mathcal{U}$. As \mathcal{T}_1 is essential we have that $\mathcal{T}_2 + \mathcal{U}$ is inessential. Let E be a disk separating the strings of $\mathcal{T}_2 + \mathcal{U}$. Note that ∂E is in ∂B_1 . As \mathcal{T}_2 is

essential, in case E intersects ∂B_2 we can reduce the number of components of $E \cap \partial B_2$ with an innermost curve/outermost arc argument, until E is disjoint of ∂B_2 . Then, ∂E is in $\partial B_1 - D$ and separates the points of $D \cap \sigma_1$ from the other two end points of σ_1 . Hence, the tangle \mathcal{U} , with respect to the fixed end points of \mathcal{T} , has the same slope as a rational tangle as ∂D in ∂B . ■

For a 2-string tangle (B, σ) , we say that a disk D properly embedded in B is a *meridian* (resp., *longitude*) *disk* if ∂D separates ∂B into a west and east disk (resp., into a north and south disk). In case D is a meridian or longitude disk of B with fixed boundary, we have the following corollary.

Corollary 4.7.1. *Let $\mathcal{T} = (B, \sigma)$ be an essential 2-string tangle and D a meridian (resp., longitude) disk of B . Suppose that the tangles separated from \mathcal{T} by D are essential. Then \mathcal{T} is unknottable if and only if $D(\mathcal{T})$ (resp., $N(\mathcal{T})$) is the unknot.*

A similar result holds for unlinkable/splittable tangles.

In particular, in the previous corollary, when D intersects σ at two points, we have the following corollary.

Corollary 4.7.2. *Let \mathcal{T}_1 and \mathcal{T}_2 be essential 2-string tangles. Then*

- (a) $\mathcal{T}_1 + \mathcal{T}_2$ is unknottable if and only if $D(\mathcal{T}_1 + \mathcal{T}_2) = \text{unknot}$.
- (b) $\mathcal{T}_1 * \mathcal{T}_2$ is unknottable if and only if $N(\mathcal{T}_1 * \mathcal{T}_2) = \text{unknot}$.

A similar result holds for unlinkable/splittable tangles.

Proof: The sum of the two 2-string tangles corresponds to a decomposition through a meridian disk in the resulting tangle. Hence, (a) is an immediate consequence of the theorem. The statement (b) is equivalent to (a) by a 90° rotation of the tangles. ■

5. Colorings

In this section we make further remarks on the coloring invariants of knots being an obstruction for a tangle to be unknottable. This observations follow in line with a remark by Silver in Krebs paper [10] on Fox colorings being able to detect when a 2-string tangle is not unknottable, and with the work of Kauffman and Lopes in [9] on involutory quandle colorings, which include Fox colorings, being able to detect when a 2-string tangle is not unknottable. Here we make observations that these statements cannot be extended to

oriented quandles. We also determine a necessary condition from coloring invariants for unlinkability and splittability of 2-string tangles.

Definition 5. A *quandle* is a set X with an operation \triangleright such that

- $\forall x \in X, x \triangleright x = x$;
- $\forall y, z \in X, \exists! x \in X : z = x \triangleright y$;
- $\forall x, y, z \in X, (x \triangleright y) \triangleright z = (x \triangleright z) \triangleright (y \triangleright z)$.

Definition 6. A *coloring* of an oriented knot or tangle by the quandle (X, \triangleright) is a labeling of its arcs by elements of X in such a way that at a crossing where the right underarc has label x , the overarc has label y , and the left underarc has label z , then $z = x \triangleright y$, as in Figure 7.




FIGURE 7. The consistency of the quandle operation and the labeling at each crossing.

We say that a quandle coloring of an oriented knot or tangle is *nontrivial* if it uses more than one color; we say that it is *trivial* otherwise. A tangle coloring such that all boundary arcs have the same color is called a *c-coloring*, and otherwise is called a *d-coloring*. We say also that an oriented knot is *polychromatic* if it has a nontrivial coloring and that an oriented tangle is *polychromatic* if it has a nontrivial *c-coloring*. Otherwise, the knot is said *monochromatic* and the tangle is said *monochromatic* if every *c-coloring* of the tangle is trivial. Since Reidemeister moves preserve the existence of a nontrivial coloring, then an oriented knot or tangle is polychromatic if and only if any equivalent oriented knot or tangle is polychromatic.

If (X, \triangleright) is a quandle, then (X, \triangleleft) , where $z \triangleleft y$ is the unique x such that $z = x \triangleright y$, is also a quandle and a coloring of an oriented knot or tangle by (X, \triangleright) determines a coloring of the knot or tangle with the opposite orientation by (X, \triangleleft) . However, if we change the orientation of some strings of a tangle while keeping the orientation of the remaining strings, it may happen that the original oriented tangle is polychromatic and the new oriented tangle is monochromatic. If the operations \triangleleft and \triangleright coincide, the quandle is called *involutory* or *unoriented*. An oriented tangle has a nontrivial coloring by an involutory quandle (X, \diamond) if and only if it has a nontrivial coloring by (X, \diamond) ,

for any orientation of its strings. The *dihedral quandle* R_n is the involutory quandle $(\mathbb{Z}/n\mathbb{Z}, \diamond)$, where $x \diamond y \equiv 2y - x \pmod{n}$ and n is a non-negative integer (for the sake of knot or tangle nontrivial colorings, we may consider only the case where n is zero or prime.)

Theorem 5.1. *If a tangle \mathcal{T} is unknottable, then \mathcal{T} is monochromatic for some orientation.*

Proof: Let \mathcal{U} be a unknotting closure tangle of \mathcal{T} . Suppose that \mathcal{T} is polychromatic for the orientation induced by the unknot $K = N(\mathcal{T} + \mathcal{U})$. Then, by assigning the boundary color of \mathcal{T} to all arcs of \mathcal{U} , we get a nontrivial coloring of K . Since the trivial diagram of the unknot is monochromatic, we obtain a contradiction. ■

The following corollary is also a remark by Silver in Krebs paper [10] on Fox colorings being able to detect when a 2-string tangle is not unknottable, and it also appears in the work of Kauffman and Lopes in [9] on involutory quandle colorings being able to detect when a 2-string tangle is not unknottable.

Corollary 5.1.1. *If a tangle \mathcal{T} is unknottable, then every c -coloring of \mathcal{T} by an involutory quandle is trivial.*

Proof: Suppose that \mathcal{T} has a nontrivial c -coloring by an involutory quandle. Then \mathcal{T} is polychromatic for all orientations, hence, by Theorem 5.1, \mathcal{T} is not unknottable. ■

By Corollary 5.1.1, if one can find a polychromatic coloring of a tangle by an involutory quandle, then this tangle is not unknottable. For instance, since the 2-string tangles 6_2 , 6_3 , 7_{13} , 7_{15} , 7_{16} , 7_{17} and 7_{18} have polychromatic colorings by dihedral quandles (see Figure 8 for one such coloring), they are not unknottable.




FIGURE 8. Polychromatic colorings of 6_2 , 6_3 , 7_{13} , 7_{15} , 7_{16} , 7_{17} and 7_{18} , respectively.

The reciprocal of Theorem 5.1 is not true. For instance, the tangle 6_4 is not unknottable (see Section 6), but it is monochromatic. To see this, suppose that there is a coloring of 6_4 by some quandle, with the four ends having the same color 0. Let the other arcs have colors 1, 2, 3, 4 as in Figure 9. Since the arc with the color 1 is in two crossings with a color 0 overarc on the same side, then the colors 2 and 3 coincide. It then follows that the coloring is trivial.




FIGURE 9. Any c -coloring of the tangle 6_4 is trivial.

Also, the conclusion of Theorem 5.1 cannot be extended to all orientations. For instance, the tangle 7_7 is unknottable (see Section 6), but it is polychromatic for one of the orientations. To see this, consider the quandle $\mathbb{Z}_2[t]/(t^2 + t + 1)$ whose multiplication table is

\triangleright	0	1	t	$t + 1$
0	0	$t + 1$	1	t
1	t	1	$t + 1$	0
t	$t + 1$	0	t	1
$t + 1$	1	t	0	$t + 1$

There is a polychromatic coloring of one orientation of 7_7 by this quandle (see Figure 10). The existence of this coloring implies that there is no unknotting closure tangle of 7_7 that connects the NW and the SE ends.




FIGURE 10. A coloring of the tangle 7_7 .

Theorem 5.2. *The 2-string tangle \mathcal{T} has a polychromatic coloring by R_n if and only if $\mathcal{T} + [\pm 2]$ has one such coloring. A similar result holds for $\mathcal{T} * [\pm 2]$.*

Proof: Any polychromatic coloring of \mathcal{T} can be extended trivially to $\mathcal{T} + [\pm 2]$. Reciprocally, consider a c -coloring of $\mathcal{T} + [\pm 2]$. Then the four arcs of $[\pm 2]$ must have the same color, hence the four boundary arcs of \mathcal{T} also have that color. If the coloring of $\mathcal{T} + [\pm 2]$ is nontrivial then the coloring of \mathcal{T} is also nontrivial. ■

This theorem shows, for instance, that 7_{16} has a polychromatic R_3 -coloring, since $7_{16} \approx \mathcal{T} * [-2]$, where $\mathcal{T} \approx 6_2$, and the tangle 6_2 has a polychromatic R_3 -coloring.

To prove that a tangle is not unknottable we can use polychromatic R_n -colorings of the tangle as this implies that any closure of the tangle would have more than n distinct colorings, being this an obstruction for the closure to be the unknot. However, this is not sufficient to prove that a tangle is not unlinkable, as an unlink with t components has n^t distinct R_n -colorings. Hence, we could consider the possibility of proving that \mathcal{T} is not unlinkable by showing that it has no non-trivial R_n -coloring for some n . The next theorem shows that this is also not a strategy to obstruct unlinkability.

The following theorems establish general results on R_n -colorings of tangles.

Theorem 5.3. *Every tangle \mathcal{T} with more than one string has a nontrivial R_n -coloring, for every n .*

Proof: Let \mathcal{T} be a tangle with k crossings and s strings. An R_n -coloring of \mathcal{T} is determined by a system of k linear equations (one for each crossing), with $k + s$ variables (one for each arc). The matrix defined by this system has rank at most k , thus has nullity at least s . Since $s > 1$, there exists a nontrivial R_n -coloring of \mathcal{T} . ■

We say that a R_n -coloring of a tangle verifies the *alternating sum rule* if, ordering the ends of the tangle clockwise, the sum of the colors of the odd ends is the same as the sum of the colors of the even ends. Note that a R_n -coloring of a 2-string tangle verifies the alternating sum rule if and only if the sum of colors of the *NW* and *SE* boundary arcs equals the sum of colors of the *NE* and *SW* boundary arcs. We say that a tangle \mathcal{T} verifies the *alternating sum rule* if every R_n -coloring of \mathcal{T} verifies the alternating sum rule.

Lemma 5.4. *Any tangle \mathcal{T} verifies the alternating sum rule.*

Proof: The tangles $[0]$, $[1]$, $[-1]$ and $[\infty]$ clearly verify the alternating sum rule.

Let \mathcal{T}_1 and \mathcal{T}_2 be tangles that verify the alternating sum rule. Consider the tangle $\mathcal{T} = \mathcal{T}_1 \cup \mathcal{T}_2$ obtained by identifying some ends of \mathcal{T}_1 and \mathcal{T}_2 . By ordering the ends of \mathcal{T}_2 in such a way that the common ends of \mathcal{T}_1 and \mathcal{T}_2 have opposite parities, we conclude that the difference of sum of the colors of the odd and even ends of \mathcal{T} equals the sum of differences coming from \mathcal{T}_1 and \mathcal{T}_2 , which is zero. Therefore, the sum of the colors of the odd ends of \mathcal{T} is the same as the sum of the colors of the even ends. ■

Definition 7. Let a, b, c, d be the *NW, NE, SW, SE* boundary colors of a R_0 d -coloring of a tangle \mathcal{T} . The *coloring fraction* of this coloring is defined by $\frac{b-a}{b-d}$.

In the definition of coloring fraction, note that as the R_0 coloring is a d -coloring, from Lemma 5.4, $b-a$ and $b-d$ cannot be simultaneously 0. Hence, the coloring fraction $\frac{b-a}{b-d}$ is well defined in $\mathbb{Q} \cup \{\pm\infty\}$.

The following theorem states that the coloring fraction of a rational tangle is the same as its arithmetical fraction.

Theorem 5.5 (Kauffman and Lambropoulou [7]). *The coloring fraction of a R_0 d -coloring of the rational tangle $\left[\frac{p}{q}\right]$ is $\frac{p}{q}$.*

Proof: For $\left[\frac{p}{q}\right] = [0]$ and $\left[\frac{p}{q}\right] = [\infty]$, the result is immediate.

Let $\mathcal{T} = \left[\frac{p}{q}\right]$ be a tangle such that $\frac{b-a}{b-d} = \frac{p}{q}$. Then the boundary colors of $\mathcal{T} + [1]$ are $a, 2b-d, c, b$. Since $\frac{2b-d-a}{2b-d-b} = \frac{b-a}{b-d} + 1 = \frac{p}{q} + 1$, the result holds for the tangle $\mathcal{T} + [1]$.

A similar reasoning (using the previous Lemma) shows that the result also holds for the tangles $\mathcal{T} + [-1]$, $\mathcal{T} * [1]$ and $\mathcal{T} * [-1]$. By successive applications of this property, the result holds for all rational tangles. ■

Theorem 5.6. *Consider a R_0 -coloring of the tangle $\mathcal{T} + \mathcal{U}$ and its restrictions to \mathcal{T} and \mathcal{U} . If these coloring are all d -colorings and the coloring fractions of \mathcal{T} and \mathcal{U} are r_1 and r_2 , then the coloring fraction of $\mathcal{T} + \mathcal{U}$ is $r_1 + r_2$.*

Proof: Let a, b, c, d be the boundary colors of \mathcal{T} and b, e, d, f be the boundary colors of \mathcal{U} . Then, since $a + d = b + c$ and $b + f = d + e$, we have

$$\frac{e - a}{e - f} - \left(\frac{b - a}{b - d} + \frac{e - b}{e - f} \right) = 0.$$

■

Theorem 5.7. *Let $L = N(\mathcal{T} + \mathcal{U})$ be a link with a nontrivial R_0 -coloring. If, for R_0 -colorings, \mathcal{T} and \mathcal{U} are monochromatic, then the coloring fractions of \mathcal{T} and \mathcal{U} are symmetric.*

Proof: If the coloring induced by L on \mathcal{T} and \mathcal{U} was a c -coloring, then it would be trivial. Therefore, the induced colorings are d -colorings. If the induced coloring of $\mathcal{T} + \mathcal{U}$ is a d -coloring, then, by the previous theorem, the coloring fraction of $\mathcal{T} + \mathcal{U}$, which is 0 for the NW and NE colors being the same, is the sum of the coloring fractions of \mathcal{T} and \mathcal{U} . If the induced coloring of $\mathcal{T} + \mathcal{U}$ is a c -coloring, then the coloring fraction of \mathcal{T} is $\pm\infty$ and the coloring fraction of \mathcal{U} is $\mp\infty$. ■

Corollary 5.7.1. *Let \mathcal{T} be a R_0 -monochromatic tangle with unlinking (or rational splitting) closure tangle $\left[\frac{p}{q} \right]$. Then \mathcal{T} has coloring fraction $\frac{-p}{q}$.*

Proof: Every split link $N\left(\mathcal{T} + \left[\frac{p}{q} \right]\right)$ has a nontrivial R_0 -coloring and $\left[\frac{p}{q} \right]$ is R_0 -monochromatic. ■

This corollary shows that a tangle with coloring invariant p/q can possibly only have $\left[\frac{-p}{q} \right]$ as its rational splitting closure tangle. However, even if the resulting link is not split, it has nevertheless zero determinant, as stated in the following theorem.

Theorem 5.8. *Let \mathcal{T} be a tangle with coloring invariant p/q . Then $N\left(\mathcal{T} + \left[\frac{-p}{q} \right]\right)$ is a link with determinant 0.*

Proof: Since there is a d -coloring of $\mathcal{T} + \left[\frac{-p}{q} \right]$ such that its north ends of have the same color, then $N\left(\mathcal{T} + \left[\frac{-p}{q} \right]\right)$ has a nontrivial R_0 -coloring. Therefore, for every prime number n larger than the colors of $\mathcal{T} + \left[\frac{-p}{q} \right]$,

$N\left(\mathcal{T} + \left[\frac{-p}{q}\right]\right)$ has a nontrivial R_n -coloring. Hence, n divides the determinant of $N\left(\mathcal{T} + \left[\frac{-p}{q}\right]\right)$, therefore this determinant must be zero. ■

6. Unknottability of essential 2-string tangles with crossing number at most 7

In this section, we classify unknottable unlinkable and splittable tangles with at most 7 crossings, with the notation of table 1 in [6].

Theorem 6.1. *An essential 2-string tangle with crossing number at most 7 is unknottable if and only if it is equivalent to 5_1 , 6_1 , 7_2 , 7_5 , 7_7 or 7_{14} .*

Proof: It can be easily checked (see Figure 11) that $N(5_1 + [-1])$, $N(6_1 + [-1])$, $N(7_2 + [-1])$, $N(7_5 + [0])$, $N(7_7 + [0])$ and $N(7_{14} + [-1])$ are unknotted.




FIGURE 11. The unknotting closure tangles of 5_1 , 6_1 , 7_2 , 7_5 , 7_7 and 7_{14} .

Reciprocally, the 2-string tangles 6_2 , 6_3 , 7_{13} , 7_{15} , 7_{16} , 7_{17} and 7_{18} have non-trivial colorings by dihedral quandles (see Section 5), therefore they are not unknottable. The remaining 2-string tangles with crossing number at most 7 are algebraic, with the expression given by Table 1, and a direct application of Theorems 4.5 and 4.6.1 shows that these tangles are not unknottable. ■

Theorem 6.2. *An essential 2-string tangle with crossing number at most 7 is unlinkable or splittable if and only if it is equivalent to 6_3 .*

Proof: It can be easily checked (see Figure 12) that $N(6_3 + [0])$ is the unlink.



FIGURE 12. The unlinking closure tangle of 6_3 .

By Theorem 4.4, the tangles 5_1 , 6_1 , 7_2 , 7_5 , 7_7 or 7_{14} are not unlinkable, since they are unknottable.

Tangle	Algebraic expression	Tangle	Algebraic expression
6_4	$([\frac{1}{3}] + [\frac{-1}{2}]) * [\frac{-2}{1}]$	7_8	$([\frac{-1}{2}] + [\frac{-1}{3}]) * [\frac{-2}{1}]$
7_1	$[\frac{1}{2}] + [\frac{1}{5}]$	7_9	$([\frac{-1}{2}] + [\frac{-2}{3}]) * [\frac{-2}{1}]$
7_3	$[\frac{1}{2}] + [\frac{2}{7}]$	7_{10}	$([\frac{-1}{2}] + [\frac{1}{3}]) * [\frac{3}{1}]$
7_4	$[\frac{1}{3}] + [\frac{1}{4}]$	7_{11}	$([\frac{-2}{3}] + [\frac{1}{2}]) * [\frac{-3}{1}]$
7_6	$[\frac{1}{3}] + [\frac{2}{5}]$	7_{12}	$([\frac{2}{3}] + [\frac{-1}{2}]) * [\frac{-2}{1}]$

TABLE 1. Algebraic expressions of 6_4 , 7_1 , 7_3 , 7_4 , 7_6 , 7_8 , 7_9 , 7_{10} , 7_{11} and 7_{12} .

The tangles 6_2 , 6_4 , 7_1 , 7_3 , 7_4 , 7_6 , 7_8 , 7_9 , 7_{10} , 7_{11} and 7_{12} are algebraic, with the expression given by Tables 1 and 2, and a direct application of Theorems 4.5 and 4.6.1 shows that these tangles are not unlinkable nor splittable.

Tangle	Algebraic expression	Tangle	Algebraic expression
6_2	$[\frac{1}{3}] + [\frac{1}{3}]$	7_{16}	$([\frac{1}{3}] + [\frac{1}{3}]) * [\frac{-2}{1}]$

TABLE 2. Algebraic expressions of 6_2 and 7_{16} .


The tangles 7_{13} , 7_{15} , 7_{17} and 7_{18} are R_0 -monochromatic and have the coloring invariants of Table 3. Hence, from Corollary 5.7.1, it remains to show that the corresponding rational tangles are not the unlinking or rational splitting closure tangles.

The link $N(7_{13} + [\frac{-3}{4}])$, as in Figure 13, is not split since its Jones polynomial, $-t^{-3} + t^{-1} - t - t^5 - t^7 + t^9 - t^{11} + t^{13}$, is different from the Jones polynomial of a split link defined by the right-handed trefoil and the unknot, $t^2 + t^6 - t^8$. The link $N(7_{15} + [\frac{-2}{3}])$, as in Figure 13, is not split since its Jones polynomial, $t^{-11} - t^{-9} - t^{-5} - t + t^3 - t^5$, is different from the Jones polynomial of the two component unlink, $-t - t^{-1}$. The links $N(7_{17} + [\frac{-8}{7}])$

Tangle	Coloring invariant	Tangle	Coloring invariant
7_{13}	$\frac{3}{4}$	7_{17}	$\frac{8}{7}$
7_{15}	$\frac{2}{3}$	7_{18}	2

TABLE 3. Coloring invariants of 7_{13} , 7_{15} , 7_{17} and 7_{18} .

and $N(7_{18} + [-2])$, as in Figure 13, are not split since their linking number is not zero. Therefore, from Corollary 5.7.1, the tangles 7_{13} , 7_{15} , 7_{17} and 7_{18} are not splittable, and hence also not unlinkable (we can also conclude that the tangle 7_{13} is not unlinkable since one of its strings is knotted). ■

FIGURE 13. The links $N(7_{13} + [\frac{-3}{4}])$, $N(7_{15} + [\frac{-2}{3}])$, $N(7_{17} + [\frac{-8}{7}])$, $N(7_{18} + [-2])$.

References

- [1] S. Bleiler, M. Scharlemann, *Tangles, property P, and a problem of J. Martin*, Math. Ann. 273 (1986), pp. 215-225.
- [2] S. Bleiler, M. Scharlemann, *A projective plane in \mathbb{R}^4 with three critical points is standard. Strongly invertible knots have property P*, Topology 27 No. 4 (1988), pp. 519-540.
- [3] T. Cochran, D. Ruberman, *Invariants of tangles*, Math. Proc. Camb. Phil. Soc. 105 (1989), pp. 299-306.
- [4] C. Ernst, D. W. Sumners, *A calculus for rational tangles: applications to DNA recombination*, Math. Proc. Camb. Phil. Soc. 108 (1990), pp. 489-515.
- [5] M. Eudave-Muñoz, *Primeness and sums of tangles*, Trans. of Amer. Math. Soc. 306 No. 2 (1988), pp. 773-790.
- [6] T. Kanenobu, H. Saito, S. Satoh, *Tangles with up to seven crossings*, Interdisciplinary Information Sciences 9 No. 1 (2003), pp. 127-140.
- [7] L. Kauffman, S. Lambropoulou, *On the classification of rational tangles*, Advances in Applied Mathematics 33 (2004), pp. 199-237.
- [8] L. Kauffman, S. Lambropoulou, *Hard unknots and collapsing tangles*, Series on Knots and Everything, Introductory Lectures on Knot Theory, World Scientific (2011), pp. 187-247.
- [9] L. Kauffman, P. Lopes, *The prevalence of persistent tangles*, Topology and its Applications, 271 (2020), 107040, 12 pp.
- [10] D. Krebes, *An obstruction to embedding 4-tangles in links*, J. Knot Theory and its Ramifications 8 No. 03 (1999), pp. 321-352.

- [11] D. Krebes, D. Silver, S. Williams, *Persistent invariants of tangles*, J. of Knot Theory and Its Ramifications 9 No. 4 (2000), pp. 471-477.
- [12] J. M. Montesinos, *Varietades de Seifert que son recubridores cclicos ramificados de dos hojas*, Bol. Soc. Mat. Mexicana 18, No. 2 (1973), pp. 1-32.
- [13] J. M. Nogueira, A. Salgueiro, *The minimum crossing number of essential tangles*, J. Knot Theory and its Ramifications 23 No. 10 (2014), 1450054, 15 pp.
- [14] J. H. Przytycki, D. D. Silver, S. G. Williams, *3-manifolds, tangles and persistent invariants*, Math. Proc. Cambridge Philos. Soc. 139 (2005), pp. 291-306.
- [15] D. Ruberman, *Embedding tangles in links*, J. of Knot Theory and its Ramifications 9 No. 4 (2000), pp. 523-530.
- [16] M. Scharlemann, *Smooth spheres in \mathbb{R}^4 with four critical points are standard*, Invent. Math. 79 (1985), pp. 125-141.
- [17] D. Silver, S. Williams, *Virtual tangles and a theorem of Krebes*, J. of Knot Theory and Its Ramifications 8 No. 7 (1999), pp. 941-945.
- [18] D. Silver, S. Williams, *Tangles and links: a view with trees*, J. of Knot Theory and Its Ramifications 27 No. 12 (2018), 1850061, 7 pp.
- [19] S. Taylor, *Boring split links and unknots*, Ph.D. Dissertation, University of California Santa Barbara, 2008.

CMUC, DEPARTMENT OF MATHEMATICS,
UNIVERSITY OF COIMBRA
E-mail: nogueira@mat.uc.pt

CMUC, DEPARTMENT OF MATHEMATICS,
UNIVERSITY OF COIMBRA
E-mail: ams@mat.uc.pt