

THE COFRAME OF D -SUBLOCALES OF A LOCALE AND THE T_D -DUALITY

IGOR ARRIETA AND ANNA LAURA SUAREZ

ABSTRACT: The notion of D -sublocale is explored. This is the notion analogue to that of sublocale in the duality of T_D -spaces. A sublocale S of a frame L is a D -sublocale if and only if the corresponding localic map preserves the property of being a covered prime. It is shown that for a frame L the system of those sublocales which are also D -sublocales form a dense sublocale $\mathbf{S}_D(L)$ of the coframe $\mathbf{S}(L)$ of all its sublocales. It is also shown that the spatialization $\mathbf{sp}_D[\mathbf{S}_D(L)]$ of $\mathbf{S}_D(L)$ consists precisely of those D -sublocales of L which are T_D -spatial. Additionally, frames such that we have $\mathbf{S}_D(L) \cong \mathcal{P}(\mathbf{pt}_D(L))$ — that is, those such that D -sublocales perfectly represent subspaces — are characterized as those T_D -spatial frames such that $\mathbf{S}_D(L)$ is the Booleanization of $\mathbf{S}(L)$.

KEYWORDS: Locale, T_D -space, coframe, totally spatial frame.

2020 MATHEMATICS SUBJECT CLASSIFICATION: 18F70, 06D22.

Introduction

For a point-free space (that is, a frame) L the ordered collection of its point-free subspaces, that is, its *sublocales*, is always a coframe usually denoted as $\mathbf{S}(L)$. Embedded in $\mathbf{S}(L)$ as a coframe we have the coframe $\mathbf{Cl}(L)$ of the *closed* sublocales of L , point-free versions of the closed subspaces of a space. We have an anti-isomorphism $L \rightarrow \mathbf{Cl}(L)$. Embedded in $\mathbf{S}(L)$ as a frame we also have $\mathbf{Op}(L)$, the ordered collection of all *open* sublocales of L ; these represent the open subspaces of the space L . It is not surprising, then, that the frame $\mathbf{Op}(L)$ is isomorphic to L . Since the system $\mathbf{S}(L)^{op}$ is a frame, it has its own sublocales, and among these there are certain special ones that encode information about the frame L . Next to the sublocales of $\mathbf{S}(L)^{op}$ that are particularly interesting are also certain subsets. Recently, some of these special sublocales and subsets have enjoyed special attention. We have the

Received December 15, 2020.

The first named author gratefully acknowledges support from the Basque Government (grant IT974-16 and predoctoral fellowship PRE-2018-1-0375), from the Spanish Ministry of Science and Innovation (reference code PID2019-103838GB-I00 (MCIU/AEI/FEDER, UE)) and from the Centre for Mathematics of the University of Coimbra - UIDB/00324/2020, funded by the Portuguese Government through FCT/MCTES.

system of all joins of closed sublocales $\mathbf{S}_c(L)$ (see [11]), which coincides with the smallest dense sublocale of $\mathbf{S}(L)$ if and only if the frame L satisfies a weak separation axiom called *subfitness*. We then have $\mathbf{S}_b(L)$, the smallest dense sublocale of $\mathbf{S}(L)$. The properties of this sublocale have been studied in [1]. We also have $\mathbf{sp}[\mathbf{S}(L)]$ (see [15]), the sublocale of $\mathbf{S}(L)$ consisting of those sublocales which are spatial — that is, those sublocales which are the frame of opens of some topological space.

We explore the relation these important subsets of the assembly $\mathbf{S}(L)$ of a frame L ; and we identify another which also encodes some special properties of L , in particular in the context of the T_D -duality. We will look at the notion of *D-sublocale* of L , a sublocale $S \subseteq L$ such that the corresponding frame surjection is a D -homomorphism (i.e. it is such that every prime covered in S is also covered in L). We take D -sublocales as being the analogues of sublocales in the T_D -duality. We show that for a frame L the D -sublocales form a coframe $\mathbf{S}_D(L)$ which is always a (D -)subcolocale of $\mathbf{S}(L)$.

We then relate the subcolocale $\mathbf{S}_D(L)$ with the subcolocale $\mathbf{sp}[\mathbf{S}(L)]$ (i.e. the ordered collection of all spatial sublocales of L); as well as with the subcolocale $\mathbf{S}_b(L)$ (i.e. the Booleanization subcolocale of $\mathbf{S}(L)$). We also relate all these structures with the subset $\mathbf{S}_c(L) \subseteq \mathbf{S}(L)$ of all joins of closed sublocales. We obtain characterizations of all possible set inclusions between these structures, as depicted in the table below.

Relations between sublocales of $\mathbf{S}(L)$	Properties of L
$\mathbf{S}_b(L) \subseteq \mathbf{sp}[\mathbf{S}(L)]$	Spatial
$\mathbf{S}_b(L) = \mathbf{sp}[\mathbf{S}(L)]$	Strongly T_D -spatial
$\mathbf{S}_b(L) = \mathbf{S}(L)$	Scattered
$\mathbf{S}(L) = \mathbf{S}_D(L)$	Primes are covered
$\mathbf{S}(L) = \mathbf{sp}(\mathbf{S}(L))$	Totally spatial
$\mathbf{S}_b(L) = \mathbf{S}_D(L)$	D -scattered
$\mathbf{S}_D(L) \subseteq \mathbf{S}_b(L)$	D -scattered
$\mathbf{S}_D(L) \subseteq \mathbf{sp}(\mathbf{S}(L))$	Totally spatial
$\mathbf{sp}[\mathbf{S}(L)] \subseteq \mathbf{S}_D(L)$	Primes are covered
$\mathbf{S}_c(L) \subseteq \mathbf{sp}[\mathbf{S}(L)]$	Spatial
$\mathbf{sp}[\mathbf{S}(L)] \subseteq \mathbf{S}_c(L)$	Primes are maximal
$\mathbf{S}_D(L) \subseteq \mathbf{S}_c(L)$	Subfit + D -scattered
$\mathbf{S}_D(L) = \mathbf{S}_c(L)$	Subfit + D -scattered
$\mathbf{S}(L) = \mathbf{S}_c(L)$	Subfit + scattered

We then show that we have an adjunction of posets

$$\begin{array}{ccc} & \mathfrak{M} & \\ & \curvearrowright & \\ \mathcal{P}(\mathbf{pt}_D(L)) & \perp & \mathbf{S}_D(L) \\ & \curvearrowleft & \\ & \mathbf{pt}_D & \end{array}$$

Here, $\mathbf{pt}_D(S)$ is the collection of covered primes of a D -sublocale S . As S is assumed to be a D -sublocale, there is no ambiguity in this phrase as primes of S are covered in S if and only if they are covered in L . The adjunct \mathbf{pt}_D , then, takes the T_D spectrum of a D -sublocale, while the adjunct \mathfrak{M} closes the subset under arbitrary meets. The fixpoints of $\mathbf{pt}_D \circ \mathfrak{M}$, then, are the equivalentents of the sober spaces in the T_D -duality (those isomorphic to the spectrum of their frame of opens); while the fixpoints of $\mathfrak{M} \circ \mathbf{pt}_D$ are those D -sublocales which play the role of spatial frames in the classical duality (i.e. they are those isomorphic to the frame of opens of their spectrum).

It is shown that the composition $\mathbf{pt}_D \circ \mathfrak{M}$ is the identity. It is shown that the fixpoints of $\mathfrak{M} \circ \mathbf{pt}_D$ are the spatial D -sublocales. We finally use this adjunction to ask ourselves how we can characterize those frames L such that the D -sublocales of L perfectly represent the subspaces of $\mathbf{pt}_D(L)$. We reach the conclusion that there is a fundamental difference in the way subspaces and sublocales behave in the T_D -duality, compared to the classical duality. In the classical spatial-sober duality, we could ask ourselves three questions. In the following $\mathbf{sob}[\mathcal{P}(\mathbf{pt}(L))]$ denotes the ordered collection of the sober subspaces of $\mathbf{pt}(L)$.

- (1) When is it that for a frame L we have that taking the spectrum induces an isomorphism $\mathbf{sp}[\mathbf{S}(L)] \cong \mathcal{P}(\mathbf{pt}(L))$? These are exactly the frames for which all subspaces of $\mathbf{pt}(L)$ are sober.
- (2) When is it that for a frame L we have that taking the spectrum induces an isomorphism $\mathbf{S}(L) \cong \mathbf{sob}[\mathcal{P}(\mathbf{pt}(L))]$? These are exactly the frames for which all sublocales of L are spatial (the *totally spatial* frames, explored for instance in [8]).
- (3) When is it that we have both of the above conditions? In other words, when is it that taking the spectrum induces an isomorphism $\mathbf{S}(L) \cong \mathcal{P}(\mathbf{pt}(L))$? Adapting a famous result by Simmons (see [14]) we find that these are the frames such that they are spatial and the space $\mathbf{pt}(L)$ is *scattered*.

In the T_D -duality, the answer to the analogue of the first question collapses to “always”, and so the second and the third question collapse to the same.

CONTENTS

Introduction	1
Preliminaries	4
The categories Loc and Frm	5
Sublocales	5
The axiom T_D	9
1. T_D -spatiality and strong T_D -spatiality	11
2. D -sublocales	13
2.1. Functoriality	15
3. The relation between D -sublocales and subspaces	17
3.1. Local and global T_D -spatialization	19
3.2. When do the D -sublocales perfectly represent the subspaces?	23
4. The relation between $\mathbf{S}_D(L)$ and $\mathbf{sp}[\mathbf{S}(L)]$	29
5. Adding $\mathbf{S}_c(L)$ to the picture	31
References	32

Preliminaries

We first recall some background on point-free topology. For more information on the categories of frames and locales, we refer the reader to Johnstone [7] or the more recent Picado-Pultr [9]. A *locale* (or *frame*) is a complete lattice L satisfying

$$a \wedge \bigvee B = \bigvee \{ a \wedge b \mid b \in B \}$$

for all $a \in L$ and $B \subseteq L$. A *frame homomorphism* is a function preserving arbitrary joins (including the bottom element 0) and finite meets (including the top element 1). Frames and their homomorphisms form a category **Frm**. For each a the map $a \wedge (-)$ preserves arbitrary joins, thus it has a right (Galois) adjoint $a \rightarrow (-)$, making L a complete Heyting algebra (i.e. a cartesian closed category, if one regards L as a thin category). This right adjoint is called the *Heyting operator*. In particular, the *pseudocomplement* of an $a \in L$ is the element $a^* = a \rightarrow 0$ and it can be characterized as the largest $b \in L$ such that $a \wedge b = 0$.

The categories \mathbf{Loc} and \mathbf{Frm} . Given a topological space X , its lattice of open sets $\Omega(X)$ is always a frame, and this construction can be upgraded to a functor $\Omega: \mathbf{Top} \rightarrow \mathbf{Frm}^{op}$ which sends a continuous map $f: X \rightarrow Y$ to the preimage operator $f^{-1}[-]: \Omega(Y) \rightarrow \Omega(X)$. An element p of a frame L is said to be *prime* if whenever $p = x \wedge y$ for some $x, y \in L$, then $p = x$ or $p = y$. The collection of all primes of L will be denoted by $\mathbf{pt}(L)$ and we shall refer to it as the *spectrum* (or the *classical spectrum* or *sober spectrum*) of L . For each $a \in L$, we set $\Sigma_a = \{p \in \mathbf{pt}(L) \mid a \not\leq p\}$. Then the family $\{\Sigma_a \mid a \in L\}$ is a topology on $\mathbf{pt}(L)$. The assignment $\Sigma(L) = (\mathbf{pt}(L), \{\Sigma_a \mid a \in L\})$ extends to an spectrum functor $\Sigma: \mathbf{Frm}^{op} \rightarrow \mathbf{Top}$ which yields an adjunction

$$\begin{array}{ccc} & \Omega & \\ \mathbf{Top} & \begin{array}{c} \curvearrowright \\ \perp \\ \curvearrowleft \end{array} & \mathbf{Frm}^{op} \\ & \Sigma & \end{array}$$

The category \mathbf{Loc} of locales is by definition the opposite category of \mathbf{Frm} :

$$\mathbf{Loc} = \mathbf{Frm}^{op},$$

and Ω restricts to a full embedding of a substantial part of \mathbf{Top} (namely the full subcategory of sober spaces) into \mathbf{Loc} . The latter can therefore be seen as a category of generalized spaces. We shall mostly speak of objects in \mathbf{Loc} as locales (instead of frames) when emphasizing the covariant approach. Morphisms in \mathbf{Loc} can be concretely represented by the right (Galois) adjoints $f_*: M \rightarrow L$ of the corresponding frame homomorphisms $f: L \rightarrow M$; these will be referred to as *localic maps*.

A frame is said to be *spatial* if $L \cong \Omega(X)$ for some space X , or equivalently if $L \cong \Omega(\Sigma(L)) = \{\Sigma_a \mid a \in L\}$. For any frame L , there is a surjective frame homomorphism (the counit of the adjunction) $L \twoheadrightarrow \Omega(\Sigma(L))$ sending a to Σ_a ; this map is usually called the *spatialization* of L .

A topological space is *sober* if every prime of $\Omega(X)$ is of the form $X \setminus \overline{\{x\}}$ for a unique $x \in X$. It is well-known that the space $\mathbf{pt}(L)$ defined above is sober for any frame L .

Sublocales. A regular subobject in \mathbf{Loc} (that is, an isomorphism class of regular monomorphisms) of a locale L is a *sublocale* of L . Sublocales of a locale L can be represented as the actual subsets $S \subseteq L$ such that

- (1) S is closed under arbitrary meets in L , and
- (2) $a \rightarrow s \in S$ for all $a \in L$ and $s \in S$.

A different, but equivalent, representation of sublocales is by means of nuclei, i.e. inflationary and idempotent maps $\nu: L \rightarrow L$ which preserve binary meets. The sublocale associated to a nucleus ν is the image $\nu[L]$, and conversely the nucleus associated to a sublocale $S \subseteq L$ is given by $\iota_S \circ \nu_S$ where ι_S denotes the inclusion of S into L and ν_S is its left adjoint frame homomorphism given by

$$\nu_S(a) = \bigwedge \{ s \in S \mid s \geq a \}.$$

A locale L is said to be *totally spatial* if each of its sublocales is spatial.

A sublocale should not be confused with a *subframe*; the latter is a subobject of a frame in the category \mathbf{Frm} . Subframes can be represented as subsets which are closed under arbitrary joins and finite meets.

Closed and open sublocales. For each $a \in L$, one has an *open sublocale* and a *closed sublocale*

$$\mathfrak{o}(a) = \{ b \mid b = a \rightarrow b \} = \{ a \rightarrow b \mid b \in L \} \quad \text{and} \quad \mathfrak{c}(a) = \uparrow a$$

which in the spatial case $L = \Omega(X)$ correspond to the open and closed subspaces.

If S is a sublocale of L , the *closure* of S in L , denoted by \overline{S} , is the smallest closed sublocale containing S , which can be computed as $\overline{S} = \mathfrak{c}(\bigwedge S)$. A sublocale S is *dense* if $\overline{S} = L$, or equivalently if $0 \in S$. A sublocale S is said to be *codense* if $\nu_S(a) = 1$ implies $a = 1$.

Boolean sublocales. For each $a \in L$, there is a sublocale $\mathfrak{b}(a) = \{ b \rightarrow a \mid b \in L \}$ which turns out to be Boolean. Moreover, every Boolean sublocale of L is of the form $\mathfrak{b}(a)$ for some $a \in L$. If $p \in L$ is a prime, it is easy to check that then $\mathfrak{b}(p) = \{ 1, p \}$, and these sublocales are usually referred to as *one-point sublocales*.

In particular, the sublocale $\mathfrak{b}(0) = \{ x^* \mid x \in L \}$ is called the *Booleanization* of L and it can be characterized either as the least dense sublocale of L or the unique Boolean dense sublocale of L .

The coframe of sublocales. The family

$$\mathbf{S}(L)$$

of all sublocales of L partially ordered by inclusion is a coframe (i.e. its dual poset $\mathbf{S}(L)^{op}$ is a frame) and lattice operations in $\mathbf{S}(L)$ are given by

$$\bigwedge_{i \in I} S_i = \bigcap_{i \in I} S_i, \quad \bigvee_{i \in I} S_i = \{ \bigwedge A \mid A \subseteq \bigcup_{i \in I} S_i \}.$$

It is not generally the case that $\mathbf{S}(L)$ is Boolean but it still has plenty of complemented sublocales: the frame $\mathbf{S}(L)^{op}$ is always zero-dimensional. More precisely, every sublocale S of L can be written $S = \bigcap_i \mathbf{c}(a_i) \vee \mathbf{o}(b_i)$ for some $\{a_i\}_{i \in I}, \{b_i\}_{i \in I}$, and closed sublocales $\mathbf{c}(a)$ and open sublocales $\mathbf{o}(a)$ are complements of each other. We also have

$$\begin{aligned} \mathbf{c}\left(\bigvee_i a_i\right) &= \bigcap_i \mathbf{c}(a_i), & \mathbf{o}\left(\bigvee_i a_i\right) &= \bigvee_i \mathbf{o}(a_i), \\ \mathbf{c}(a \wedge b) &= \mathbf{c}(a) \vee \mathbf{c}(b), & \mathbf{o}(a \wedge b) &= \mathbf{o}(a) \cap \mathbf{o}(b). \end{aligned}$$

Further, a sublocale is said to be *locally closed* if it is of the form $\mathbf{o}(a) \cap \mathbf{c}(b)$ for some $a, b \in L$.

Since $\mathbf{S}(L)$ is a coframe, there is a *co-Heyting operator* giving the *difference* $S \setminus T$ of two sublocales $S, T \in \mathbf{S}(L)$; this operator is characterized by the condition

$$S \setminus T \subseteq R \iff S \subseteq T \vee R.$$

In particular, the *supplement* of $S \in \mathbf{S}(L)$ is $S^\# = L \setminus S$, i.e. the smallest sublocale of L whose join with S is L . We shall freely use some its properties, e.g. the ones listed below

- (1) $S \setminus T \subseteq S$;
- (2) $S \setminus T = 0$ iff $S \subseteq T$;
- (3) $S \setminus C = S \cap C^\#$;
- (4) $S \setminus \bigcap_i S_i = \bigvee_i (S \setminus S_i)$;
- (5) $(S \setminus T) \setminus R = (S \setminus R) \setminus T$;

for each $S, T, R \in \mathbf{S}(L)$, $\{S_i\}_{i \in I} \subseteq \mathbf{S}(L)$ and complemented sublocale $C \subseteq L$. A comprehensive list of its properties may be found in [6].

For several notions related to coframes, we shall use terminology from frame theory modified just by adding the prefix “co-”. For example, a subset S of a coframe L will be said to be a *subcolocale* of L if S^{op} is a sublocale of the frame L^{op} (i.e. iff it is a subset of L closed under arbitrary joins and the co-Heyting operator). Sometimes we will omit the prefix “co-” in order to avoid confusion: we will speak of a dense (resp. spatial, T_D -spatial) subcolocale

S of a coframe L for meaning that S^{op} is a dense (resp. spatial, T_D -spatial) sublocale of L^{op} .

Images and preimages. Every localic map $f: L \rightarrow M$ gives, by pulling back in \mathbf{Loc} , an *inverse image* map $f_{-1}[-]: \mathbf{S}(M) \rightarrow \mathbf{S}(L)$ which turns out to be a coframe homomorphism (i.e. a function which preserves arbitrary meets and finite joins). Set-theoretic direct image yields a *direct image* map $f[-]: \mathbf{S}(L) \rightarrow \mathbf{S}(M)$ which is additionally a colocalic map (i.e. a left adjoint of a coframe homomorphism). In this context, the usual adjunction $f[-] \dashv f_{-1}[-]$ is satisfied.

Joins of closed (resp. complemented) sublocales. Let $\mathbf{S}_c(L)$ denote the subset of $\mathbf{S}(L)$ consisting of joins of closed sublocales i.e.

$$\mathbf{S}_c(L) = \left\{ \bigvee_{a \in A} \mathfrak{c}(a) \mid A \subseteq L \right\},$$

endowed with the inclusion order inherited from $\mathbf{S}(L)$. In the recent paper [11], Picado, Pultr and Tozzi show (a.o.) that $\mathbf{S}_c(L)$ is *always* a frame which is embedded as a join-sublattice in the coframe $\mathbf{S}(L)$. One of the main results from [11] is that

if L is subfit, and only in that case, $\mathbf{S}_c(L)$ is a Boolean algebra and coincides precisely with the Booleanization of $\mathbf{S}(L)$.

More generally, if L is not necessarily subfit, it is also of interest to consider the Booleanization of $\mathbf{S}(L)$, denoted by $\mathbf{S}_b(L)$. The latter is precisely the system consisting *smooth* sublocales, that is, those which are joins of complemented sublocales (or equivalently joins of locally closed sublocales). A study of the properties of $\mathbf{S}_b(L)$ may be found in [1].

Subspaces and sublocales. In [15], the relation between sublocales of a frame L and subspaces of its spectrum $\mathbf{pt}(L)$ is explored. Spatial sublocales and sober subspaces are seen as fixpoints of the following adjunction of posets.

$$\begin{array}{ccc} & \mathfrak{M} & \\ & \curvearrowright & \\ \mathcal{P}(\mathbf{pt}(L)) & \perp & \mathbf{S}(L) \\ & \curvearrowleft & \\ & \mathbf{pt} & \end{array}$$

Here, $\mathbf{pt}: \mathbf{S}(L) \rightarrow \mathcal{P}(\mathbf{pt}(L))$ maps each sublocale to the collection of primes contained in it, while $\mathfrak{M}: \mathcal{P}(\mathbf{pt}(L)) \rightarrow \mathbf{S}(L)$ takes a collection of primes of L and it closes it under arbitrary meets. For every frame L , we have that

its spatialization is a surjection, and so it corresponds to a certain sublocale that we will denote by $\mathbf{sp}(L)$. For a sublocale $S \subseteq L$, its spatialization sublocale is $\mathfrak{M}(\mathbf{pt}(S))$. Hence, the composition $\mathfrak{M} \circ \mathbf{pt}$ of the two adjoints above is spatialization. Similarly, the composition $\mathbf{pt} \circ \mathfrak{M}$ is sobrification, up to homeomorphism. For every frame L we have the following commuting diagram in the category of coframes. The bottom arrow sends a sublocale S to its prime spectrum $\mathbf{pt}(S)$, which simply consists of the collection $\mathbf{pt}(L) \cap S$.

$$\begin{array}{ccc} \mathbf{sp}[\mathbf{S}(L)] & \xrightarrow{\mathbf{pt}(\cong)} & \mathbf{sob}[\mathcal{P}(\mathbf{pt}(L))] \\ \mathbf{sp} \uparrow & & \downarrow \\ \mathbf{S}(L) & \xrightarrow{\mathbf{pt}} & \mathcal{P}(\mathbf{pt}(L)) \end{array}$$

The coframe $\mathbf{sp}[\mathbf{S}(L)]$ denotes the ordered collection of spatial sublocales of L , while $\mathbf{sob}[\mathcal{P}(\mathbf{pt}(L))]$ denotes the ordered collection of sober subspaces of $\mathbf{pt}(L)$. Finally, the vertical arrow maps each sublocale S to its spatialization $\mathfrak{M}(\mathbf{pt}(S))$.

The axiom T_D . A space X is defined to be T_D if every point $x \in X$ has some neighborhood U with $U \setminus \{x\}$ open. This is an axiom stronger than T_0 and weaker than T_1 and it was introduced in [2]. It has been used (for instance in [12]) in order to answer the question of when a topological space can be completely recovered from its frame of opens. The *Skula space* of a space X , denoted as $Sk(X)$, is the space defined as follows. The set of points is the same set of points as X , while the topology is the *Skula topology*, that is the one generated by the opens of X together with their complements. The following characterizations of T_D -spaces can be found in [9].

Proposition 0.1. *The following are equivalent for a T_0 space X .*

- (1) *The space is T_D .*
- (2) *For no $x \in X$ do we have that the dualization of the inclusion $X \setminus \{x\} \subseteq X$ is a frame isomorphism.*
- (3) *If $Y \not\subseteq Z$ then $\Omega'(Y) \not\subseteq \Omega'(Z)$, for all subspaces $Y, Z \subseteq X$.*
- (4) *The Skula space $Sk(X)$ is discrete.*

One can exhibit the role of the axiom T_D in point-free topology by comparing it to sobriety. This has been done in detail in [4]. In particular, in the context of T_0 spaces, the two axioms mirror each other in that sober spaces are maximal in the same sense in which T_D -spaces are minimal.

- A space X is sober if and only if we can never have a nontrivial subspace inclusion $X \hookrightarrow Y$ such that its dualization is an isomorphism.
- A space X is T_D if and only if we can never have a nontrivial subspace inclusion $Y \hookrightarrow X$ such that its dualization is an isomorphism.

Comparing the two axioms with respect to their implications for subspaces and sublocales yields the following. First, define $\mathbf{sp}[\mathbf{S}(\Omega(X))]$ as being the ordered collection of spatial sublocales of $\Omega(X)$. We always have a map $\Omega': \mathcal{P}(X) \rightarrow \mathbf{sp}[\mathbf{S}(\Omega(X))]$, which to each subspace of X assigns the sublocale that it induces on $\Omega(X)$. The following is well-known:

- A space X is sober if and only if the map $\Omega': \mathcal{P}(X) \rightarrow \mathbf{sp}[\mathbf{S}(\Omega(X))]$ is a surjection.
- A space X is T_D if and only if the map $\Omega': \mathcal{P}(X) \rightarrow \mathbf{sp}[\mathbf{S}(\Omega(X))]$ is an injection.

A frame L is said to be T_D -spatial if $L \cong \Omega(X)$ for some T_D -topological space X . A *covered prime* of L is a $p \neq 1$ such that whenever $p = \bigwedge_i x_i$ for some $\{x_i\}_{i \in I} \subseteq L$, then there is an $i \in I$ with $p = x_i$. In general, it is not true that localic maps map covered primes into covered primes. Following [4] a frame homomorphism f will be said to be a D -homomorphism if its associated localic map f_* sends covered primes into covered primes. The subset $\mathbf{pt}_D(L)$ will denote the set of covered primes of L , and we will refer to it as the T_D spectrum of L . For each $a \in L$, set $\Sigma'_a = \{p \in \mathbf{pt}_D(L) \mid a \not\leq p\}$. Then, the family $\{\Sigma'_a \mid a \in L\}$ is a topology on $\mathbf{pt}_D(L)$, and in fact $(\mathbf{pt}_D(L), \{\Sigma'_a \mid a \in L\})$ is a T_D -space [4].

Let \mathbf{Top}_D be the full subcategory of \mathbf{Top} consisting of T_D -spaces and let \mathbf{Frm}_D be the non-full subcategory of \mathbf{Frm} whose morphisms are the D -homomorphisms. In [4], the authors showed that the assignment

$$\Sigma'(L) = (\mathbf{pt}_D(L), \{\Sigma'_a \mid a \in L\})$$

yields an adjunction

$$\begin{array}{ccc} & \Omega & \\ \mathbf{Top}_D & \begin{array}{c} \curvearrowright \\ \perp \\ \curvearrowleft \end{array} & \mathbf{Frm}_D^{op} \\ & \Sigma' & \end{array}$$

Analogously to the case of the classical spectrum, a frame L is T_D -spatial iff $L \cong \Omega(\mathbf{pt}_D(L))$, and there is always a surjective frame homomorphism

(the counit of the adjunction) $L \rightarrow \Omega(\mathbf{pt}_D(L))$ sending a to Σ'_a . We will refer to $\Omega(\mathbf{pt}_D(L))$ as the T_D -spatialization of L .

1. T_D -spatiality and strong T_D -spatiality

Among the matters that we will tackle in this paper there is that of studying the system of all T_D -spatial sublocales of a frame. This is why we begin with an analysis of T_D -spatiality, and of a natural strengthening of this property which we will call “strong T_D -spatiality”. We recall that a frame L is T_D -spatial if it is isomorphic to the frame of opens of the subspace of its prime spectrum consisting of the covered primes. A natural strengthening of this condition is the following. We say that a frame L is *strongly T_D -spatial* if it is spatial and all its primes are covered. Thus, a strongly T_D -spatial is isomorphic to the frame of opens of its T_D spectrum, and furthermore its T_D spectrum coincides with the classical (sober) spectrum. In this section we will characterize both these conditions on a frame; in particular we will show that a frame is T_D -spatial if and only if the Booleanization of its assembly is spatial, and that a frame is strongly T_D -spatial if and only if the Booleanization of its assembly coincides with its spatialization.

Lemma 1.1 ([6, Proposition 10.2]). *For a frame L and a prime $p \in \mathbf{pt}(L)$ we have that $\mathbf{b}(p)$ is complemented in $\mathbf{S}(L)$ if and only if p is a covered prime.*

From the fact that $\mathbf{S}_b(L)$ is the collection of joins of sublocales complemented in $\mathbf{S}(L)$, we deduce the following corollary.

Corollary 1.2. *For a frame L and a prime $p \in \mathbf{pt}(L)$ we have that $\mathbf{b}(p) \in \mathbf{S}_b(L)$ if and only if $\mathbf{b}(p)$ is a covered prime. Then, for a frame L and for $\{p_i\}_{i \in I}$ a collection of covered primes, we have that all joins of the form $\bigvee_i \mathbf{b}(p_i)$ are in $\mathbf{S}_b(L)$.*

We recall that a frame L is spatial if and only if every element is a meet of primes. We have the following analogous characterization of T_D -spatiality.

Lemma 1.3. *A frame is T_D -spatial if and only if all its elements are meets of covered primes.*

Proof: For the “only if” part, if $L = \Omega(X)$ with X a T_D -space and U is an open, then $U = \bigwedge_{x \notin U} X - \overline{\{x\}}$, where each $X - \overline{\{x\}}$ is covered. Conversely, it is clear that if every element is a meet of covered primes, then the T_D -spatialization surjection $L \twoheadrightarrow \Omega(\mathbf{pt}_D(L))$ defined in the Preliminaries is

also one-to-one and hence an isomorphism. Since $\mathbf{pt}_D(L)$ is a T_D -space, then L is T_D -spatial. ■

We are now ready to prove our characterization theorem for T_D -spatiality.

Theorem 1.4. *For a frame L , the following are equivalent:*

- (1) L is T_D -spatial;
- (2) Every element of L is a meet of covered primes;
- (3) $\mathbf{S}_b(L)$ is spatial;
- (4) $\mathbf{S}_b(L) \cong \mathcal{P}(\mathbf{pt}_D(L))$.

Proof: The equivalence between (1) and (3) appears in [1, Theorem 6.2] and the equivalence between (1) and (2) follows by Lemma 1.3. Let us now show that (3) implies (4). If $\mathbf{S}_b(L)$ is spatial, as this is a Boolean algebra it must be isomorphic to the powerset of its primes. By Corollary 1.2, there is a bijection $\mathbf{pt}(\mathbf{S}_b(L)) \cong \mathbf{pt}_D(L)$. Finally note that (4) implies (3) is trivial. ■

We now move on to looking at strong T_D -spatiality.

Lemma 1.5. *A frame L is spatial if and only if the spatialization of $\mathbf{S}(L)^{op}$ is a dense frame surjection.*

Proof: A frame L is spatial if and only if L is a fixpoint of the interior $\mathbf{sp}: \mathbf{S}(L) \rightarrow \mathbf{S}(L)$, that is, if and only if the bottom element $L \in \mathbf{S}(L)^{op}$ is a fixpoint of the spatialization nucleus $\mathbf{sp}: \mathbf{S}(L)^{op} \rightarrow \mathbf{S}(L)^{op}$. ■

Since $\mathbf{S}_b(L)$ is the least dense sublocale of $\mathbf{S}(L)$ then one has the following:

Corollary 1.6. *For a frame L the following are equivalent:*

- (1) L is spatial;
- (2) $\mathbf{S}_b(L) \subseteq \mathbf{sp}[\mathbf{S}(L)]$.

We may now prove the characterization theorem for strong T_D -spatiality.

Theorem 1.7. *For a frame L , the following are equivalent:*

- (1) L is strongly T_D -spatial;
- (2) Every element of L is a meet of covered primes and every prime of L is covered;
- (3) The frame L is spatial and $\mathbf{pt}(L)$ is a T_D -space;
- (4) The frame L is isomorphic to $\Omega(X)$ for some sober T_D -space X ;
- (5) The frame L is spatial and we have an isomorphism $\mathbf{sp}[\mathbf{S}(L)] \cong \mathcal{P}(\mathbf{pt}(L))$;
- (6) $\mathbf{S}_b(L) = \mathbf{sp}[\mathbf{S}(L)]$.

Proof: (1) \implies (2). This follows from Lemma 1.3.

(2) \implies (3). If every element of L is a meet of primes, the frame L is spatial. Furthermore, if all primes of L are covered we have that $\mathbf{pt}(L)$ is a T_D -space, as it coincides with the T_D spectrum of L .

(3) \implies (4). This is clear as $\mathbf{pt}(L)$ is always sober.

(4) \implies (5). We recall that if a space X is T_D then the map $\Omega': \mathcal{P}(X) \rightarrow \mathbf{sp}[\mathbf{S}(\Omega(X))]$ is injective, while if it is sober it is surjective (see the subsection on the T_D axiom in the preliminaries). Thus, if $L \cong \Omega(X)$ for some T_D and sober space X we have an isomorphism $\mathcal{P}(\mathbf{pt}(L)) \cong \mathbf{sp}[\mathbf{S}(\mathbf{pt}(L))]$.

(5) \implies (6). If the frame L is spatial and we have an isomorphism $\mathbf{sp}[\mathbf{S}(L)] \cong \mathcal{P}(\mathbf{pt}(L))$, then in particular the subcolocale $\mathbf{sp}[\mathbf{S}(L)] \subseteq \mathbf{S}(L)$ is Boolean. As L is spatial it is also dense (see Lemma 1.5), and so it must be its Booleanization.

(6) \implies (1). By Corollary 1.6, L is spatial. Let p be a prime of L . Then $\mathfrak{b}(p) \in \mathbf{sp}[\mathbf{S}(L)] \subseteq \mathbf{S}_b(L)$, and so by Corollary 1.2 it follows that p is covered. \blacksquare

2. D -sublocales

We now investigate the notion equivalent to that of sublocale in the T_D -duality; the notion of D -sublocale. For a sublocale to count as a D -sublocale, we will require that its corresponding localic inclusion is a morphism in the category \mathbf{Loc}_D . This constraint amounts to the condition that the localic map ought to send covered primes to covered primes. In this section we will analyze the collection $\mathbf{S}_D(L) \subseteq \mathbf{S}(L)$ of D -sublocales of a frame L . In particular, we will show that $\mathbf{S}_D(L) \subseteq \mathbf{S}(L)$ is always a dense subcolocale. We then will explore the question of how close the assignment $\mathbf{S}_D(L)(-): \mathbf{Obj}(\mathbf{Frm}) \rightarrow \mathbf{Obj}(\mathbf{Frm})$ is to being functorial. It is known that localic maps do not generally send covered primes into covered primes; those frame homomorphisms such that their right adjoint localic maps send covered primes into covered primes were referred to as D -homomorphisms in [9]. Now we introduce one of the central notions in this paper: a sublocale S of L will be said to be a D -sublocale if the corresponding frame surjection $L \twoheadrightarrow S$ is a D -homomorphism, i.e. iff the equality $\mathbf{pt}_D(S) = \mathbf{pt}_D(L) \cap S$ (equivalently the inclusion $\mathbf{pt}_D(S) \subseteq \mathbf{pt}_D(L)$) holds.

In general, not every sublocale is a D -sublocale, and neither is an intersection of two D -sublocales, as the following example shows:

Remark 2.1. One of the simplest examples of an intersections of two D -sublocales which is not a D -sublocale seems to be the following: let $L = [0, 1]$ (the unit interval with its usual total order). A subset of a totally ordered set is a sublocale iff it is closed under meets, so the following two subsets are indeed sublocales:

$$S = \{0, 1\} \cup \bigcup_{n \in \mathbb{N}} \left[\frac{1}{2n}, \frac{1}{2n-1} \right) \quad \text{and} \quad T = \{0, 1\} \cup \bigcup_{n \in \mathbb{N}} \left[\frac{1}{2n+1}, \frac{1}{2n} \right).$$

Obviously, for every $x \neq 1$ in S , one has $x = \bigwedge_{x < t \in S} t$, and this shows that $\text{pt}_D(S) = \emptyset$, so S is a D -sublocale of L . Similarly, $\text{pt}_D(T) = \emptyset$ and so it is also a D -sublocale of L . Now, observe that $S \cap T = \{0, 1\}$ and then one has $\text{pt}_D(S \cap T) = \text{pt}(S \cap T) = \{0\}$. But 0 is not a covered prime in L as $0 = \bigwedge_{t > 0} t$. Hence $S \cap T$ is not a D -sublocale.

Let $\mathbf{S}_D(L)$ be the subset of $\mathbf{S}(L)$ consisting of D -sublocales of L , equipped with the inherited order. We observe that we have the following.

$$\mathbf{S}_D(L) = \{ S \in \mathbf{S}(L) \mid \text{pt}_D(S) \subseteq \text{pt}_D(L) \}.$$

In what follows, we investigate the structure of $\mathbf{S}_D(L)$:

Proposition 2.2. *The system $\mathbf{S}_D(L)$ is closed under arbitrary joins in $\mathbf{S}(L)$.*

Proof: Let $\{S_i\}_{i \in I} \subseteq \mathbf{S}_D(L)$, i.e. $\text{pt}_D(S_i) \subseteq \text{pt}_D(L)$ for all $i \in I$. We have to check that $\bigvee_i S_i \in \mathbf{S}_D(L)$, that is, $\text{pt}_D(\bigvee_i S_i) \subseteq \text{pt}_D(L)$. Let then $p \in \text{pt}_D(\bigvee_i S_i)$. Since $p \in \text{pt}_D(\bigvee_i S_i) \subseteq \bigvee_i S_i$, there is a family $\{a_i\}_{i \in I} \subseteq L$ such that $a_i \in S_i$ for each $i \in I$ and with $p = \bigwedge_i a_i$. Since for all $i \in I$ one has $a_i \in S_i \subseteq \bigvee_i S_i$ and $p \in \text{pt}_D(\bigvee_i S_i)$, then there is an $i_0 \in I$ such that $p = a_{i_0} \in S_{i_0}$. We claim that $p \in \text{pt}_D(S_{i_0})$. Indeed, let $\{b_j\}_{j \in J} \subseteq S_{i_0}$ a family such that $p = \bigwedge_j b_j$. Then $b_j \in S_{i_0} \subseteq \bigvee_i S_i$ and since $p \in \text{pt}_D(\bigvee_i S_i)$, there is a $j_0 \in J$ with $p = b_{j_0}$. Then $p \in \text{pt}_D(S_{i_0}) \subseteq \text{pt}_D(L)$, as desired. \blacksquare

Proposition 2.3. *If $S \in \mathbf{S}_D(L)$ and $T \in \mathbf{S}(L)$, then $S \setminus T \in \mathbf{S}_D(L)$.*

Proof: Let $p \in \text{pt}_D(S \setminus T)$. Since $\mathbf{S}(L)^{op}$ is zero-dimensional, write $T = \bigcap_i C_i$ with C_i complemented. Then $S \setminus T = S \setminus \bigcap_i C_i = \bigvee_i S \setminus C_i = \bigvee_i S \cap C_i^c$. Now, $p \in S \setminus T = \bigvee_i S \cap C_i^c$ and so there is a family $\{a_i\}_{i \in I}$ with $a_i \in S \cap C_i^c$ and $p = \bigwedge_i a_i$. Since $a_i \in S \cap C_i^c \subseteq S \setminus T$ and $p \in \text{pt}_D(S \setminus T)$, then there is an $i_0 \in I$ with $p = a_{i_0} \in C_{i_0}^c$. Since $T = \bigcap_i C_i$, then we conclude $p \notin T$. Now, $p \notin T$

is equivalent to $\mathfrak{b}(p) \not\subseteq T$, which is in turn equivalent to $\mathfrak{b}(p) \setminus T \neq 0$. But $\mathfrak{b}(p) \setminus T \subseteq \mathfrak{b}(p) = \{1, p\}$, and so $\mathfrak{b}(p) \setminus T = \mathfrak{b}(p)$. I claim that $\mathfrak{b}(p) \not\subseteq S \setminus \mathfrak{b}(p)$. By way of contradiction assume otherwise. Then $(S \setminus T) \setminus \mathfrak{b}(p) = (S \setminus \mathfrak{b}(p)) \setminus T \supseteq \mathfrak{b}(p) \setminus T = \mathfrak{b}(p)$. Then $0 \neq \mathfrak{b}(p) = \mathfrak{b}(p) \cap [(S \setminus T) \setminus \mathfrak{b}(p)]$, which is a contradiction since $\mathfrak{b}(p)$ is complemented in $S \setminus T$. Therefore, we must have $\mathfrak{b}(p) \not\subseteq S \setminus \mathfrak{b}(p)$, which is equivalent to $0 = \mathfrak{b}(p) \cap (S \setminus \mathfrak{b}(p))$. Then $\mathfrak{b}(p)$ is complemented in S , i.e. $p \in \mathbf{pt}_D(S) \subseteq \mathbf{pt}_D(L)$, as we wanted. \blacksquare

Corollary 2.4. $\mathbf{S}_D(L)$ is a dense D -sublocale of $\mathbf{S}(L)$. In particular it is a coframe.

Proof: The fact that it is a sublocale follows from the two previous propositions. Moreover, density follows from the obvious fact that L (i.e. the bottom element of $\mathbf{S}(L)^{op}$) belongs to $\mathbf{S}_D(L)$. Finally, since $\mathbf{S}(L)^{op}$ is a zero-dimensional frame, it is in particular regular; and it is well-known that primes in any regular frame are maximal (so in particular they are covered). Thus, every sublocale of $\mathbf{S}(L)^{op}$ is a D -sublocale. \blacksquare

Density of $\mathbf{S}_D(L)$ has an important consequence: it implies that $\mathbf{S}_b(L) \subseteq \mathbf{S}_D(L)$. Hence, all open sublocales and closed sublocales are D -sublocales, and so are locally closed sublocales or more generally smooth sublocales.

The system $\mathbf{S}_D(L)$ contains several different families of sublocales, for example:

- (1) All smooth sublocales, as we have just pointed out;
- (2) All pointless sublocales of L (as $\mathbf{pt}_D(S) \subseteq \mathbf{pt}(S) = \emptyset$);
- (3) Any join of pointless sublocales (joins of pointless sublocales may contain points, but their join will still be a D -sublocale because of Proposition 2.2);
- (4) All codense sublocales of L in which primes are maximal (these include codense naturally Hausdorff sublocales, codense fit sublocales, ...). This only of interest if L not subfit, cf. [5]. In order to show this assertion, one easily checks that if p is a maximal element in a codense sublocale S , then it is maximal in L , and in particular it belongs to $\mathbf{pt}_D(L)$.

2.1. Functoriality. As we will see below, the assignment $L \mapsto \mathbf{S}_D(L)^{op}$ cannot be made functorial in the whole of the category of frames in such a way that there is a natural transformation $\mathbf{c}: 1_{\mathbf{Frm}} \rightarrow \mathbf{S}_D(-)^{op}$. Therefore, one has to deal with *lifts* of individual frame homomorphisms. Let $f: L \rightarrow M$ be

a frame homomorphism and suppose there is a further frame homomorphism $\mathbf{S}_D(f): \mathbf{S}_D(L)^{op} \rightarrow \mathbf{S}_D(M)^{op}$ together with a commutative square

$$\begin{array}{ccc} \mathbf{S}_D(L)^{op} & \xrightarrow{\mathbf{S}_D(f)} & \mathbf{S}_D(M)^{op} \\ \uparrow \mathfrak{c}_L & & \uparrow \mathfrak{c}_M \\ L & \xrightarrow{f} & M \end{array}$$

If S is any sublocale of L , then one can write $S = \bigcap_{S \subseteq \mathfrak{o}(a) \vee \mathfrak{c}(b)} \mathfrak{o}(a) \vee \mathfrak{c}(b)$. Each of the sublocales $\mathfrak{o}(a) \vee \mathfrak{c}(b)$ belongs to $\mathbf{S}_D(L)$ (as open sublocales and closed sublocales are D -sublocales and $\mathbf{S}_D(L)$ is closed under joins). Now, if S happens to be a D -sublocale, then the intersection $\bigcap_{S \subseteq \mathfrak{o}(a) \vee \mathfrak{c}(b)} \mathfrak{o}(a) \vee \mathfrak{c}(b)$ coincides with the meet $\bigwedge_{S \subseteq \mathfrak{o}(a) \vee \mathfrak{c}(b)} \mathfrak{o}(a) \vee \mathfrak{c}(b)$ taken in $\mathbf{S}_D(L)$ (incidentally, this, together with the fact that $\mathfrak{o}(a) \vee \mathfrak{c}(b)$ are complemented in $\mathbf{S}_D(L)$, shows that $\mathbf{S}_D(L)^{op}$ is a zero-dimensional frame). Hence, each $S \in \mathbf{S}_D(L)$ can be written as $S = \bigvee_{\mathfrak{o}(a) \wedge \mathfrak{c}(b) \leq S} \mathfrak{o}(a) \wedge \mathfrak{c}(b)$ (here we are using the reverse order), where \bigvee is nothing but the join in $\mathbf{S}(L)^{op}$. Now, applying the frame homomorphism $\mathbf{S}_D(f)$ and using the fact that frame homomorphisms commute with complements, we obtain $\mathbf{S}_D(f)(S) = \bigvee_{\mathfrak{o}(a) \wedge \mathfrak{c}(b) \leq S} \mathfrak{c}(f(a)) \wedge \mathfrak{o}(f(a))$, where \bigvee denotes join in $\mathbf{S}_D(M)^{op}$. Denote $g = f_*$. Then, one has that $\mathbf{S}_D(f)(S)$ can be computed as the largest D -sublocale of M contained in $g_{-1}[S]$.

Proposition 2.5. *Let L a frame and $f: L \twoheadrightarrow S$ a surjection onto a sublocale S . Then f lifts if and only if it is a D -homomorphism (i.e. iff S is a D -sublocale of L).*

Proof: For the “if” implication, assume that S is a D -sublocale of L and define a map $h: \mathbf{S}_D(L) \rightarrow \mathbf{S}_D(S)$ given by $h(T) = T \wedge S$ (where \wedge stands for meet in $\mathbf{S}_D(L)$). We note that h is well-defined because since $T \wedge S \in \mathbf{S}_D(L)$ in particular we have $T \wedge S \in \mathbf{S}_D(S)$. Also, since S is a D -sublocale of L , it is easy to check that h preserves arbitrary meets (and obviously finite joins as well). Hence it is a (surjective) coframe homomorphism. For showing that the relevant naturality square commutes it suffices to check that $S \cap \mathfrak{c}(a)$ is a D -sublocale of L . Let $p \in \mathbf{pt}_D(S \cap \mathfrak{c}(a))$. If $p = \bigwedge_i x_i$ for a family $\{x_i\}_{i \in I} \subseteq S$, then since $p \in \mathfrak{c}(a)$, one has $a \leq p \leq x_i$ for each $i \in I$ and so $x_i \in S \cap \mathfrak{c}(a)$ for each $i \in I$. Hence there is an $i_0 \in I$ with $p = x_{i_0}$. Thus $p \in \mathbf{pt}_D(S) \subseteq \mathbf{pt}_D(L)$.

Let us now show the converse, so assume that there is a frame homomorphism

$h: \mathbf{S}_D(L)^{op} \rightarrow \mathbf{S}_D(S)^{op}$ making the square

$$\begin{array}{ccc} \mathbf{S}_D(L)^{op} & \xrightarrow{h} & \mathbf{S}_D(S)^{op} \\ \uparrow \mathfrak{c}_L & & \uparrow \mathfrak{c}_S \\ L & \xrightarrow{f} & S \end{array}$$

commutative. We claim that h is a surjection too. Indeed, we noted in the discussion preceding the statement that $\mathbf{S}_D(S)^{op}$ is generated by $\{ \mathfrak{c}_S(a), \mathfrak{o}_S(b) \mid a, b \in S \}$, and thus it suffices to observe that $h(\mathfrak{c}_L(a)) = \mathfrak{c}_S(a)$ and $h(\mathfrak{o}_L(a)) = \mathfrak{o}_S(a)$ for each $a, b \in S$. Hence h is a surjection. Let us finally check that S is a D -sublocale of L . Let $p \in \mathbf{pt}_D(S)$. Then by the converse implication, the surjection $S \twoheadrightarrow \mathfrak{b}(p)$ lifts, and so we obtain a commutative diagram as follows:

$$\begin{array}{ccccc} \mathbf{S}_D(L)^{op} & \xrightarrow{h} & \mathbf{S}_D(S)^{op} & \twoheadrightarrow & \mathbf{S}_D(\mathfrak{b}(p))^{op} \\ \uparrow \mathfrak{c}_L & & \uparrow \mathfrak{c}_S & & \cong \uparrow \\ L & \xrightarrow{f} & S & \twoheadrightarrow & \mathfrak{b}(p) \end{array}$$

where the top composite is also a surjection. Hence there is a surjection $\mathbf{S}_D(L)^{op} \twoheadrightarrow \mathfrak{b}(p)$ which displays $\mathfrak{b}(p)$ as a point of $\mathbf{S}_D(L)^{op}$, but Lemma 3.9 below asserts that then p must be a covered prime of L . \blacksquare

Remark 2.6. Characterizing monomorphisms which lift seems to be a more difficult task. An analogous situation happens when studying the functoriality of the assignment $L \mapsto \mathbf{S}_b(L)$ (cf. [1, Proposition 5.10]).

3. The relation between D -sublocales and subspaces

In [15] the matter of the relation between subspaces and sublocales is explored. Recall that there it is proven that we have an adjunction

$$\begin{array}{ccc} \mathcal{P}(\mathbf{pt}(L)) & \begin{array}{c} \xrightarrow{\mathfrak{m}} \\ \perp \\ \xleftarrow{\mathbf{pt}} \end{array} & \mathbf{S}(L) \end{array}$$

in which the fixpoints are precisely the spatial sublocales of $\mathbf{S}(L)$ on one side, and the sober subspaces of $\mathbf{pt}(L)$ on the other. It is also shown that for every frame L we have a commuting diagram, as follows, in the category of coframes.

$$\begin{array}{ccc}
\text{sp}[\mathbf{S}(L)] & \xrightarrow{\text{pt}(\cong)} & \text{sob}[\mathcal{P}(\text{pt}(L))] \\
\uparrow \text{sp} & & \downarrow \\
\mathbf{S}(L) & \xrightarrow{\text{pt}} & \mathcal{P}(\text{pt}(L))
\end{array}$$

The top horizontal arrow is always an isomorphism. It is then meaningful to ask three questions.

- (1) Firstly we can ask ourselves when is it that the left vertical arrow is an isomorphism. The frames for which this holds are the *totally spatial* frames, i.e. those such that all their sublocales are spatial. These are precisely the frames L such that $\mathbf{S}(L)^{op}$ is spatial, and they have also been characterized in several other ways in [8]. By looking at the diagram, we see that the totally spatial frames are exactly those such that there is a perfect correspondence between the sublocales of L and the sober subspaces of $\text{pt}(L)$.
- (2) We may also ask when it is that the right vertical arrow is an isomorphism. The frames for which this holds are exactly those such that $\text{pt}(L)$ is a T_D -space, as shown in [15]. By looking at the diagram, we see that these are exactly the frames for which there is a perfect correspondence between the spatial sublocales of L and the subspaces of $\text{pt}(L)$.
- (3) We may ask ourselves when it is that both vertical arrows are isomorphisms. These frames are the spatial frames such that $\text{pt}(L)$ is a *scattered* space (see [14]). These are the frames such that there is a perfect correspondence between sublocales of L and subspaces of $\text{pt}(L)$.

We want to tackle the analogues of these three questions in the setting of the T_D -duality. Let us start from the main adjunction.

Proposition 3.1. *There is an adjunction*

$$\begin{array}{ccc}
& \mathfrak{M} & \\
& \curvearrowright & \\
\mathcal{P}(\text{pt}_D(L)) & \perp & \mathbf{S}_D(L) \\
& \curvearrowleft & \\
& \text{pt}_D &
\end{array}$$

The fixpoints of $\mathfrak{M} \circ \text{pt}_D$ are the T_D -spatial D -sublocales, and $\text{pt}_D \circ \mathfrak{M}$ is the identity.

Proof: The map \mathbf{pt}_D is well-defined because if $S \in \mathbf{S}_D(L)$ then $\mathbf{pt}_D(S) \subseteq \mathbf{pt}_D(L)$. Moreover, clearly both \mathbf{pt}_D and \mathfrak{M} are monotone. For the adjunction, we have to check that $\mathfrak{M}(Y) \subseteq S$ if and only if $Y \subseteq \mathbf{pt}_D(S)$ for each $S \in \mathbf{S}_D(L)$ and $Y \subseteq \mathbf{pt}_D(L)$. The “only if” implication is trivial because $Y \subseteq \mathfrak{M}(Y)$, and the converse clearly follows from the fact that sublocales are closed under meets. Now, By Lemma 1.3, the fixpoints of $\mathfrak{M} \circ \mathbf{pt}_D$ are precisely those D -sublocales which are T_D -spatial. The only task remaining is to show that $\mathbf{pt}_D \circ \mathfrak{M}$ is the identity, so let $Y \subseteq \mathbf{pt}_D(L)$ and $p \in \mathbf{pt}_D(\mathfrak{M}(Y))$. Since $p \in \mathbf{pt}_D(\mathfrak{M}(Y)) \subseteq \mathfrak{M}(Y)$, there is a family $\{y_i\}_{i \in I} \subseteq Y$ with $p = \bigwedge y_i$. But p is a covered prime in Y so there is an $i_0 \in I$ with $p = y_{i_0} \in Y$. The other inclusion follows by adjunction. \blacksquare

Lemma 3.2. *For a frame L , the sublocale of L associated with its T_D -spatialization is $\mathfrak{M}(\mathbf{pt}_D(L))$.*

Proof: By the universal property of the T_D -spatialization, this is the largest D -sublocale of L such that it is T_D -spatial. Suppose that $S \subseteq L$ is a T_D -spatial D -sublocale. Because S is a D -sublocale, we have $\mathbf{pt}_D(S) \subseteq \mathbf{pt}_D(L)$, and because \mathfrak{M} is monotone we also have $\mathfrak{M}(\mathbf{pt}_D(S)) \subseteq \mathfrak{M}(\mathbf{pt}_D(L))$. Thus, by T_D -spatiality of S , we have proven that $S \subseteq \mathfrak{M}(\mathbf{pt}_D(L))$. \blacksquare

3.1. Local and global T_D -spatialization. It follows from Lemma 3.2 that the map $\mathfrak{M} \circ \mathbf{pt}_D: \mathbf{S}_D(L) \rightarrow \mathbf{S}_D(L)$ assigns to each D -sublocale its T_D -spatialization. From now on, we will call this map $\mathbf{sp}_D: \mathbf{S}_D(L) \rightarrow \mathbf{S}_D(L)$. Let us see what more we can learn about this operator.

Lemma 3.3. *If L is a complete lattice and $\iota: L \rightarrow L$ is an interior operators, then $\iota[L]$ is a complete lattice where the joins in are computed as in L , and the meets are computed as $\bigwedge^\iota \iota(x_i) = \iota(\bigwedge_i \iota(x_i))$.*

Proof: Suppose that L is a complete lattice and that $\iota: L \rightarrow L$ is an interior operator. Suppose that $x_i \in L$. We claim that $\bigvee_i \iota(x_i)$ is the join of $\{\iota(x_i) \mid i \in I\}$ in $\iota[L]$. first, we show that $\iota(\bigvee_i \iota(x_i)) = \bigvee_i \iota(x_i)$. One inequality holds since ι is deflationary. For the other, we notice that $\iota(x_i) \leq \bigvee_i \iota(x_i)$. By monotonicity and idempotence of ι , it follows that $\iota(x_i) \leq \iota(\bigvee_i \iota(x_i))$. The desired result follows. Let us show that $\bigvee_i^\iota \iota(x_i) = \iota(\bigvee_i \iota(x_i))$. On the one hand we showed before that $\iota(x_i) \leq \iota(\bigvee_i \iota(x_i))$. On the other hand, if we have $\iota(x_i) \leq \iota(y)$ for some $y \in L$ we also have that $\iota(\bigvee_i \iota(x_i)) \leq \iota(y)$, by monotonicity and idempotence of ι . Then, indeed $\bigvee_i^\iota \iota(x_i) = \iota(\bigvee_i \iota(x_i))$.

Now let us show that $\bigwedge_i \iota(x_i) = \iota(\bigwedge_i \iota(x_i))$. On the one hand we have $\bigwedge_i \iota(x_i) \leq \iota(x_i)$ and so $\iota(\bigwedge_i \iota(x_i)) \leq \iota(x_i)$ by monotonicity and idempotence. On the other hand if $\iota(y) \leq \iota(x_i)$ then we also have $\iota(y) \leq \bigwedge_i \iota(x_i)$ and so $\iota(y) \leq \iota(\bigwedge_i \iota(x_i))$ by monotonicity and idempotence. ■

Lemma 3.4. *If L is a complete lattice and $\iota: L \rightarrow L$ an interior operator on it, the surjection $\iota: L \twoheadrightarrow \iota[L]$ preserves arbitrary meets. If $\iota: L \rightarrow L$ preserves finite joins, this surjection preserves finite joins, too.*

Proof: Suppose that $\iota: L \rightarrow L$ is an interior operator. Suppose that $x_i \in L$. We first claim that $\iota(\bigwedge_i x_i) = \iota(\bigwedge_i \iota(x_i))$. We have that $\iota(x_i) \leq x_i$ since ι is deflationary, and thus $\bigwedge_i \iota(x_i) \leq \bigwedge_i x_i$, from which we deduce $\iota(\bigwedge_i \iota(x_i)) \leq \iota(\bigwedge_i x_i)$ by monotonicity of ι . For the reverse inequality, we start from the inequality $\bigwedge_i x_i \leq x_i$. By monotonicity of ι we deduce $\iota(\bigwedge_i x_i) \leq \bigwedge_i \iota(x_i)$. Finally, by monotonicity and idempotence, we deduce $\iota(\bigwedge_i x_i) \leq \iota(\bigwedge_i \iota(x_i))$. We have shown that $\iota(\bigwedge_i x_i) = \iota(\bigwedge_i \iota(x_i)) = \bigwedge_i \iota(x_i)$, where the last equality holds by Lemma 3.3. Suppose now that $\iota: L \rightarrow L$ preserves finite joins. For $x, y \in L$ we have that $\iota(x \vee y) = \iota(x) \vee \iota(y) = \iota(x) \vee^{\iota} \iota(y)$, where the last equality holds by Lemma 3.3. ■

Proposition 3.5. *For a frame L , the map $\text{sp}_D: \mathbf{S}_D(L) \rightarrow \mathbf{S}_D(L)$ is an interior operator which preserves finite joins.*

Proof: The map is monotone since it is the composition of the monotone maps $\text{pt}_D: \mathbf{S}_D(L) \rightarrow \mathcal{P}(\text{pt}_D(L))$ and $\mathfrak{M}: \mathcal{P}(\text{pt}_D(L)) \rightarrow \mathbf{S}_D(L)$. The map is clearly deflationary. Idempotence follows from the adjunction identity $\mathfrak{M} \circ \text{pt}_D \circ \mathfrak{M} = \mathfrak{M}$. For preservation of finite joins, suppose that we have a finite collection $S_m \in \mathbf{S}_D(L)$. Since these are D -sublocales and by primality of the elements in $\text{pt}_D(L)$, we have that $\text{pt}_D(S_1 \vee \dots \vee S_n) = \text{pt}_D(L) \cap (S_1 \vee \dots \vee S_n) = \text{pt}_D(L) \cap (S_1 \cup \dots \cup S_n) = (\text{pt}_D(L) \cap S_1) \cup \dots \cup (\text{pt}_D(L) \cap S_n) = \text{pt}_D(S_1) \cup \dots \cup \text{pt}_D(S_n)$. We have shown that $\text{pt}_D: \mathbf{S}_D(L) \rightarrow \mathcal{P}(\text{pt}_D(L))$ preserves finite joins. As \mathfrak{M} is a left adjoint, it preserves all joins, in particular finite ones. Thus the composition $\mathfrak{M} \circ \text{pt}_D: \mathbf{S}_D(L) \rightarrow \mathbf{S}_D(L)$ preserves finite joins. ■

Corollary 3.6. *Suppose that L is a frame. The map $\text{sp}_D: \mathbf{S}_D(L) \twoheadrightarrow \text{sp}_D[\mathbf{S}_D(L)]$ is a coframe surjection whose codomain is the ordered collection of the T_D -spatial D -sublocales of L .*

Proof: The fact that we have a coframe surjection follows from Lemma 3.4 and from Proposition 3.5. The fact that the codomain is the ordered collection of the T_D -spatial D -sublocales follows from Corollary 3.1. ■

Remark 3.7. The operator $\mathbf{sp}_D: \mathbf{S}_D(L) \rightarrow \mathbf{S}_D(L)$ actually preserves arbitrary joins, too. In the proof of Lemma 3.5, we have shown that for a finite collection S_m of sublocales we have that $\mathbf{pt}_D(S_1 \vee \dots \vee S_n) = \mathbf{pt}_D(S_1) \cup \dots \cup \mathbf{ppt}_D(S_n)$. Using coveredness of primes, this argument can be strengthened to show that $\mathbf{pt}_D(\bigvee_i S_i) = \bigcup_i \mathbf{pt}_D(S_i)$ for an arbitrary collection S_i of sublocales. This tells us that in the frame $\mathbf{S}_D(L)^{op}$ the congruence associated with the sublocale $\mathbf{sp}_D[\mathbf{S}_D(L)]$ is a *complete* congruence (see [5]).

We then have a local T_D -spatialization surjection $\mathbf{sp}_D: \mathbf{S}_D(L) \rightarrow \mathbf{sp}_D[\mathbf{S}_D(L)]$. As $\mathbf{S}_D(L)^{op}$ is a frame, it has its own T_D -spatialization sublocale; its global T_D -spatialization subcolocale, so to speak. We want to know how the local T_D -spatialization subcolocale $\mathbf{sp}[\mathbf{S}_D(L)] \subseteq \mathbf{S}_D(L)$ relates to the global one.

Lemma 3.8. *For a frame L , a D -sublocale $S \subseteq L$ is T_D -spatial if and only if $S = \bigvee \{ \mathbf{b}(p) \mid p \in \mathbf{pt}_D(S) \}$.*

Proof: By Corollary 3.1, a D -sublocale $S \subseteq L$ is T_D -spatial if and only if $\mathfrak{M}(\mathbf{pt}_D(S)) = S$. But $\mathfrak{M}(\mathbf{pt}_D(S)) = \bigvee \{ \mathbf{b}(p) \mid p \in \mathbf{pt}_D(S) \}$ and so we are done. \blacksquare

Lemma 3.9. *The covered prime elements of $\mathbf{S}_D(L)^{op}$ are exactly the sublocales of the form $\mathbf{b}(p)$ with $p \in \mathbf{pt}_D(L)$.*

Proof: Suppose that $p \in \mathbf{pt}_D(L)$. We claim that $\mathbf{b}(p) \in \mathbf{S}_D(L)^{op}$ is a covered prime. Indeed, we observe that if we have $\mathbf{b}(p) = \bigvee_i S_i$ we clearly must have $S_j = \{1, p\}$ for some $j \in I$.

For the converse, suppose that $S \in \mathbf{S}_D(L)$ and that $S \neq \mathbf{b}(p)$ for each $p \in \mathbf{pt}_D(L)$. Since for each prime $p \in \mathbf{pt}(L)$ we have that $\mathbf{b}(p) \in \mathbf{S}_D(L)$ if and only if p is covered, we must have that S cannot be a two-element sublocale, and so it contains at least three elements. Suppose, then, that $\{x, y, 1\}$ are pairwise distinct and that $\{x, y\} \subseteq S$. Suppose, in particular, that $x \not\leq y$. First, recall that all open and all closed sublocales of L are elements of $\mathbf{S}_D(L)$. We claim that $S \subseteq \mathbf{o}(x) \vee \mathbf{c}(x)$ is a nontrivial decomposition of S . We have that $x \rightarrow x = 1 \neq x$, and so $x \notin \mathbf{o}(x)$, hence $S \not\subseteq \mathbf{o}(x)$. On the other hand, we have that $y \notin \mathbf{c}(x)$ and so $S \not\subseteq \mathbf{c}(x)$. But since $\mathbf{c}(x) \vee \mathbf{o}(x) = 1$ we have $S \subseteq \mathbf{o}(x) \vee \mathbf{c}(x)$. \blacksquare

Observation 3.10. *In the proof above, starting from the assumption that a sublocale $S \in \mathbf{S}_D(L)$ is not of the form $\mathbf{b}(p)$ for a covered prime $p \in L$ we have shown that S is not a prime. This means that in $\mathbf{S}_D(L)$ if an element is not a covered prime it is not a prime.*

We have then proved the following.

Proposition 3.11. *In the frame $\mathbf{S}_D(L)^{op}$ all primes are covered. So, the frame $\mathbf{S}_D(L)^{op}$ is T_D -spatial if and only if it is spatial.*

Proof: This follows from Proposition 3.9 and from Observation 3.10. \blacksquare

We are ready to show that the local and the global spatialization of $\mathbf{S}_D(L)$ coincide.

Proposition 3.12. *For any frame L we have that $\mathfrak{M}(\mathbf{pt}_D(\mathbf{S}_D(L))) = \mathbf{sp}_D[\mathbf{S}_D(L)]$. In other words, global and local T_D -spatialization of $\mathbf{S}_D(L)$ coincide.*

Proof: By Lemma 3.9, we have that $\mathbf{pt}_D(\mathbf{S}_D(L)) = \{ \mathbf{b}(p) \mid p \in \mathbf{pt}_D(L) \}$. Hence the elements of $\mathfrak{M}(\mathbf{pt}_D(\mathbf{S}_D(L)))$ are exactly the joins of elements of the form $\mathbf{b}(p)$ for $p \in \mathbf{pt}_D(L)$. By Lemma 3.8, these are exactly the T_D -spatial D -sublocales of L . \blacksquare

Corollary 3.13. *For a frame L , we have that all its D -sublocales are T_D -spatial if and only if $\mathbf{S}_D(L)$ is T_D -spatial, and this holds if and only if $\mathbf{S}_D(L)$ is spatial.*

Proof: This follows from Propositions 3.12 and 3.11. \blacksquare

Remark 3.14. It should be noted that $\mathbf{sp}_D[\mathbf{S}_D(L)]$ is not only spatial, but it is also Boolean for any frame L . Indeed, it follows by virtue of the adjunction in Proposition 3.1 and the fact that $\mathbf{pt}_D \circ \mathfrak{M}$ is the identity that it is actually isomorphic to $\mathcal{P}(\mathbf{pt}_D(L))$. More precisely, it corresponds to the Boolean sublocale of $\mathbf{S}(L)^{op}$ determined by the element $\mathbf{sp}_D(L)$, i.e. we have $\mathbf{sp}_D[\mathbf{S}_D(L)] = \mathbf{b}(\mathbf{sp}_D(L))$ (to see this, observe that a Boolean sublocale S of a frame L always equals $\mathbf{b}(\bigwedge S)$, hence $\mathbf{sp}_D[\mathbf{S}_D(L)]$ is determined by the largest D -sublocale which is T_D -spatial, i.e. the T_D -spatialization of L).

We wish to stress that the behaviour of the operator $\mathbf{pt}_D \circ \mathfrak{M}: \mathcal{P}(\mathbf{pt}_D(L)) \rightarrow \mathcal{P}(\mathbf{pt}_D(L))$ exhibits a difference between the spatial-sober duality and the T_D -duality. In the classical duality (see [15]) we have an adjunction

$$\begin{array}{ccc} & \mathfrak{M} & \\ & \curvearrowright & \\ \mathcal{P}(\mathbf{pt}(L)) & \perp & \mathbf{S}(L) \\ & \curvearrowleft & \\ & \mathbf{pt} & \end{array}$$

in which the fixpoints of $\mathfrak{M} \circ \mathbf{pt}$ are precisely the spatial sublocales, and the fixpoints of $\mathbf{pt} \circ \mathfrak{M}$ are the sober subspaces of $\mathbf{pt}(L)$. None of the two

compositions is the identity. In particular, since not all subspaces of $\mathbf{pt}(L)$ are sober, this means that the map $\mathbf{pt}: \mathbf{S}(L) \rightarrow \mathcal{P}(\mathbf{pt}(L))$ will not in general be a surjection: some subspaces of $\mathbf{pt}(L)$ do not arise as spectra of sublocales of L . This accounts for part of the mismatch between subspaces and sublocales. In the T_D -duality, this does not happen. Since the operator $\mathbf{pt}_D \circ \mathfrak{M}$ is the identity, it means that all subspaces of $\mathbf{pt}_D(L)$ arise as spectra of D -sublocales of L . We shall see that this means that in the T_D -duality the mismatch between subspaces and sublocales is only given by the map $\mathbf{pt}_D: \mathbf{S}_D(L) \rightarrow \mathcal{P}(\mathbf{pt}_D(L))$ not being injective: in the T_D -duality we have more sublocales than subspaces.

Let us now look at the T_D analogue of the square in the beginning of this section.

$$\begin{array}{ccc}
 (\mathfrak{M} \circ \mathbf{pt}_D)[\mathbf{S}_D(L)] & \xrightarrow{\mathbf{pt}_D(\cong)} & (\mathbf{pt}_D \circ \mathfrak{M})[\mathcal{P}(\mathbf{pt}_D(L))] \\
 \mathfrak{M} \circ \mathbf{pt}_D \uparrow & & \downarrow = \\
 \mathbf{S}_D(L) & \xrightarrow{\mathbf{pt}_D} & \mathcal{P}(\mathbf{pt}_D(L))
 \end{array}$$

We know that the diagram commutes in the category of coframes as the composition $\mathbf{pt}_D \circ \mathfrak{M}$ is the identity and so indeed $\mathbf{pt}_D \circ \mathfrak{M} \circ \mathbf{pt}_D = \mathbf{pt}_D$. We observe that the right vertical arrow is always the identity, by Corollary 3.1. Thus, in the T_D -duality, we are left with only one of the three questions. We shall then characterize those frames L such that all the D -sublocales of L are T_D -spatial, and these coincide with the frames L such that there is a perfect correspondence between D -sublocales of L and subspaces of $\mathbf{pt}_D(L)$.

3.2. When do the D -sublocales perfectly represent the subspaces?

For a frame L and a meet $\bigwedge M$, we say that $m \in M$ is an *essential* element of the meet $\bigwedge M$ if we have $\bigwedge M \setminus \{m\} \neq \bigwedge M$. If L is a frame and $a \in L$ is the meet of the primes above it, we say that $p \in \mathbf{pt}(\uparrow a)$ is an *essential* prime of a if and only if $\bigwedge \mathbf{pt}(\uparrow a) \setminus \{p\} \neq a$. We say that $p \in \mathbf{pt}(\uparrow a)$ is an *absolutely essential* prime of a if $\bigwedge \mathbf{pt}(\uparrow a) \setminus \{p\} \neq a$. It is clear that absolute essentiality is a condition stronger than essentiality. The terminology “absolutely” essential is justified by the following result.

Lemma 3.15. *For a frame L and for $a \in L$ with $a = \bigwedge \text{pt}(\uparrow a)$ and for $p \in \text{pt}(\uparrow a)$, the following are equivalent.*

- (1) *The prime p is an absolutely essential prime of a ;*
- (2) *Whenever $a = \bigwedge P$ for $P \subseteq \text{pt}(L)$ we must have $p \in P$;*
- (3) *The prime p is weakly covered and it is an essential prime of a .*

Proof: Let L be a frame, and let $a \in L$ be an element such that it is the meet of the primes above it. Let $p \in \text{pt}(\uparrow a)$.

(1) \implies (2). Suppose that p is an absolutely essential prime of a , and that $P \subseteq \text{pt}(L)$ is such that $a = \bigwedge P$. We must have that $P \subseteq \text{pt}(\uparrow a)$ and so $P \setminus \{p\} \subseteq \text{pt}(\uparrow a) \setminus \{p\}$, from which we deduce that $\bigwedge \text{pt}(\uparrow a) \setminus \{p\} \leq \bigwedge P \setminus \{p\}$. From this fact, and by absolute essentiality of p , we deduce that $\bigwedge P \setminus \{p\} \not\leq a$. As $\bigwedge P \leq a$, we must have $p \in P$.

(2) \implies (3). Suppose that $p \in P$ whenever $\bigwedge P = a$ for $P \subseteq \text{pt}(L)$. In particular, we have that $\bigwedge \text{pt}(\uparrow a) \setminus \uparrow p \not\leq a$ as $p \notin \text{pt}(\uparrow a) \setminus \uparrow p$. Now suppose towards contradiction that p is not weakly covered, and so that $p = \bigwedge \text{pt}(\uparrow p) \setminus \{p\}$. We then have that $\bigwedge \{q \in \text{pt}(\uparrow a) \mid p \neq q\} = \bigwedge \{q \in \text{pt}(\uparrow a) \mid p < q\} \wedge \bigwedge \{q \in \text{pt}(\uparrow a) \mid p \not\leq q\} = p \wedge \bigwedge \text{pt}(\uparrow a) \setminus \uparrow p \leq a$. We have then contradicted (2).

(3) \implies (1). Suppose that p is an essential prime of a and that it is weakly covered. if we had $\bigwedge \text{pt}(\uparrow a) \setminus \uparrow p \leq p$ then we would have that $\bigwedge \text{pt}(\uparrow a) \setminus \uparrow p \leq q$ whenever $q \in \text{pt}(\uparrow a)$, contradicting essentiality of p . Then, we must have $\bigwedge \text{pt}(\uparrow a) \setminus \uparrow p \not\leq p$. We also have $\bigwedge \{q \in \text{pt}(\uparrow a) \mid p < q\} \not\leq p$ by weak coveredness of p . By primality of p , this means that $\bigwedge \text{pt}(\uparrow a) \setminus \uparrow p \wedge \bigwedge \{q \in \text{pt}(\uparrow a) \mid p < q\} = \bigwedge \text{pt}(\uparrow a) \setminus \{p\} \not\leq p$, and so $\bigwedge \text{pt}(\uparrow a) \setminus \{p\} \not\leq a$. Thus, p is an absolutely essential prime of a . \blacksquare

Thus, we have called this condition “absolute” essentiality as opposed to relative essentiality: an essential prime of a is only essential relative to the particular meet of primes $\bigwedge \text{pt}(\uparrow a) \setminus \{q \in \text{pt}(\uparrow a) \mid p < q\}$; whereas an absolutely essential prime is essential with respect to any meet of primes $\bigwedge P$ with $\bigwedge P = a$. It is crucial that for covered primes these two conditions collapse together.

Lemma 3.16. *If p is a covered prime and p is an essential prime of $a \in L$, then p is also an absolutely essential prime of a .*

Proof: If p is an essential prime of a and it is covered, in particular it is weakly covered and so it is absolutely essential, by Lemma 3.15. \blacksquare

In [15], the following is shown. This is a choice-free adaptation of a famous result of [8].

Proposition 3.17. *For a frame L , we have that L is totally spatial if and only if every element of L is the meet of its essential primes.*

Let us prove some general facts about essential and absolutely essential primes.

Lemma 3.18. *For a frame L and an element a which is the meet of the primes above it, we have that $p \in \mathbf{pt}(\uparrow a)$ is an essential prime of a if and only if it is a prime of $\mathbf{b}(a)$.*

Proof: Suppose that L is a frame and that $a \in L$ is the meet of the primes above it. Suppose first that p is an essential prime of a . We claim that $p \in \mathbf{b}(a)$ because we have

$$p = \bigwedge \mathbf{pt}(\uparrow a) \setminus \uparrow p \rightarrow a = \bigwedge \{ \bigwedge \mathbf{pt}(\uparrow a) \setminus \uparrow p \rightarrow q \mid q \in \mathbf{pt}(\uparrow a) \}.$$

The right hand side is a conjunction, in which every conjunct is either 1 or a prime above a . Then, this is equal to

$$\bigwedge \{ q \in \mathbf{pt}(\uparrow a) \mid \bigwedge \mathbf{pt}(\uparrow a) \setminus \uparrow p \not\leq q \}$$

As p is an essential prime, it is certainly in the set $\{ q \in \mathbf{pt}(\uparrow a) \mid \bigwedge \mathbf{pt}(\uparrow a) \setminus \uparrow p \not\leq q \}$. It must be the minimal element of this set, as for an element q to be in this set we need to have $p \leq q$. Then, indeed, $p = \bigwedge \mathbf{pt}(\uparrow a) \setminus \uparrow p \rightarrow a$ and so $p \in \mathbf{b}(a)$. For the converse, suppose that we have $p \in \mathbf{b}(a)$. Then, we have that $(p \rightarrow a) \rightarrow a = p$. This means that

$$p = \bigwedge \{ (p \rightarrow a) \rightarrow q \mid q \in \mathbf{pt}(\uparrow a) \}$$

The right hand side is again a conjunct of primes, and in particular it is

$$\bigwedge \{ q \in \mathbf{pt}(\uparrow a) \mid p \rightarrow a \not\leq q \}$$

Now, let us observe that $p \rightarrow a$ is exactly $\bigwedge \{ q \in \mathbf{pt}(\uparrow a) \mid p \not\leq q \} = \bigwedge \mathbf{pt}(\uparrow a) \setminus \uparrow p$. Then, the expression above equals

$$\bigwedge \{ q \in \mathbf{pt}(\uparrow a) \mid \bigwedge \mathbf{pt}(\uparrow a) \setminus \uparrow p \not\leq q \}.$$

Since this expression equals p , if we had $\bigwedge \mathbf{pt}(\uparrow a) \leq p$, we would also have $\bigwedge \mathbf{pt}(\uparrow a) \leq q$ for every element q of the set $\{ q \in \mathbf{pt}(\uparrow a) \mid \bigwedge \mathbf{pt}(\uparrow a) \setminus \uparrow p \not\leq q \}$,

as p is a lower bound for this set. This is a contradiction, by the membership condition of this set. The prime p must then be essential. ■

Lemma 3.19. *For a frame L and for $x, y \in L$ we have that $\mathfrak{b}(x \rightarrow y) = \mathfrak{o}(x) \cap \mathfrak{b}(y)$.*

Proof: Suppose that $z \in \mathfrak{b}(x \rightarrow y)$. Then, we have that $z = z' \rightarrow (x \rightarrow y)$ for some $z' \in L$. Then, $z = (z' \wedge x) \rightarrow y$ and so $z \in \mathfrak{b}(y)$. Additionally, we have $x \rightarrow z = x \rightarrow (z' \rightarrow (x \rightarrow y)) = (x \wedge z') \rightarrow (x \rightarrow y) = (x \wedge z') \rightarrow y = z' \rightarrow (x \rightarrow y) = z$. Then, $z \in \mathfrak{o}(x)$. For the converse, suppose that $z \in \mathfrak{o}(x) \cap \mathfrak{b}(y)$. Then, we have $x \rightarrow z = z$ and also that $z = z' \rightarrow y$ for some $z' \in L$. Then, $z = x \rightarrow (z' \rightarrow y) = x \wedge z' \rightarrow y = z' \rightarrow (x \rightarrow y)$, and so $z \in \mathfrak{b}(x \rightarrow y)$. ■

Lemma 3.20. *For a frame L , we have that it is spatial and every element other than 1 has an absolutely essential prime above it if and only if every element is the meet of the essential primes above it.*

Proof: If L is a frame in which every element is the meet of the essential primes above it, in particular every element $a \in L$ with $a \neq 1$ must have at least an essential prime above it. The frame is also spatial. For the other implication, suppose that L is spatial and that every element other than 1 has an absolutely essential prime above it. Let $a \in L$ and let $\mathbf{AbsEss}(a)$ be the collection of absolutely essential primes of a . Suppose towards contradiction that $\bigwedge \mathbf{AbsEss}(a) \not\leq a$. Then, $\bigwedge \mathbf{AbsEss}(a) \rightarrow a \neq 1$ and so there is an absolutely essential prime q of $\bigwedge \mathbf{AbsEss}(a) \rightarrow a$. This means that we have $q \in \mathfrak{o}(\bigwedge \mathbf{AbsEss}(a)) \cap \mathfrak{b}(a)$ (see Lemma 3.18 and 3.19) and that q is weakly covered. Since $q \in \mathfrak{o}(\bigwedge \mathbf{AbsEss}(a))$ we have $q \notin \mathfrak{c}(\bigwedge \mathbf{AbsEss}(a))$, and so $\bigwedge \mathbf{AbsEss}(a) \not\leq q$. Since $q \in \mathfrak{b}(a)$ and so it is an absolutely essential prime of a . This contradicts that $\bigwedge \mathbf{AbsEss}(a) \not\leq q$. ■

We are ready to prove the main theorems of this section.

Theorem 3.21. *For a frame L , the following are equivalent.*

- (1) *All sublocales of L are T_D -spatial;*
- (2) *All D -sublocales of L are T_D -spatial;*
- (3) *$\mathfrak{M} \circ \mathbf{pt}_D$ is the identity on $\mathbf{S}_D(L)$;*
- (4) *There is an isomorphism $\mathbf{S}_D(L) \cong \mathcal{P}(\mathbf{pt}_D(L))$;*
- (5) *$\mathbf{S}_D(L)$ is spatial;*
- (6) *$\mathbf{S}_D(L)$ is spatial and Boolean (i.e. a complete and atomic Boolean algebra);*

(7) $\mathbf{S}_D(L) = \mathbf{S}_b(L)$ and L is T_D -spatial.

(8) Every nonzero sublocale of L contains a covered prime in itself.

Proof: (1) obviously implies (2).

(2) is equivalent to (3). This holds by Lemma 3.2

(3) \implies (4). The isomorphism is obtained by composing $\mathfrak{M} \circ \mathbf{pt}_D$ and the isomorphism $(\mathfrak{M} \circ \mathbf{pt}_D)[\mathbf{S}_D(L)] \rightarrow (\mathbf{pt}_D \circ \mathfrak{M})[\mathcal{P}(\mathbf{pt}_D(L))]$.

(4) \implies (5). This is clear.

(5) \implies (6). This implication follows from Corollary 3.13 and Remark 3.14.

(6) \implies (7). Suppose that $\mathbf{S}_D(L)$ is spatial and Boolean. If $\mathbf{S}_D(L)$ is Boolean, since it is also dense, it must coincide with the unique Boolean dense sublocale of $\mathbf{S}(L)$, namely $\mathbf{S}_b(L)$. Now, if $\mathbf{S}_D(L) = \mathbf{S}_b(L)$ is also spatial, by Theorem 1.4 we conclude that L is T_D -spatial.

(7) \implies (8). Notice that every non-smooth sublocale has a covered prime in itself (otherwise, we would have a non-smooth sublocale S with $\mathbf{pt}_D(S) = \emptyset$, then S is obviously a D -sublocale so by assumption it has to be smooth, contradiction). But L being T_D -spatial, say $L = \Omega(X)$ with X a T_D -space, every (nonzero) smooth sublocale of it is induced by a subspace $A \subseteq X$ and hence it contains a covered prime of L , in particular a covered prime in itself. Hence every nonzero sublocale contains a covered prime in itself.

(8) \implies (1). Clearly, the condition (8) is hereditary with respect to any sublocale. Hence, it suffices to show that L is T_D -spatial. If for all complemented sublocale $C \subseteq L$ such that $\mathbf{sp}_D(L) \subseteq C$ one has $C = L$, then by zero-dimensionality of $\mathbf{S}(L)^{op}$, it follows that $L = \mathbf{sp}_D(L)$ is T_D -spatial. Hence assume by way of contradiction that there is a complemented sublocale $C \subseteq L$ such that $C \neq L$ and $\mathbf{sp}_D(L) \subseteq C$. Since $C \neq L$, then $C^c \neq 0$, so by assumption there is a $p \in \mathbf{pt}_D(C^c)$ with $\mathbf{b}(p) \subseteq C^c$. But C^c is complemented; in particular it is a D -sublocale, hence $p \in \mathbf{pt}_D(C^c) \subseteq \mathbf{pt}_D(L)$. But then $\mathbf{b}(p) \subseteq \mathbf{sp}_D(L) \subseteq C$, whence $\mathbf{b}(p) \subseteq C \cap C^c = 0$, contradiction. \blacksquare

We now move on to looking at the condition of a frame having all its sublocales *strongly* T_D -spatial. This condition is indeed strictly stronger than those occurring in Theorem 3.21; as the following example based on [3, Example 7.8] shows.

Consider the totally ordered set of the natural numbers with the Alexandroff topology (i.e. the topology consisting of upsets), that is, $\Omega(\mathbb{N}) = \{\emptyset\} \cup$

$\{\uparrow n \mid n \in \mathbb{N}\}$. Observe that for any $n > 1$ the element $\uparrow n = \mathbb{N} - \overline{\{n-1\}}$ is a covered prime in $\Omega(\mathbb{N})$ and \emptyset is a prime which is not covered. Hence condition (1) in Theorem 3.22 is not satisfied. Moreover, it was pointed out in [3] that $\Omega(\mathbb{N})$ is totally spatial. In fact, one has that every sublocale is T_D -spatial. In order to see this, let S be a nonzero sublocale. If $S = \{\mathbb{N}, \emptyset\}$, obviously S contains a covered prime in itself (namely, \emptyset). Otherwise, S contains an element of the form $\uparrow n$ with $n > 1$, but all such elements are covered in $\Omega(\mathbb{N})$, and in particular they are covered in S . Thus condition (8) in Theorem 3.21 holds.

The fact that T_D -spatiality of D -sublocales implies T_D -spatiality of *all* sublocales may seem somewhat surprising, specially because the conditions of Theorem 3.21 do not imply that every sublocale is a D -sublocale (indeed, by Theorem 4.1 below, that would imply the conditions of Theorem 3.22, which we have just seen to be a strictly stronger condition).

Theorem 3.22. *For a frame L , the following are equivalent.*

- (1) *The frame L is totally spatial and all its primes are covered;*
- (2) *The frame L is totally spatial and strongly T_D -spatial;*
- (3) *All sublocales of L are strongly T_D -spatial;*
- (4) *There is an isomorphism $\mathbf{S}(L) \cong \mathcal{P}(\mathbf{pt}_D(L))$;*
- (5) *The coframe $\mathbf{S}(L)$ is spatial and Boolean;*
- (6) *Every element of L is the meet of the covered essential primes above it;*
- (7) *Every element of L is the meet of the covered absolutely essential primes above it;*
- (8) *The frame L is spatial and every element other than 1 has a covered absolutely essential prime above it;*
- (9) *Every nonzero sublocale of L contains a covered prime of L .*

Proof: (1) \implies (2). If L is totally spatial, then in particular it is spatial; if additionally all its primes are covered the frame is also strongly T_D -spatial.

(2) \implies (3). If L is totally spatial, then every sublocale is spatial, so every element of S is a meet of primes for each $S \in \mathbf{S}(L)$. But every $p \in \mathbf{pt}(S)$ is a covered prime in L , and in particular in S , so every element of S is a meet of covered primes in S . Now use Lemma 1.3.

(3) \implies (4). If all sublocales of L are strongly T_D -spatial, in particular all of the primes of L are covered and so we have $\mathbf{S}_D(L) = \mathbf{S}(L)$. Furthermore, all of the sublocales being strongly T_D -spatial means, by commutativity of the main

square of this section, that we have an isomorphism $\mathbf{S}_D(L) \cong \mathcal{P}(\text{pt}_D(L))$.

(4) \implies (5). This is clear.

(5) \implies (6). If $\mathbf{S}(L)$ is spatial, the frame L is totally spatial, and so every element is the meet of its essential primes by Lemma 3.17. Furthermore, if $\mathbf{S}(L)$ is Boolean, in particular all two-element sublocales are complemented, and so every prime is covered.

(6) \implies (7). This follows from Lemma 3.15 (if a prime is essential for one element and it is covered, it is also absolutely essential).

(7) \implies (8). If all elements are the meets of the absolutely essential primes above it, then in particular this must be true for elements other than 1, and so their corresponding meet of absolutely essential primes must be nonempty.

(8) and (9) are equivalent. If (8) holds, then by Lemma 3.18 all nonzero Boolean sublocales have at least a point and since all sublocales are unions of Boolean sublocales this also implies that all nonzero sublocales have at least a point. For the converse, suppose that every nonzero sublocale of L has a covered prime. In particular, every Boolean sublocale $\mathfrak{b}(a)$ with $a \neq 1$ has a prime $p \in \mathfrak{b}(a)$, and this by Lemma 3.18 means that a has an essential prime which is covered. But since p is covered, by Lemma 3.16 this implies that it is also an absolutely essential prime of a .

(8) \implies (1). Suppose that L is spatial, and that every element other than 1 has a covered absolutely essential prime above it. By Lemma 3.20, we know that the frame is totally spatial, as every element is the meet of the absolutely essential primes above it and so, a fortiori, it is the meet of the essential primes above it. Let us show that all primes of L are covered. Every prime p has an absolutely essential prime and so, in particular, the meet $\bigwedge\{p\}$ has an essential prime, which must be p itself. Furthermore, our assumption also tells us that the absolutely essential prime p must be covered. ■

4. The relation between $\mathbf{S}_D(L)$ and $\text{sp}[\mathbf{S}(L)]$

We begin this section with a non-spatial version of Theorem 1.7, that is, we explore locales in which every prime is covered. Note that this is the T_D analogue of the point-free notion of T_1 locale introduced by Rosický and Šmarda [13].

Theorem 4.1. *For a frame L , the following are equivalent.*

- (1) *All primes of L are covered;*
- (2) $\text{sp}[\mathbf{S}(L)] \subseteq \mathbf{S}_b(L)$;
- (3) $\mathbf{S}_D(L) = \mathbf{S}(L)$;
- (4) $\mathbf{S}_D(L)$ *is closed under arbitrary intersections in $\mathbf{S}(L)$;*
- (5) $\text{sp}[\mathbf{S}(L)] \subseteq \mathbf{S}_D(L)$.

Proof: Suppose that (1) holds. Then, (2) holds by Corollary 1.2. Suppose that (2) holds and let S an arbitrary sublocale. If $p \in \text{pt}_D(S)$, then in particular $p \in \text{pt}(S)$ and so $\mathfrak{b}(p) \in \text{sp}[\mathbf{S}(L)] \subseteq \mathbf{S}_b(L)$. By Lemma 1.1, one has that p is covered in L , and so $S \in \mathbf{S}_D(L)$. That (3) implies (4) is obvious and (4) implies (3) because every sublocale is an intersection of complemented sublocales (and hence of D -sublocales). Assume now that (3) holds and let p be a prime. Then $\mathfrak{b}(p) \in \text{sp}[\mathbf{S}(L)] \subseteq \mathbf{S}_D(L)$ and so $p \in \text{pt}_D(\mathfrak{b}(p)) \subseteq \text{pt}_D(L)$. Hence (1) follows. \blacksquare

We can use $\mathbf{S}_D(L)$ for characterizing a number of well-known properties for a frame L . For example we have the following (but see also Theorem 3.21 or Theorem 4.1).

Proposition 4.2. *For a frame L , the following are equivalent:*

- (1) *L is totally spatial;*
- (2) $\mathbf{S}_D(L) \subseteq \text{sp}[\mathbf{S}(L)]$.

Proof: (1) \implies (2). This is trivial since if L is totally spatial then $\text{sp}[\mathbf{S}(L)] = \mathbf{S}(L)$.

(2) \implies (1). By the observation preceding the proposition, every pointless sublocale of L is in $\mathbf{S}_D(L)$, and so by assumption, every pointless sublocale is spatial. Thus L has no nontrivial pointless sublocales, i.e. every nontrivial sublocale of L has a point. By [8, p. 269], it follows that L is totally spatial. \blacksquare

Next we explore the inclusion $\mathbf{S}_D(L) \subseteq \mathbf{S}_b(L)$ (note that the inclusion is actually equivalent to the equality $\mathbf{S}_D(L) = \mathbf{S}_b(L)$, since every frame has a unique Boolean and dense sublocale). We first give a necessary condition for it:

Proposition 4.3. *If $\mathbf{S}_D(L) \subseteq \mathbf{S}_b(L)$ then $\text{sp}(L)$ is totally spatial.*

Proof: Let S be a pointless sublocale contained in $\text{sp}(L)$. Then $S \in \mathbf{S}_D(L) \subseteq \mathbf{S}_b(L)$. It is easy to show that one always has $\mathbf{S}_b(\text{sp}(L)) \supseteq \downarrow^{\mathbf{S}_b(L)} \text{sp}(L)$. It

follows that $S \in \mathbf{S}_b(\mathbf{sp}(L))$. But $\mathbf{sp}(L)$ is spatial and so $S \in \mathbf{S}_b(\mathbf{sp}(L)) \subseteq \mathbf{sp}[\mathbf{S}(\mathbf{sp}(L))]$, but a spatial pointless sublocale must be trivial. Hence $\mathbf{sp}(L)$ does not contain any nontrivial pointless sublocales, which by [8, p. 269] implies that $\mathbf{sp}(L)$ is totally spatial. ■

We now characterize frames for which the inclusion $\mathbf{S}_D(L) \subseteq \mathbf{S}_b(L)$ holds. Despite the fact that we have not been able to find a clear geometric condition as those occurring in other results of the paper (spatiality, total spatiality, scatteredness, T_D -spatiality, . . .), condition (2) below indeed tells us something: the assumption in (1) (i.e. that a given sublocale S is a D -sublocale) is relative to L , whereas the assumption in (2) is absolute, in the sense that it does not depend on the ambient frame L :

Proposition 4.4. *For a frame L , the following are equivalent:*

- (1) $\mathbf{S}_D(L) \subseteq \mathbf{S}_b(L)$ (i.e. every D -sublocale is smooth);
- (2) Sublocales of L without covered primes in themselves are smooth in L .

Proof: (1) \implies (2). Let S be a sublocale such that $\mathbf{pt}_D(S) = \emptyset$. Then $S \in \mathbf{S}_D(L) \subseteq \mathbf{S}_b(L)$ so it is smooth.

(2) \implies (1). Let $S \in \mathbf{S}_D(L)$, and consider the decomposition $S = \mathbf{sp}_D(S) \vee (S \setminus \mathbf{sp}_D(S))$. Observe that by Proposition 2.3, $S \setminus \mathbf{sp}_D(S)$ is a D -sublocale as well, i.e. $\mathbf{pt}_D(S \setminus \mathbf{sp}_D(S)) \subseteq \mathbf{pt}_D(L)$. Assume that there exists some $p \in \mathbf{pt}_D(S \setminus \mathbf{sp}_D(S))$. Then $p \in \mathbf{pt}_D(L) \cap S \subseteq \mathbf{pt}_D(S)$ and hence $\mathbf{b}(p) \subseteq \mathbf{sp}_D(S)$. Therefore, $\mathbf{b}(p) \subseteq S \setminus \mathbf{sp}_D(S) \subseteq S \setminus \mathbf{b}(p) \subseteq L \setminus \mathbf{b}(p) = \mathbf{b}(p)^c$, which yields a contradiction. Hence $\mathbf{pt}_D(S \setminus \mathbf{sp}_D(S)) = \emptyset$ and so $S \setminus \mathbf{sp}_D(S)$ is smooth in L . Moreover, $\mathbf{sp}_D(S) = \bigvee_{p \in \mathbf{pt}_D(S)} \mathbf{b}(p)$ is also a smooth sublocale because $\mathbf{pt}_D(S) \subseteq \mathbf{pt}_D(L)$ and Lemma 1.1. Hence $S = \mathbf{sp}_D(S) \vee (S \setminus \mathbf{sp}_D(S))$ is smooth, as it is a join of smooth sublocales. ■

By analogy with the classical spectrum, frames L for which $\mathbf{S}_D(L) \subseteq \mathbf{S}_b(L)$ holds are going to be called *D -scattered*.

5. Adding $\mathbf{S}_c(L)$ to the picture

First, let us connect the frame of joins of closed sublocales with the spatialization sublocale of $\mathbf{S}(L)$.

Lemma 5.1. *A frame L is spatial if and only if $\mathbf{S}_c(L) \subseteq \mathbf{sp}[\mathbf{S}(L)]$.*

Proof: Suppose that L is spatial. For a closed sublocale $\uparrow a \subseteq L$, suppose that we have $b \in \uparrow a$. By spatiality, $b = \bigwedge \mathbf{pt}(\uparrow b)$, but since all primes above

b are also above a , this is the same as $\bigwedge(\text{pt}(\uparrow a) \cap \uparrow a) = \bigwedge\{p \in \uparrow a \mid b \leq p\}$. Then, $\uparrow a \subseteq \mathfrak{M}(\text{pt}(\uparrow a))$, and this means that $\uparrow a$ is spatial. Conversely, if $\mathbf{S}_c(L) \subseteq \text{sp}[\mathbf{S}(L)]$ we have that every closed sublocale is spatial, and in particular L itself is. ■

Lemma 5.2. *A frame L is such that all its primes are maximal if and only if $\text{sp}[\mathbf{S}(L)] \subseteq \mathbf{S}_c(L)$.*

Proof: Suppose that for a frame L all its primes are maximal. This means that for every prime $p \in \text{pt}(L)$ we have that $\mathfrak{b}(p)$ is closed, and since the elements of $\text{sp}[\mathbf{S}(L)]$ are all joins of elements of the form $\mathfrak{b}(p)$, we have $\text{sp}[\mathbf{S}(L)] \subseteq \mathbf{S}_c(L)$. Conversely, if we have a prime $p \in L$ which is not maximal, say $p < x < 1$, we have that $\mathfrak{b}(p) \neq \mathfrak{c}(p)$, and since $\mathfrak{b}(p)$ does not equal its closure it is not closed. Then, $\text{sp}[\mathbf{S}(L)] \not\subseteq \mathbf{S}_c(L)$. ■

Secondly, we look at the relation between $\mathbf{S}_c(L)$ and $\mathbf{S}_D(L)$. First, we observe that we always have $\mathbf{S}_c(L) \subseteq \mathbf{S}_D(L)$ (we know that all closed sublocales are D -sublocales, and the collection of D -sublocales is closed under joins).

Proposition 5.3. *For a frame L , we have that $\mathbf{S}_D(L) \subseteq \mathbf{S}_c(L)$ if and only if L is subfit and D -scattered.*

Proof: Suppose that for a frame L we have $\mathbf{S}_D(L) \subseteq \mathbf{S}_c(L)$. We have that all closed sublocales are complemented, and so $\mathbf{S}_c(L) \subseteq \mathbf{S}_b(L)$. We then have that $\mathbf{S}_D(L) \subseteq \mathbf{S}_b(L)$, which means that L is D -scattered. Furthermore, as this implies also that $\mathbf{S}_D(L) = \mathbf{S}_b(L)$, we also have that $\mathbf{S}_c(L) = \mathbf{S}_b(L)$, which means that L is subfit (see [10]). Conversely, suppose that L is D -scattered and subfit. Since L is subfit, we have that $\mathbf{S}_c(L) = \mathbf{S}_b(L)$ (see [10]), and since L is D -scattered, we have that $\mathbf{S}_D(L) = \mathbf{S}_b(L)$. Hence, $\mathbf{S}_D(L) = \mathbf{S}_c(L)$. ■

References

- [1] ARRIETA, I. On joins of complemented sublocales, 2020. Preprint available at “<http://www.mat.uc.pt/preprints/ps/p2023.pdf>”.
- [2] AULL, C. E., AND THRON, W. J. Separation Axioms Between T_0 and T_1 . *Indagationes Mathematicae* 24 (1963), 26–37.
- [3] ÁVILA, F., BEZHANISHVILI, G., MORANDI, P., AND ZALDÍVAR, A. When is the frame of nuclei spatial: A new approach. *Journal of Pure and Applied Algebra* 224 (2020), 1–20.
- [4] BANASCHEWSKI, B., AND PULTR, A. Pointfree Aspects of the T_D Axiom of Classical Topology. *Quaestiones Mathematicae* 33 (2010), 369–385.
- [5] CLEMENTINO, M. M., PICADO, J., AND PULTR, A. The Other Closure and Complete Sublocales. *Applied Categorical Structures* 26 (2018), 891–908.
- [6] FERREIRA, M. J., PICADO, J., AND PINTO, S. Remainders in pointfree topology. *Topology and its Applications* 245 (2018), 21–45.

- [7] JOHNSTONE, P. T. *Stone Spaces*, vol. 3 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, 1982.
- [8] NIEFIELD, S. B., AND ROSENTHAL, K. I. Spatial sublocales and essential primes. *Topology and its Applications* 26 (1987), 263–269.
- [9] PICADO, J., AND PULTR, A. *Frames and Locales: Topology without points*. Springer-Birkhäuser Basel, 2011.
- [10] PICADO, J., AND PULTR, A. Axiom T_D and the Simmons sublocale theorem. *Commentationes Mathematicae Universitatis Carolinae* 60 (2019), 541–551.
- [11] PICADO, J., PULTR, A., AND TOZZI, A. Joins of closed sublocales. *Houston Journal of Mathematics* 45 (2019), 21–38.
- [12] PULTR, A., AND TOZZI, A. Separation axioms and frame representation of some topological facts. *Applied Categorical Structures* 2 (1994), 107–118.
- [13] ROSICKÝ, J., AND ŠMARDA, B. T_1 -locales. *Mathematical Proceedings of the Cambridge Philosophical Society* 98 (1985), 81–86.
- [14] SIMMONS, H. Spaces with Boolean assemblies. *Colloquium Mathematicae* 43 (1980), 23–39.
- [15] SUAREZ, A. L. Revisiting the relation between subspaces and sublocales, 2020. Preprint available at arXiv:2010.05284.

IGOR ARRIETA

UNIVERSITY OF COIMBRA, CMUC, DEPARTMENT OF MATHEMATICS, PORTUGAL, AND UNIVERSIDAD DEL PAÍS VASCO UPV/EHU, DEPARTAMENTO DE MATEMÁTICAS, SPAIN

E-mail address: iarrieta024@ikasle.ehu.eus

ANNA LAURA SUAREZ

UNIVERSITY OF BIRMINGHAM, SCHOOL OF COMPUTER SCIENCE, UK, AND UNIVERSITY OF COIMBRA, CMUC, DEPARTMENT OF MATHEMATICS, PORTUGAL

E-mail address: annalaura.suarez@hotmail.it