

HOW PROTOMODULAR IS YOUR FAVOURITE CATEGORY?

MARIA MANUEL CLEMENTINO AND DIANA RODELO

ABSTRACT. In this work we investigate a relative notion of protomodularity. We say that a finitely complete category $\mathbf{C}$ satisfies the $\mathcal{M}$ -version of *protomodularity* when the change-of-base functors (concerning the fibration of points) are conservative with respect to a class of morphisms $\mathcal{M}$ of $\mathbf{C}$. If $\mathcal{M}$ is the class of all morphisms or the class of all monomorphisms of $\mathbf{C}$, then this notion is precisely that of an “ordinary” protomodular category. We analyse what conditions the class $\mathcal{M}$ should have to obtain the relative $\mathcal{M}$ -versions of some well-known properties of protomodular categories, such as the Split Short Five Lemma (when $\mathbf{C}$ is also pointed) and their relation with Mal’tsev categories. We also give diverse examples of categories satisfying the $\mathcal{M}$ -version of protomodularity for different choices of $\mathcal{M}$.

INTRODUCTION

Protomodular categories were introduced in [4] with the purpose to provide a good categorical setting to capture group-like features. The main example of a (pointed) protomodular category is **Grp**. Any other variety (of universal algebras) with more algebraic structure on its objects is still protomodular. On the other hand, the algebraically weaker variety **Mon** of monoids is not protomodular (for more examples of protomodular categories see [4, 5, 2]).

A finitely complete category $\mathbf{C}$ is called *protomodular* [4] when the change-of-base functors (with respect to the fibration of points $\Pi: \text{Pt}(\mathbf{C}) \rightarrow \mathbf{C}$) $h^*: \text{Pt}_Y(\mathbf{C}) \rightarrow \text{Pt}_Z(\mathbf{C})$ are conservative for any morphism $h: Z \rightarrow Y$. When $\mathbf{C}$ is also pointed, we can reduce our attention to morphisms of the type $h = 0_Y: 0 \rightarrow Y$ and the conservativeness of the kernel functors $0_Y^* = \ker: \text{Pt}_Y(\mathbf{C}) \rightarrow \mathbf{C}$ for any object Y . It easily follows that a pointed finitely complete category is protomodular if and only if the Split Short Five Lemma holds in $\mathbf{C}$ (see Theorem 1.3)

$$(0.i) \quad \begin{array}{ccccc} K' & \xrightarrow{\ker(f')} & X' & \xrightleftharpoons[s']{f'} & Y' \\ \downarrow \cong & & \downarrow m & & \downarrow \cong \\ K & \xrightarrow{\ker(f)} & X & \xrightleftharpoons[s]{f} & Y \end{array} \quad \Rightarrow \quad m \text{ is an isomorphism,}$$

which is well known to hold in the variety **Grp** of groups.

Since the change-of-base functors preserve finitely limits, any h^* is conservative if and only if it is conservative on monomorphisms. More precisely, if m is a monomorphism (in $\text{Pt}_Y(\mathbf{C})$) and $h^*(m)$ is an isomorphism (in $\text{Pt}_Z(\mathbf{C})$), then m is an isomorphism (in $\text{Pt}_Y(\mathbf{C})$). As a consequence, we only need to check the validity of the Split Short Five Lemma when m is a monomorphism.

2020 *Mathematics Subject Classification.* 18E08, 18E13, 18D20, 08C05, 18B25.

Key words and phrases. regular categories, categories enriched in the category of preorders, protomodular categories, Mal’tsev categories.

The authors acknowledge partial financial support by CMUC (doi.org/10.54499/UID/00324/2025), funded by the Portuguese Foundation for Science and Technology (FCT), grants UID/00324/2025 and UID/PRR/00324/2025. The second author also acknowledges financial support by CIDMA (https://ror.org/05pm2mw36), funded by the Portuguese Foundation for Science and Technology (FCT, https://ror.org/00snfq58), grants UID/04106/2025 (https://doi.org/10.54499/UID/04106/2025) and UID/PRR/04106/2025.

The category $\mathbf{OrdGrp}$ of (pre)ordered groups and continuous (=order-preserving) group homomorphisms is not protomodular, even though its objects are groups. It was shown in [11] that the Split Short Five Lemma fails in $\mathbf{OrdGrp}$. Still, $\mathbf{OrdGrp}$ satisfies a weak version of it: given the commutative diagram (0.i), then m is an isomorphism provided that it is a strong monomorphism. This led us to study categories satisfying a relative notion of protomodularity, placing a class $\mathcal{M}$ of morphisms where all morphisms were considered in the notion of protomodular category. This gave rise to the notion of category satisfying an $\mathcal{M}$ -version of protomodularity, which we simply call $\mathcal{M}$ -protomodular, proposed in this paper (see Definition 1.1). This should not be confused with the notion of $\mathcal{S}$ -protomodularity introduced by Bourn, Martins-Ferreira, Montoli and Sobral in [7], where $\mathcal{S}$ refers to a class of points (or split extensions).

The aim of this work is to investigate $\mathcal{M}$ -protomodularity for different classes $\mathcal{M}$ of (mono)morphisms in $\mathbf{C}$, with particular interest in finding examples of categories that are $\mathcal{M}$ -protomodular, where $\mathcal{M}$ is not the class of all (mono)morphisms (i.e. $\mathbf{C}$ is not protomodular in the usual sense). Although these categories are not “truly” protomodular, they still exhibit some sort of protomodular flavour. Of course, any finitely complete category $\mathbf{C}$ is always $\mathcal{M}$ -protomodular when $\mathcal{M}$ is the class of isomorphisms of $\mathbf{C}$. If this is the only class for which it is $\mathcal{M}$ -protomodular, then $\mathbf{C}$ is far from being protomodular. If $\mathbf{C}$ is $\mathcal{M}$ -protomodular, with $\mathcal{M}$ being the class of all (mono)morphisms, then $\mathbf{C}$ is truly protomodular. This led us to the title of this work.

In Section 1 we translate to our relative context well-known characterisations of protomodular category, showing that their $\mathcal{M}$ -versions are equivalent under very mild conditions on $\mathcal{M}$.

Sections 2 and 3 are dedicated to special properties and examples of $\mathcal{M}$ -protomodular categories. As expected, in most of our examples the class of strong monomorphisms plays a crucial role. Still, the more general setting described in Section 3, where $\mathcal{M}$ is the class $\mathbf{Spl}$ of split monomorphisms, turns out to be very interesting; in particular, it is shown that in $\mathbf{Spl}$ -protomodular categories pullbacks reflect monomorphisms.

In this context it is natural to introduce an $\mathcal{M}$ -version of Mal'tsev categories. That is the goal of Section 4, where it is shown that, as in the classical setting, $\mathcal{M}$ -protomodular categories are $\mathcal{M}$ -Mal'tsev categories.

In the last section we analyse $\mathcal{M}$ -protomodularity from the point of view of $\mathbf{Ord}$ -enriched categories. In the examples studied throughout this paper it turns out that the class of full and faithful (ff) monomorphisms is not strong enough to provide relative protomodularity for the $\mathbf{Ord}$ -enrichments studied. Hence it is an open question whether there is an interesting $\mathcal{M}$ -protomodular $\mathbf{Ord}$ -category for $\mathcal{M}$ the class of ff-monomorphisms.

1. $\mathcal{M}$ -PROTOMODULARITY

In this section we will consider $\mathbf{C}$ to be a finitely complete category, although the existence of pullbacks along split epimorphisms will suffice for most of the results. Given an object Y of $\mathbf{C}$, a *point over Y* is given by a pair of morphisms (f, s) , where $f: X \rightarrow Y$ is a split epimorphism with a chosen splitting $s: Y \rightarrow X$. We denote by $\mathbf{Pt}_Y(\mathbf{C})$ the category of points over Y in $\mathbf{C}$. Given a morphism $h: Z \rightarrow Y$, its corresponding *change-of-base functor* h^* is defined as

$$\begin{array}{ccc}
h^*: \text{Pt}_Y(\mathcal{C}) & \longrightarrow & \text{Pt}_Z(\mathcal{C}), \\
\begin{array}{ccc} X' & & \\ \swarrow f' & & \searrow s' \\ & Y & \\ \nwarrow m & & \nearrow s \\ X & & \end{array} & \longmapsto & \begin{array}{ccc} Z \times_Y X' & & \\ \swarrow \pi'_Z & & \searrow i'_Z \\ & Z & \\ \nwarrow h^*(m) & & \nearrow \pi_Z \\ Z \times_Y X & & \end{array}
\end{array}$$

where the front and back faces of the following diagram are pullbacks

$$\begin{array}{ccccc}
Z \times_Y X' & \xrightarrow{\pi_{X'}} & X' & & \\
\swarrow h^*(m) & & \searrow m & & \\
& & X & \xrightarrow{\pi_X} & Y \\
\swarrow \pi'_Z & & \nwarrow f' & & \nwarrow s' \\
& & Z & \xrightarrow{h} & Y \\
& & \uparrow i_Z & & \uparrow s \\
& & \pi_Z & & f
\end{array}$$

(1.i)

$i_Z = \langle 1_Z, sh \rangle$ and $i'_Z = \langle 1_Z, s'h \rangle$. Note that a morphism $m: (f', s') \rightarrow (f, s)$ is an isomorphism (resp. monomorphism) in $\text{Pt}_Y(\mathcal{C})$ if and only if $m: X' \rightarrow X$ is an isomorphism (resp. monomorphism) in $\mathcal{C}$.

We will be interested in studying the conservativeness of the change-of-base functors with respect to a certain class $\mathcal{M}$ of morphisms of $\mathcal{C}$. We say that h^* is *conservative on $\mathcal{M}$* when for any $m: (f', s') \rightarrow (f, s)$ in $\text{Pt}_Y(\mathcal{C})$ with $m \in \mathcal{M}$, if $h^*(m)$ is an isomorphism, then m itself is an isomorphism. If $\mathcal{M} = \text{Mono}$ is the class of all monomorphisms of $\mathcal{C}$, we say that h^* is conservative on monomorphisms. It is well-known that the conservativeness of a left exact functor is equivalent to its conservativeness on monomorphisms. This applies to the change-of-base functors. Indeed, it is easy to check that h^* preserves finite limits (this follows from the fact that $\mathcal{C}$ admits pullbacks of split epimorphisms).

In the proofs concerning the conservativeness of h^* , we eventually assume that the isomorphism $h^*(m)$ is an identity morphism.

Definition 1.1. Let $\mathcal{C}$ be a finitely complete category and $\mathcal{M}$ a class of morphisms of $\mathcal{C}$. We say that $\mathcal{C}$ satisfies the *$\mathcal{M}$ -version of protomodularity* when, for any morphism $h: Z \rightarrow Y$, the change-of-base functor $h^*: \text{Pt}_Y(\mathcal{C}) \rightarrow \text{Pt}_Z(\mathcal{C})$ is conservative on $\mathcal{M}$.

For simplicity, we will often say instead that $\mathcal{C}$ is *$\mathcal{M}$ -protomodular*. This should not be confused with the well-known notion of $\mathcal{S}$ -protomodular category introduced in [7], where $\mathcal{S}$ refers to a class of points of $\mathcal{C}$.

- Example 1.2.**
1. When $\mathcal{M}$ is the class Mor of all morphisms or the class Mono of all monomorphisms in $\mathcal{C}$, $\mathcal{C}$ satisfies the $\mathcal{M}$ -version of protomodularity exactly when $\mathcal{C}$ is protomodular [4].
 2. When $\mathcal{M}$ is the class $\mathcal{SMono}$ of all strong monomorphisms in $\mathcal{C}$, an $\mathcal{M}$ -protomodular category is known as a *weakly protomodular category* [11].
 3. For examples concerning other classes $\mathcal{M}$ see Sections 2 and 3.

We want to explore the several known characterisations of protomodular categories in the context of $\mathcal{M}$ -protomodular categories. Mainly, we are interested in the following ones (see [4, 2]):

Theorem 1.3. *Let $\mathbf{C}$ be a finitely complete category. The following conditions are equivalent:*

- (i) $\mathbf{C}$ is protomodular, i.e. the change-of-base functors $h^*: \text{Pt}_Y(\mathbf{C}) \rightarrow \text{Pt}_Z(\mathbf{C})$ are conservative for every $h: Z \rightarrow Y$;
- (ii) if $\mathbf{C}$ is also pointed, the Split Short Five Lemma holds: given a commutative diagram of split short exact sequences

$$(1.ii) \quad \begin{array}{ccccc} K' & \xrightarrow{\ker(f')} & X' & \xrightleftharpoons[s']{f'} & Y' \\ \downarrow \cong & & \downarrow m & & \downarrow \cong \\ K & \xrightarrow{\ker(f)} & X & \xrightleftharpoons[s]{f} & Y \end{array}$$

it follows that m is also an isomorphism;

- (iii) for any pullback of a point (f, s) along an arbitrary morphism

$$(1.iii) \quad \begin{array}{ccc} Z \times_Y X & \xrightarrow{\pi_X} & X \\ \pi_Z \downarrow \uparrow i_Z & \lrcorner & \downarrow f \uparrow s \\ Z & \xrightarrow{h} & Y \end{array}$$

the pair (π_X, s) is jointly extremally epimorphic;

- (iv) for any commutative diagram involving points (f, s) and (g, t)

$$(1.iv) \quad \begin{array}{ccccc} U & \xrightarrow{u} & X & \xrightarrow{x} & A \\ g \uparrow \downarrow t & \boxed{A} & f \uparrow \downarrow s & \boxed{B} & \downarrow q \\ V & \xrightarrow{v} & Y & \xrightarrow{y} & B \end{array}$$

(also commuting with the upward sections) where $\boxed{A}$ and $\boxed{A} \boxed{B}$ are pullbacks, then $\boxed{B}$ is also a pullback.

Proof. (i) $\Leftrightarrow$ (ii) $\Leftrightarrow$ (iii): see [4, 2].

(i) $\Leftrightarrow$ (iv): see [4, 6, 2]. □

When $\mathbf{C}$ is a pointed category, we say that the Split Sort Five Lemma holds *with respect to* $\mathcal{M}$ when the property stated in Theorem 1.3(ii) holds for $m \in \mathcal{M}$. Without loss of generality, and to simplify proofs, we shall consider the isomorphisms of the diagram of the Split Short Five Lemma to be the identity morphisms. As a consequence, in diagram (1.i), we take $Z \times_Y X' = Z \times_Y X$, $\pi'_Z = \pi_Z$ and $i'_Z = i_Z$.

The proofs of the $\mathcal{M}$ -versions in the following results are similar to those for “ordinary” protomodular categories.

Proposition 1.4. *Let $\mathbf{C}$ be a pointed finitely complete category and $\mathcal{M}$ any class of morphisms of $\mathbf{C}$. The following conditions are equivalent:*

- (i) $\mathbf{C}$ is $\mathcal{M}$ -protomodular;
- (ii) the Split Short Five Lemma holds with respect to $\mathcal{M}$.

Proof. For $\mathcal{M} = \text{Mor}$ or $\mathcal{M} = \text{Mono}$ this result is the equivalence between (i) and (ii) of Theorem 1.3.

(i) $\Rightarrow$ (ii) We consider diagram (1.ii) with $K' = K$ and $Y' = Y$, so that the isomorphisms are the identity morphisms; suppose that $m \in \mathcal{M}$. This diagram shows that $0_Y^*(m) = 1_K$ which, by assumption, implies that m is an isomorphism.

(ii) $\Rightarrow$ (i) Consider an arbitrary morphism $h: Z \rightarrow Y$ of $\mathcal{C}$ and a morphism $m: X' \rightarrow X$ in $\mathcal{M}$ such that $h^*(m) = 1_P$

(1.v)

$$\begin{array}{ccccc}
 P & \xrightarrow{\pi_{X'}} & X' & & \\
 \swarrow \scriptstyle h^*(m)=1_P & & \searrow \scriptstyle m \in \mathcal{M} & & \\
 P & \xrightarrow{\pi_X} & X & & \\
 \swarrow \scriptstyle \pi_Z & \nearrow \scriptstyle i_Z & \searrow \scriptstyle f' & \nearrow \scriptstyle s' & \\
 & P & & X & \\
 & \uparrow \scriptstyle \pi_Z & & \uparrow \scriptstyle f & \\
 & Z & \xrightarrow{h} & Y & \\
 & & & \uparrow \scriptstyle s &
 \end{array}$$

here $P = Z \times_Y X = Z \times_Y X'$. By taking the kernel of π_Z we get $\pi_{X'} \ker(\pi_Z) = \ker(f')$ and $\pi_X \ker(\pi_Z) = \ker(f)$

$$\begin{array}{ccccc}
 K & \xrightarrow{\ker(f')} & X' & \xrightleftharpoons[s']{f'} & Y \\
 \parallel & & \downarrow \scriptstyle m \in \mathcal{M} & & \parallel \\
 K & \xrightarrow{\ker(f)} & X & \xrightleftharpoons[s]{f} & Y,
 \end{array}$$

so that we can conclude that m is an isomorphism, by assumption. $\square$

Proposition 1.5. *Let $\mathcal{C}$ be a finitely complete category and $\mathcal{M}$ a class of monomorphisms of $\mathcal{C}$. The following conditions are equivalent:*

- (i) $\mathcal{C}$ is $\mathcal{M}$ -protomodular;
- (ii) for any pullback of a point (f, s) along an arbitrary morphism as in (1.iii), the pair (π_X, s) is orthogonal to the class $\mathcal{M}$.

Proof. For $\mathcal{M} = \mathbf{Mono}$ this result is the equivalence between (i) and (iii) of Theorem 1.3.

(i) $\Rightarrow$ (ii) To prove that (π_X, s) is orthogonal to $\mathcal{M}$, we consider the commutative diagram

$$\begin{array}{ccccc}
 & & M & & \\
 & \nearrow \scriptstyle a & \downarrow \scriptstyle m \in \mathcal{M} & \nwarrow \scriptstyle b & \\
 Z \times_Y X & \xrightarrow{\pi_X} & X & \xleftarrow{s} & Y.
 \end{array}$$

This gives the commutative diagram

$$\begin{array}{ccccc}
 Z \times_Y X & \xrightarrow{a} & M & & \\
 \swarrow \scriptstyle h^*(m)=1_{Z \times_Y X} & & \searrow \scriptstyle m \in \mathcal{M} & & \\
 Z \times_Y X & \xrightarrow{\pi_X} & X & & \\
 \swarrow \scriptstyle \pi_Z & \nearrow \scriptstyle i_Z & \searrow \scriptstyle fm & \nearrow \scriptstyle b & \\
 & Z \times_Y X & & X & \\
 & \uparrow \scriptstyle \pi_Z & & \uparrow \scriptstyle f & \\
 & Z & \xrightarrow{h} & Y & \\
 & & & \uparrow \scriptstyle s &
 \end{array}$$

where the front and top faces are pullbacks (the top face is a pullback since m is a monomorphism). Note that $mai_Z = \pi_X i_Z = sh = mbh$ implies that $ai_Z = bh$, since m is a monomorphism. This proves that the back face commutes with respect to the sections. By assumption we may conclude that m is an isomorphism.

(ii) $\Rightarrow$ (i) Consider diagram (1.v) with m a monomorphism in $\mathcal{M}$. We get

$$\begin{array}{ccccc} & & X' & & \\ & \nearrow \pi'_X & \downarrow m \in \mathcal{M} & \nwarrow s' & \\ P & \xrightarrow{\pi_X} & X & \xleftarrow{s} & Y; \end{array}$$

by assumption, m is an isomorphism. $\square$

Proposition 1.6. *Let $\mathcal{C}$ be a finitely complete category and $\mathcal{M}$ a class of morphisms of $\mathcal{C}$ which is left-cancellative, that is, $ba \in \mathcal{M} \Rightarrow a \in \mathcal{M}$. The following conditions are equivalent:*

- (i) $\mathcal{C}$ is $\mathcal{M}$ -protomodular;
- (ii) for any commutative diagram involving points (f, s) and (g, t)

$$\begin{array}{ccccc} U & \xrightarrow{u} & X & \xrightarrow{m \in \mathcal{M}} & A \\ \uparrow g & & \uparrow f & & \downarrow q \\ \downarrow t & \boxed{A} & \downarrow s & \boxed{B} & \\ V & \xrightarrow{v} & Y & \xrightarrow{y} & B \end{array}$$

(also commuting with the upward sections) where $\boxed{A}$ and $\boxed{A \mid B}$ are pullbacks, then $\boxed{B}$ is also a pullback.

Proof. For $\mathcal{M} = \text{Mor}$ or $\mathcal{M} = \text{Mono}$ this result is the equivalence between (i) and (iv) of Theorem 1.3.

(i) $\Rightarrow$ (ii) Take the pullback of (y, q) and the induced morphism z in

$$\begin{array}{ccccc} X & \xrightarrow{m \in \mathcal{M}} & A & & \\ & \searrow z & \downarrow \pi_A & & \\ & & Z & \xrightarrow{\pi_A} & A \\ & \nearrow f & \downarrow \pi_Y & \nearrow zs & \downarrow q \\ & & Y & \xrightarrow{y} & B. \end{array}$$

Note that $z \in \mathcal{M}$, because $\mathcal{M}$ is left-cancellative. We get

$$\begin{array}{ccccc} U & \xrightarrow{zu} & Z & \xrightarrow{\pi_A} & A \\ \uparrow g & & \uparrow \pi_Y & & \downarrow q \\ \downarrow t & \boxed{D} & \downarrow zs & \boxed{C} & \\ V & \xrightarrow{v} & Y & \xrightarrow{y} & B, \end{array}$$

where $\boxed{D \mid C} = \boxed{A \mid B}$ and $\boxed{C}$ are pullbacks, so that $\boxed{D}$ is a pullback. This gives the following commutative diagram

$$\begin{array}{ccccc} U & \xrightarrow{u} & X & & \\ \searrow g & & \searrow f & & \\ & \nearrow v^*(z) = 1_U & \nearrow z \in \mathcal{M} & & \\ & & Z & \xrightarrow{\pi_Y} & Y \\ & \nearrow t & \downarrow \pi_Y & \nearrow zs & \\ & & Y & \xrightarrow{y} & Y, \end{array}$$

from which we deduce that z is an isomorphism; consequently $\boxed{B}$ is a pullback.

(ii) $\Rightarrow$ (i) Consider diagram (1.v). We get

$$\begin{array}{ccccc}
 & & \pi_X & & \\
 & \curvearrowright & & \curvearrowright & \\
 P & \xrightarrow{\pi_{X'}} & X' & \xrightarrow{m \in \mathcal{M}} & X \\
 \uparrow \pi_Z & \downarrow i_Z & \downarrow f' & \downarrow s' & \downarrow f \\
 & \boxed{A} & & \boxed{B} & \\
 Z & \xrightarrow{h} & Y & \xrightarrow{1_Y} & Y \\
 & \curvearrowleft & & \curvearrowleft & \\
 & & h & &
 \end{array}$$

where $\boxed{A}$ and $\boxed{A \mid B}$ are pullbacks, so that $\boxed{B}$ is also a pullback; thus m is an isomorphism. $\square$

Proposition 1.7. *Let $\mathcal{C}$ be a regular category and $\mathcal{M}$ a class of morphisms of $\mathcal{C}$ with the properties*

- (P1) $\mathcal{M}$ is left-cancellative;
- (P2) $\mathcal{M}$ is closed under composition;
- (P3) $m \in \mathcal{M} \Rightarrow m \times m \in \mathcal{M}$;
- (P4) for any pullback projections π_Z, π_X , the monomorphism $\langle \pi_Z, \pi_X \rangle: Z \times_Y X \rightarrow Z \times X$ is in $\mathcal{M}$.

The following conditions are equivalent:

- (i) $\mathcal{C}$ is $\mathcal{M}$ -protomodular;
- (ii) for any commutative diagram

$$\begin{array}{ccccc}
 U & \xrightarrow{u} & X & \xrightarrow{m \in \mathcal{M}} & A \\
 \downarrow g & & \downarrow f & & \downarrow q \\
 & \boxed{A} & & \boxed{B} & \\
 V & \xrightarrow{v} & Y & \xrightarrow{y} & B
 \end{array}$$

where f, g are regular epimorphisms, if $\boxed{A}$ and $\boxed{A \mid B}$ are pullbacks, then $\boxed{B}$ is also a pullback.

Proof. For $\mathcal{M} = \text{Mor}$ or $\mathcal{M} = \text{Mono}$ the proof of the equivalence between (i) and (ii) can be found in [4].

(i) $\Rightarrow$ (ii) Take vertical kernel pairs as in

$$\begin{array}{ccccc}
 \text{Eq}(g) & \xrightarrow{\bar{u}} & \text{Eq}(f) & \xrightarrow{\bar{m}} & \text{Eq}(q) \\
 \downarrow g_1 & \downarrow g_2 & \downarrow f_1 & \downarrow f_2 & \downarrow q_1 \\
 & \boxed{C} & & \boxed{D} & \\
 U & \xrightarrow{u} & X & \xrightarrow{m \in \mathcal{M}} & A \\
 \downarrow g & & \downarrow f & & \downarrow q \\
 & \boxed{A} & & \boxed{B} & \\
 V & \xrightarrow{v} & Y & \xrightarrow{y} & B
 \end{array}$$

Since $\boxed{A}$ and $\boxed{A \mid B}$ are pullbacks, then so are $\boxed{C}$ and $\boxed{C \mid D}$. From properties (P2), (P3) and (P4), $(m \times m)\langle f_1, f_2 \rangle = \langle q_1, q_2 \rangle \bar{m} \in \mathcal{M}$; thus $\bar{m} \in \mathcal{M}$ by property (P1). We can apply Proposition 1.6 to $\boxed{C \mid D}$, to conclude that $\boxed{D}$ is a pullback. By the Barr-Kock Theorem [1] we conclude that $\boxed{B}$ is a pullback as well.

(ii) $\Rightarrow$ (i) Follows immediately from Proposition 1.6. $\square$

Proposition 1.8. *Let $\mathcal{C}$ be a pointed regular category and $\mathcal{M}$ a class of morphisms of $\mathcal{C}$ with the properties*

- (P1) $\mathcal{M}$ is left-cancellative;
- (P2) $\mathcal{M}$ is closed under composition;
- (P3) $m \in \mathcal{M} \Rightarrow m \times m \in \mathcal{M}$;
- (P4) for any pullback projections π_Z, π_X , the monomorphism $\langle \pi_Z, \pi_X \rangle: Z \times_Y X \rightarrow Z \times X$ is in $\mathcal{M}$.

The following conditions are equivalent:

- (i) $\mathcal{C}$ is $\mathcal{M}$ -protomodular;
- (ii) the Short Five Lemma holds with respect to $\mathcal{M}$, i.e. given a commutative diagram of short exact sequences,

$$\begin{array}{ccccc} K & \xrightarrow{\ker(f')} & X' & \xrightarrow{f'} & Y \\ \parallel & & \downarrow m \in \mathcal{M} & & \parallel \\ K & \xrightarrow{\ker(f)} & X & \xrightarrow{f} & Y, \end{array}$$

m is necessarily an isomorphism.

Proof. For $\mathcal{M} = \mathcal{M}\text{or}$ or $\mathcal{M} = \mathcal{M}\text{ono}$ this result is the equivalence between (i) and (iv) can be found in [4, 6, 2].

(i) $\Rightarrow$ (ii) We get the following commutative diagram

$$\begin{array}{ccccc} & & \text{ker}(f) & & \\ & \text{ker}(f') & \searrow & & \\ K & \xrightarrow{\quad} & X' & \xrightarrow{m \in \mathcal{M}} & X \\ \downarrow & \boxed{A} & \downarrow f' & \boxed{B} & \downarrow f \\ 0 & \xrightarrow{\quad} & Y & \xlongequal{\quad} & Y, \end{array}$$

where $\boxed{A}$ and $\boxed{A} \boxed{B}$ are pullbacks. By Proposition 1.7 we conclude that $\boxed{B}$ is a pullback, thus m is an isomorphism.

(ii) $\Rightarrow$ (i) Follows from Proposition 1.4. $\square$

Remark 1.9. In the following sections we are mainly interested in studying $\mathcal{M}$ -protomodularity for the classes of monomorphisms $\mathcal{M} = \mathcal{S}\mathcal{M}\text{ono}$, $\mathcal{M} = \mathcal{S}\mathcal{P}\mathcal{l}$ the class of split monomorphisms and $\mathcal{M} = \mathcal{P}\mathcal{S}\mathcal{P}\mathcal{l}$ the class of protosplit monomorphisms (=kernels of split epimorphisms). $\mathcal{S}\mathcal{M}\text{ono}$ has the properties (P1)-(P4), so any of the above equivalences hold for $\mathcal{S}\mathcal{M}\text{ono}$ -protomodular categories, as long as $\mathcal{C}$ satisfies their initial assumptions. $\mathcal{S}\mathcal{P}\mathcal{l}$ has properties (P1)-(P3) but not (P4), so only the equivalences in Propositions 1.4, 1.5 and 1.6 hold for $\mathcal{S}\mathcal{P}\mathcal{l}$ -protomodular (pointed) categories. $\mathcal{P}\mathcal{S}\mathcal{P}\mathcal{l}$ only satisfies (P2); thus only the equivalences in Propositions 1.4 and 1.5 hold for $\mathcal{P}\mathcal{S}\mathcal{P}\mathcal{l}$ -protomodular pointed categories.

2. EXAMPLES OF $\mathcal{M}$ -PROTOMODULAR CATEGORIES

We shall use the following result to obtain several interesting examples of $\mathcal{M}$ -protomodular categories.

Theorem 2.1. *Consider finitely complete categories $\mathcal{C}$ and $\mathcal{D}$, with classes of morphisms $\mathcal{M}$ and $\mathcal{N}$, respectively. Let $F: \mathcal{C} \rightarrow \mathcal{D}$ be a functor which preserves pullbacks, is conservative on $\mathcal{M}$ and such that $F(\mathcal{M}) \subseteq \mathcal{N}$. If $\mathcal{D}$ is an $\mathcal{N}$ -protomodular category, then $\mathcal{C}$ is an $\mathcal{M}$ -protomodular category.*

Proof. The proof is a straightforward adaptation of the well-known result concerning “ordinary” protomodularity [4]. $\square$

Corollary 2.2. *Let $F: \mathcal{C} \rightarrow \mathcal{D}$ be a topological functor between finitely complete categories, and $\mathcal{M}$ a class of F -initial morphisms of $\mathcal{C}$. Then $\mathcal{C}$ is $\mathcal{M}$ -protomodular provided that $\mathcal{D}$ is $F(\mathcal{M})$ -protomodular.*

In particular, if $\mathcal{D}$ is protomodular, then $\mathcal{C}$ is $\mathcal{SMono}$ -protomodular.

Proof. A topological functor preserves limits and colimits, it is conservative on initial morphisms (see [22]), and every strong monomorphism is initial. Hence the statements of our corollary follow from the theorem above. $\square$

Examples 2.3. It is well-known that:

- $\mathbf{Set}^{\text{op}}$ is a (co)complete protomodular category [4, 2];
- the forgetful functors $\mathbf{Top} \rightarrow \mathbf{Set}$, $\mathbf{Ord} \rightarrow \mathbf{Set}$, and more generally $V\text{-Cat} \rightarrow \mathbf{Set}$ (see [16]) are topological, therefore also $\mathbf{Top}^{\text{op}} \rightarrow \mathbf{Set}^{\text{op}}$, $\mathbf{Ord}^{\text{op}} \rightarrow \mathbf{Set}^{\text{op}}$, $(V\text{-Cat})^{\text{op}} \rightarrow \mathbf{Set}^{\text{op}}$ are topological since the notion of topological functor is self-dual. (Here $\mathbf{Top}$ denotes the category of topological spaces and continuous maps, $\mathbf{Ord}$ denotes the category of preordered sets and monotone maps, and $V\text{-Cat}$ denotes the category of V -categories and V -functors, for a commutative and unital quantale V .)

Hence Corollary 2.2 guarantees that $\mathbf{Ord}^{\text{op}}$, $\mathbf{Top}^{\text{op}}$, $(V\text{-Cat})^{\text{op}}$ are $\mathcal{SMono}$ -protomodular.

None of the categories $\mathbf{Top}^{\text{op}}$, $\mathbf{Ord}^{\text{op}}$, and consequently $(V\text{-Cat})^{\text{op}}$ in general, are protomodular. To prove this claim we give an example of a non-strong monomorphism m with $h^*(m)$ isomorphism. We describe our diagram in $\mathbf{Ord}$, for simplicity. In the following diagram

$$\begin{array}{ccc}
 A & \xrightarrow{h} & C \\
 \alpha \updownarrow & & \updownarrow \alpha' \\
 B & \xrightarrow{g} & D \\
 m \downarrow & & \downarrow \cong \\
 B' & \longrightarrow & D'
 \end{array}$$

$A = \{x_1, x_2, y_1, y_2\}$ with $x_1 \geq x_2 \leq y_2 \geq y_1$, $C = \{1, x_2, y_2\}$ with $1 \geq x_2 \leq y_2 \geq 1$, B and B' have as underlying set $\{x_1, x_2, y_1, y_2, x_0, y_0\}$, the order on B is given by the order on A together with $x_0 \leq x_1$, $y_1 \leq y_0$, and the order on B' has the extra relation $x_0 \leq y_0$. The morphism $h: A \rightarrow C$ is defined by $h(x_1) = h(y_1) = 1$, $h(x_2) = x_2$, $h(y_2) = y_2$. The inclusions $\alpha: A \rightarrow B$, $\alpha': C \rightarrow D$ are split monomorphisms, with retractions $\beta: B \rightarrow A$, $\beta': D \rightarrow C$ where both identify x_0 with x_1 and y_0 with y_1 . Now it is easy to check that the bijection $m: B \rightarrow B'$ is not an isomorphism, although its pushout along g is an isomorphism, since the identification of x_1 and y_1 by h implies that $x_0 \leq y_0$ in D .

Examples 2.4. Analogously:

- $\mathbf{Grp}$ is a (co)complete protomodular category;
- as shown in [11, 12, 13] respectively, the forgetful functors $\mathbf{OrdGrp} \rightarrow \mathbf{Grp}$, $\mathbf{ROrdGrp} \rightarrow \mathbf{Grp}$, $V\text{-Grp} \rightarrow \mathbf{Grp}$ are topological, where $\mathbf{ROrdGrp}$ denotes the category of right-ordered groups and monotone group homomorphisms, and $V\text{-Grp}$ has as objects V -groups and as morphisms V -functors which are group homomorphisms, for a suitable quantale V (as introduced in [12]).

Hence, using again Corollary 2.2, we conclude that the following categories are $\mathcal{SMono}$ -protomodular:

$$\mathbf{OrdGrp}, \mathbf{ROrdGrp}, \mathbf{V-Grp}.$$

None of these categories is protomodular, as shown in [11, 12, 13] respectively.

Example 2.5. It was shown in [3] that the category $\mathbf{TopGrp}$ of topological groups and continuous group homomorphisms is protomodular. If in the definition of topological group we drop the continuity of the inversion map we end up with the notion of *paratopological group*. Let $\mathbf{PTopGrp}$ be the category of paratopological groups and continuous group homomorphisms. It is a complete and cocomplete category, and the forgetful functor $\mathbf{PTopGrp} \rightarrow \mathbf{Grp}$ is topological; indeed the proof of topologicity of this functor uses the same arguments used to show that the forgetful functor $\mathbf{TopGrp} \rightarrow \mathbf{Grp}$ is topological. Hence, by Corollary 2.2, $\mathbf{PTopGrp}$ is $\mathcal{SMono}$ -protomodular.

However, in contrast to the case of $\mathbf{TopGrp}$, $\mathbf{PTopGrp}$ is not protomodular, showing that the inversion map of each group is essential for protomodularity. Let us show that the Split Short Five Lemma fails.

Let Y be the group $(\mathbb{R}, +)$ equipped with the topology $\mathcal{T}_S$ of the Sorgenfrey line; that is, the topology having $\{[0, a[; a \in \mathbb{R}, a > 0\}$ as basic neighbourhoods of 0. It is a paratopological group but not a topological group.

In $(\mathbb{R}, +) \times (\mathbb{R}, +)$ we may consider two different topologies: the product topology $\mathcal{T}_1 = \mathcal{T}_S \times \mathcal{T}_S$, and $\mathcal{T}_2$ having as basic neighbourhoods of 0

$$\{U_a = \mathbb{R} \times]0, a[\cup [0, a[\times \{0\}; a \in \mathbb{R}, a > 0\}.$$

It is straightforward to check that both make $\mathbb{R} \times \mathbb{R}$ a non-topological paratopological group.

Both $\mathcal{T}_1$ and $\mathcal{T}_2$ make the following split short exact sequence in $\mathbf{Grp}$

$$Y \xrightarrow{\langle 1, 0 \rangle} (\mathbb{R} \times \mathbb{R}, \mathcal{T}_i) \xrightleftharpoons[\langle 0, 1 \rangle]{\pi_2} Y$$

a split short exact sequence in $\mathbf{PTopGrp}$; indeed, as the product topology, the case of $\mathcal{T}_1$ clearly works. Let us check our claim for $\mathcal{T}_2$:

- $k = \langle 1, 0 \rangle$ is a strong monomorphism: $k^{-1}(U_a) = [0, a[$.
- $s = \langle 0, 1 \rangle$ is continuous: $s^{-1}(U_a) = [0, a[$.
- To show that π_2 is continuous it is enough to notice that π_2 is continuous in 0, and this follows from $\pi_2(U_a) = [0, a[$.

Now, for each $a > 0$, $U_a \supseteq [0, a[\times [0, a[$, hence U_a is clearly a neighbourhood of 0 in $\mathcal{T}_1$, which implies that the identity map $(\mathbb{R} \times \mathbb{R}, \mathcal{T}_1) \rightarrow (\mathbb{R} \times \mathbb{R}, \mathcal{T}_2)$ is continuous. Since it is not an isomorphism, this example shows that the Split Short Five Lemma fails.

Example 2.6. In $\mathbf{Mon}$, consider the class of morphisms $\mathcal{M}$ to be all isomorphisms in $\mathbf{Mon}$ and all group homomorphisms. Then $\mathbf{Mon}$ is $\mathcal{M}$ -protomodular. Indeed, consider diagram (1.i) with m a group homomorphism and $h^*(m)$ an isomorphism. Since f is surjective, then Y is also a group, so that the right face of this diagram lives in $\mathbf{Grp}$. The proof that m is an isomorphism follows as in $\mathbf{Grp}$, since the only inverses we must take are in X or X' .

3. Spl-PROTOMODULAR CATEGORIES

Consider the class $\mathbf{Spl}$ of split monomorphisms of $\mathbf{C}$. Every $\mathcal{SMono}$ -protomodular category is in particular a $\mathbf{Spl}$ -protomodular category. Next we will present an example of an $\mathbf{Spl}$ -protomodular category which is not $\mathcal{SMono}$ -protomodular. Before this example we note that:

Proposition 3.1. *If $\mathcal{C}$ is Spl-protomodular, then every change-of-base functor h^* reflects monomorphisms.*

Proof. Consider diagram (1.i) such that $h^*(m)$ is a monomorphism. We extend the diagram by taking kernel pairs of $h^*(m)$ and of m . Since the top part of diagram (1.i) is a pullback, so are the following downward commutative squares

$$\begin{array}{ccc} Z \times_Y X' & \xrightarrow{\quad} & \text{Eq}(m) \\ \parallel & \lrcorner & \downarrow m_1 \uparrow s_0 \downarrow m_2 \\ Z \times_Y X' & \xrightarrow{\pi_{X'}} & X', \end{array}$$

where $s_0 = \langle 1_{X'}, 1_{X'} \rangle$. Consequently, the upward square is also a pullback. We get

$$\begin{array}{ccccc} Z \times_Y X' & \xrightarrow{\pi_{X'}} & X' & & \\ \parallel & \searrow h^*(s_0) = 1_{Z \times_Y X'} & \searrow s_0 & & \\ & & & \searrow f' & \\ & & & & \text{Eq}(m) \\ & \searrow \pi'_Z & \searrow i'_Z & \searrow f'm_1 & \parallel s_0 s' \\ & & Z & \xrightarrow{h} & Y. \end{array}$$

Since s_0 is a split monomorphism such that $h^*(s_0)$ is an isomorphism, then s_0 is an isomorphism, by assumption. It easily follows that each m_i is a isomorphism; thus m is a monomorphism. $\square$

Lemma 3.2. *In a finitely complete category $\mathcal{C}$, the following conditions are equivalent:*

- (i) *any h^* reflects monomorphisms;*
- (ii) *pullbacks in $\mathcal{C}$ reflect monomorphisms.*

Proof. (ii) $\Rightarrow$ (i) Given diagram (1.i) with $h^*(m)$ a monomorphism, we conclude that m is a monomorphism since the top part of this diagram is a pullback in $\mathcal{C}$.

(i) $\Rightarrow$ (ii) Consider the following diagram where the bottom square is a pullback in $\mathcal{C}$, m is a monomorphism and the top is obtain by taking vertical kernel pairs

$$\begin{array}{ccc} A & \longrightarrow & \text{Eq}(n) \\ \parallel & & \downarrow n_1 \uparrow s_0 \downarrow n_2 \\ A & \longrightarrow & C \\ \downarrow m & \lrcorner & \downarrow n \\ B & \longrightarrow & D; \end{array}$$

here $s_0 = \langle 1_C, 1_C \rangle$. It follows that the downward top squares are pullbacks. This implies that $n_1: (n_1, s_0) \rightarrow (1_C, 1_C)$ is a morphism in $\text{Pt}_{\mathcal{C}}(\mathcal{C})$ such that $f^*(n_1) = 1_A$. By assumption, n_1 is a monomorphism, which implies it is an isomorphism in $\mathcal{C}$ (since it is also a split epimorphism). It easily follows that n is a monomorphism. $\square$

Corollary 3.3. *Let $\mathcal{C}$ be a pointed Spl-protomodular category. If $\ker(n: C \rightarrow D) = 0$, then n is a monomorphism.*

Proof. This easily follows by combining Proposition 3.1 and Lemma 3.2. $\square$

Remark 3.4. It is well-known that in every protomodular category pullbacks reflect monomorphisms [4]. We do not know whether the equivalent properties of Lemma 3.2 characterise Spl-protomodular categories.

Example 3.5. Mon is not a Spl-protomodular category since there exist monoid homomorphisms with trivial kernel that are not monomorphisms. So, we find pullbacks that do not reflect monomorphisms (see Corollary 3.3).

A category is called *semi-abelian* [19] when it is pointed, finitely (co)complete, Barr-exact [1] and protomodular. Let $\mathcal{C}$ be a semi-abelian category and $\mathcal{M}$ a class of monomorphisms in $\mathcal{C}^{\text{op}}$ such that all morphisms in $\mathcal{M}^{\text{op}}$ are cokernels. By Proposition 1.5, $\mathcal{C}^{\text{op}}$ is $\mathcal{M}$ -protomodular if and only if for any pushout diagram in $\mathcal{C}$

$$(3.i) \quad \begin{array}{ccc} A & \xrightarrow{h} & C \\ \alpha \uparrow \beta & & \uparrow \alpha' \beta' \\ B & \xrightarrow{u} & D, \end{array}$$

where α is a split monomorphism with retraction β , the pair of morphisms (u, β) is orthogonal to the class $\mathcal{M}^{\text{op}}$. This means that: if (u, β) factors through a morphism $e \in \mathcal{M}^{\text{op}}$ as in the commutative diagram

$$\begin{array}{ccccc} & & D & \xleftarrow{u} & B & \xrightarrow{\beta} & A, \\ & & & & \downarrow e \in \mathcal{M}^{\text{op}} & & \\ & & & & E & & \end{array}$$

then e is an isomorphism. Since e is a cokernel, it must be the cokernel of its kernel; thus the vertical arrows in the following diagram give a short exact sequence. By taking the kernels of β, e and u , there exists a factorisation as above if and only if there exist the induced morphisms $\bar{a}, \bar{b}$

$$(3.ii) \quad \begin{array}{ccccc} & & \text{Ker}(e) & & \\ & \swarrow \bar{b} & \downarrow \text{ker}(e) & \searrow \bar{a} & \\ & \text{Ker}(\beta) & & & \text{Ker}(u) \\ & \swarrow \text{ker}(\beta) & \downarrow \text{ker}(u) & \searrow \beta & \\ D & \xleftarrow{u} & B & \xrightarrow{\beta} & A. \\ & & \downarrow e \in \mathcal{M}^{\text{op}} & & \\ & & E & & \end{array}$$

Lemma 3.6. Let $\mathcal{C}$ be a semi-abelian category and $\mathcal{M}$ a class of monomorphisms in $\mathcal{C}^{\text{op}}$ such that all morphisms in $\mathcal{M}^{\text{op}}$ are cokernels. Then $\mathcal{C}^{\text{op}}$ is $\mathcal{M}$ -protomodular if and only if, for any pushout of type (3.i) and cokernel $e \in \mathcal{M}^{\text{op}}$, we have

$$(3.iii) \quad (\text{Ker}(e) \subseteq \text{Ker}(\beta) \cap \text{Ker}(u)) \Rightarrow \text{Ker}(e) = 0.$$

Proof. Immediate from diagram (3.ii) and the fact that $\text{Ker}(e) = 0$ implies that (the cokernel) e is a monomorphism, since $\mathcal{C}$ is a protomodular category. Thus, e is an isomorphism. $\square$

Given a group G , we consider a descending chain of normal subgroups of G

$$G = \gamma_1(G) \supseteq \gamma_2(G) \supseteq \cdots$$

where $\gamma_1(G) = G$, $\gamma_2(G) = [G, G] = \{xyx^{-1}y^{-1} : x, y \in G\}$, $\gamma_3(G) = [\gamma_2(G), G]$, $\dots$. Given a natural number $c \geq 1$, we say that G is a *nilpotent group of class c* if $\gamma_{c+1}(G) = 0$. A nilpotent group of class 1 is just an abelian group. We fix a natural number $c \geq 2$ and denote by Nil_c the variety of nilpotent groups of class (at most) c . It is easy to check that Nil_c is a semi-abelian variety such that $\text{Ab} \subseteq \text{Nil}_c \subseteq \text{Grp}$. As a consequence, the abelianisation functor for Nil_c is defined the same way as for Grp

$$\begin{aligned} (-)^{\text{ab}} : \text{Nil}_c &\longrightarrow \text{Ab} \\ G &\longmapsto G^{\text{ab}} = G/[G, G] \end{aligned}$$

The reflection morphism with respect to abelianisation is given by a surjective homomorphism denoted by $\eta_G : G \rightarrow G^{\text{ab}}$, for any G in Nil_c .

Lemma 3.7. *Under the conditions of Lemma 3.6 applied to $\mathbf{C} = \text{Grp}$ or $\mathbf{C} = \text{Nil}_c$, we have $\text{Ker}(\beta) \cap \text{Ker}(u) \subseteq [B, B]$.*

Proof. We give the proof for $\mathbf{C} = \text{Grp}$; the proof for $\mathbf{C} = \text{Nil}_c$ is similar. The abelianisation functor preserves the pushout (3.i) (since it is a left adjoint), so that

$$\begin{array}{ccc} A^{\text{ab}} & \xrightarrow{h^{\text{ab}}} & C^{\text{ab}} \\ \alpha^{\text{ab}} \downarrow & \beta^{\text{ab}} \uparrow & \downarrow \alpha'^{\text{ab}} \\ B^{\text{ab}} & \xrightarrow{u^{\text{ab}}} & D^{\text{ab}}, \end{array}$$

is a pushout in Ab . It is easy to check that the kernel of β^{ab} is isomorphic to the reflection η_B restricted to $\text{Ker}(\beta)$; call it K . This leads to the isomorphic split short exact sequence in Ab

$$\begin{array}{ccccc} \text{Ker}(\beta^{\text{ab}}) & \xrightarrow{\text{ker}(\beta^{\text{ab}})} & B^{\text{ab}} & \xrightleftharpoons[\alpha^{\text{ab}}]{\beta^{\text{ab}}} & A^{\text{ab}} \\ \parallel & & \downarrow \cong & & \parallel \\ \text{Ker}(\beta^{\text{ab}}) & \xrightarrow{\langle 1, 0 \rangle} & \text{Ker}(\beta^{\text{ab}}) \oplus A^{\text{ab}} & \xrightleftharpoons[\langle 0, 1 \rangle]{\pi_2} & A^{\text{ab}} \\ \downarrow \cong & & \downarrow \cong & & \parallel \\ K & \xrightarrow{\langle 1, 0 \rangle} & K \oplus A^{\text{ab}} & \xrightleftharpoons[\langle 0, 1 \rangle]{\pi_2} & A^{\text{ab}}. \end{array}$$

Then, the downward pushout from above may be simplified to

$$\begin{array}{ccc} A^{\text{ab}} & \xrightarrow{h^{\text{ab}}} & C^{\text{ab}} \\ \langle 0, 1 \rangle \downarrow & & \downarrow \langle 0, 1 \rangle \\ K \oplus A^{\text{ab}} & \xrightarrow{1_K \oplus h^{\text{ab}}} & K \oplus C^{\text{ab}}. \end{array}$$

Now we prove that $\text{Ker}(\beta) \cap \text{Ker}(u) \subseteq [B, B]$. Given $x \in \text{Ker}(\beta) \cap \text{Ker}(u)$, we have $\eta_B(x) \in K$. On the other hand, $u^{\text{ab}}(\eta_B(x)) = \eta_D(u(a)) = 0$. Using the fact that $u^{\text{ab}} \cong 1_K \oplus h^{\text{ab}}$, this last equality is equivalent to $1_K \oplus h^{\text{ab}}(\eta_B(x), 0) = (0, 0)$, from which we conclude that $\eta_B(x) = 0$; thus $x \in [B, B]$. $\square$

Example 3.8. We show that Nil_c^{op} is Spl -protomodular and $\mathcal{P}\text{Spl}$ -protomodular, where $\mathcal{P}\text{Spl}$ denotes the class of protosplit monomorphisms, i.e. kernels of split epimorphisms. We also show that Nil_c^{op} is not SMono -protomodular. Since Nil_c is a semi-abelian variety, Nil_c^{op} is finitely complete.

Nil_c^{op} is Spl-protomodular

In this case the morphisms $e \in \mathcal{M}^{\text{op}}$ as in diagram (3.ii) are split epimorphisms. In Nil_c (and also in Grp) it is easy to check that a quotient e admits a splitting if and only if $\text{Ker}(e)$ admits a complement in B , i.e. a subgroup $H \leq B$ such that $\text{Ker}(e) \cap H = 0$ and $B = \text{Ker}(e)H$. By Lemma 3.6, we must prove that for any pushout of type (3.i) and split epimorphism $e \in \mathcal{M}^{\text{op}}$, we have

$$(\text{Ker}(e) \subseteq \text{Ker}(\beta) \cap \text{Ker}(u)) \Rightarrow \text{Ker}(e) = 0.$$

We know that $\text{Ker}(e)$ admits a complement H in B and $\text{Ker}(e) \subseteq [B, B]$ by Lemma 3.7. We get

$$B = \text{Ker}(e)H \subseteq [B, B]H \subseteq B \Rightarrow B = [B, B]H \Rightarrow H = B.$$

The last implication only holds in Nil_c . By induction on c one can prove that, if B is a nilpotent group of class c and $H \leq B$ such that $B = [B, B]H$, then $B = H$. To finish, $\text{Ker}(e) = \text{Ker}(e) \cap B = \text{Ker}(e) \cap H = 0$.

Nil_c^{op} is PSpl-protomodular

In this case e is the cokernel of a split monomorphism i (with retraction r), whose kernel $\langle\langle Q \rangle\rangle$ is the normal closure of Q in B . We consider i to be an inclusion

$$\begin{array}{ccccc} \langle\langle Q \rangle\rangle & \xrightarrow{\text{ker}(e)} & B & \xrightarrow{e=\text{coker}(i)} & B/\langle\langle Q \rangle\rangle \\ & \searrow r & \nearrow i & & \\ & Q & & & \end{array}$$

Suppose that $\langle\langle Q \rangle\rangle \subseteq \text{Ker}(\beta) \cap \text{ker}(u)$. By Lemma 3.7, $\langle\langle Q \rangle\rangle \subseteq [B, B]$. Considering Q a subgroup of B , we have

$$Q = r(Q) \subseteq r(\langle\langle Q \rangle\rangle) \subseteq r([B, B]) = [r(B), r(B)] = [Q, Q] \subseteq Q.$$

We conclude that $Q = [Q, Q]$, i.e. it is a perfect (nilpotent) group. Up to this point, our arguments hold both in Nil_c and in Grp . If Q is a nilpotent group of class c , then $0 = \gamma_{c+1}(Q) = \gamma_c(Q) = \dots = \gamma_1(Q) = Q$; thus $\langle\langle Q \rangle\rangle = 0$.

Nil_c^{op} is not SMono-protomodular

In this case e is just a strong epimorphism, and taking its kernel gives a short exact sequence (with no extra conditions on the basic setting of diagram (3.ii)). In particular, we can consider a pushout (3.i) where u is a strong epimorphism and take $e: B \rightarrow A^B + D$ to be the diagonal (strong epimorphism) of the pushout of (u, β) . This implies that $\text{Ker}(e) = \text{Ker}(\beta) \cap \text{Ker}(u)$. If Nil_c^{op} is SMono-protomodular, then $\text{Ker}(e) = 0$. However, this is not true in general. For example, in the pushout (3.i), take $B = D_4 = \langle r, s : r^4 = s^2 = 1, srs = r^{-1} \rangle$ the dihedral group of order 8 (which is nilpotent of class 2), $A = \langle s \rangle$, $C = 0$ and α the inclusion. In this setting, u is the cokernel of α , so that $u: D_4 \rightarrow D_4/\langle\langle s \rangle\rangle$, with $\langle\langle s \rangle\rangle = \{1, s, r^2, r^2s\}$. Consider the normal subgroup $L = \{1, r^2, rs, r^3s\}$ (because it has order 4 in a group of order 8) and define $\beta: D_4 \rightarrow \langle s \rangle$ by $\beta(L) = \{1\}$ and $\beta(D_4 \setminus L) = \{s\}$. We have $\text{Ker}(\beta) \cap \text{Ker}(u) = L \cap \langle\langle s \rangle\rangle = \{1, r^2\} \neq 0$.

Example 3.9. Grp^{op} is not SMono-protomodular, since neither is Nil_c . Although Lemma 3.7 and some of the arguments of Example 3.8 hold for Nil_c and Grp , we want to emphasize that Grp^{op} is not Spl-protomodular nor PSpl-protomodular.

Grp^{op} is not Spl-protomodular

In the pushout (3.i) take $B = S_3$, $A = \langle (12) \rangle$, $C = 0$, α the inclusion and $\beta(A_3) = \{\text{id}\}$, $\beta(S_3 \setminus A_3) = \{(12)\}$, where $A_3 = \{\text{id}, (123), (132)\}$ is the alternating group of order 3 (i.e. of even permutations). In this setting, u is the cokernel of the split monomorphism α . Since $\langle\langle (12) \rangle\rangle = S_3$, then $u: S_3 \rightarrow S_3/\langle\langle (12) \rangle\rangle = 0$. In diagram (3.ii), take $e = \beta$, which is a split epimorphism in

Grp. We have, $\text{Ker}(e) = \text{Ker}(\beta) = A_3$, $\text{Ker}(u) = S_3$. So, $\text{Ker}(e) \subseteq \text{Ker}(\beta) \cap \text{Ker}(u)$, but $\text{Ker}(e) = A_3 \neq 0$.

Grp^{op} is not $\mathcal{P}\text{Spl}$ -protomodular

Let Q be a non-trivial perfect group, i.e. $0 \neq Q = [Q, Q]$. In the pushout (3.i) take $B = Q \times Q$, $A = \Delta_Q = \{(q, q) : q \in Q\} \cong Q$, $C = 0$, α to be the inclusion and $\beta(q, q') = (q', q')$. In this setting, u is the cokernel of the split monomorphism α . Since $\langle\langle \Delta_Q \rangle\rangle = Q \times Q$ (because Q is perfect), then $u: Q \times Q \rightarrow Q \times Q / \langle\langle \Delta_Q \rangle\rangle = 0$. In diagram (3.ii), take e to be the cokernel of the split monomorphism given by the inclusion $Q \times 0 \hookrightarrow Q \times Q$, so that $e: Q \times Q \rightarrow Q \times Q / Q \times 0 \cong Q \cong \Delta_Q$. We have, $\text{Ker}(e) = \text{Ker}(\beta) = Q \times 0 \cong Q$, $\text{Ker}(u) = Q \times Q$. So, $\text{Ker}(e) \subseteq \text{Ker}(\beta) \cap \text{Ker}(u)$, but $\text{Ker}(e) \cong Q \neq 0$.

4. $\mathcal{M}$ -PROTOMODULAR CATEGORIES VS. $\mathcal{M}$ -MAL'TSEV CATEGORIES

It is well-known from [5] that every protomodular category is necessarily a Mal'tsev category. Recall from [9, 10, 8] that a *Mal'tsev category* $\mathcal{C}$ is defined through the property that every reflexive relation $R: X \rightrightarrows X$ in $\mathcal{C}$ is an equivalence relation. An equivalent definition is to ask that any relation $D: X \rightrightarrows Y$ in $\mathcal{C}$ is *difunctional*. The original definitions asks for the base category $\mathcal{C}$ to be regular or just finitely complete. However, these properties on relations can be stated through generalised elements without any assumption on $\mathcal{C}$. In this work we shall focus on the Mal'tsev property concerning difunctionality of relations.

Definition 4.1. A relation $X \xleftarrow{d_1} D \xrightarrow{d_2} Y$ in $\mathcal{C}$ is called *difunctional* when the relation

$$(4.i) \quad \mathcal{C}(W, X) \xleftarrow{\mathcal{C}(W, d_1)} \mathcal{C}(W, D) \xrightarrow{\mathcal{C}(W, d_2)} \mathcal{C}(W, Y)$$

in **Set** is difunctional [21], for every object W of $\mathcal{C}$. More precisely, given morphisms $x, u: W \rightarrow X$, $y, v: W \rightarrow Y$, we have $(xDy \wedge uDy \wedge uDv) \Rightarrow xDv$, that is

$$(4.ii) \quad \begin{array}{c} \begin{array}{ccc} & X & \\ x \nearrow & & \nwarrow d_1 \\ W & \cdots \langle x, y \rangle \cdots & D \\ y \searrow & & \swarrow d_2 \\ & Y & \end{array} & \begin{array}{ccc} & X & \\ u \nearrow & & \nwarrow d_1 \\ W & \cdots \langle u, y \rangle \cdots & D \\ y \searrow & & \swarrow d_2 \\ & Y & \end{array} & \begin{array}{ccc} & X & \\ u \nearrow & & \nwarrow d_1 \\ W & \cdots \langle u, v \rangle \cdots & D \\ v \searrow & & \swarrow d_2 \\ & Y & \end{array} \Rightarrow \begin{array}{ccc} & X & \\ x \nearrow & & \nwarrow d_1 \\ W & \cdots \langle x, v \rangle \cdots & D \\ v \searrow & & \swarrow d_2 \\ & Y & \end{array} \end{array}$$

Given a category with binary products and a class $\mathcal{M}$ of monomorphisms, we say that a relation $X \xleftarrow{d_1} D \xrightarrow{d_2} Y$ is in $\mathcal{M}$ when the monomorphism $\langle d_1, d_2 \rangle: D \rightarrow X \times Y$ is in $\mathcal{M}$.

Definition 4.2. Let $\mathcal{C}$ be a category with binary products and $\mathcal{M}$ a class of monomorphisms of $\mathcal{C}$. We call $\mathcal{C}$ an $\mathcal{M}$ -Mal'tsev category when any relation $X \xleftarrow{d_1} D \xrightarrow{d_2} Y$ in $\mathcal{M}$ is difunctional.

Example 4.3. 1. When $\mathcal{M} = \text{Mono}$, an $\mathcal{M}$ -Mal'tsev category is just an “ordinary” Mal'tsev category.
2. When $\mathcal{M} = \text{SMono}$, an $\mathcal{M}$ -Mal'tsev category is a weakly Mal'tsev category [17] (see definition in Remark 4.7).

We show next that, if a finitely complete category $\mathcal{C}$ is $\mathcal{M}$ -protomodular, where $\mathcal{M}$ is a class of monomorphisms (with certain properties), then $\mathcal{C}$ is an $\mathcal{M}$ -Mal'tsev category. The proofs of the $\mathcal{M}$ -versions in the following results are similar to those for “ordinary” protomodular and Mal'tsev categories.

Proposition 4.4. *Let $\mathcal{C}$ be an $\mathcal{M}$ -protomodular category, where $\mathcal{M}$ is a class of monomorphisms of $\mathcal{C}$. Given an arbitrary pullback of points*

$$(4.iii) \quad \begin{array}{ccccc} Z \times_Y X & \xrightarrow{\pi_X} & X & & \\ \pi_Z \downarrow & \lrcorner & \downarrow i_X & & \downarrow f \\ & & Z & \xrightarrow{h} & Y, \\ & & \uparrow t & & \uparrow s \end{array}$$

the pair (i_Z, i_X) is orthogonal to the class $\mathcal{M}$.

Proof. Since the total and right diagrams in

$$\begin{array}{ccccc} & & 1_X & & \\ & \searrow & \text{---} & \searrow & \\ X & \xrightarrow{i_X} & Z \times_Y X & \xrightarrow{\pi_X} & X \\ f \downarrow & & \downarrow \pi_Z & & \downarrow f \\ Y & \xrightarrow{t} & Z & \xrightarrow{h} & Y \\ & \swarrow & \text{---} & \swarrow & \\ & & 1_Y & & \end{array}$$

are pullbacks, then so is the left diagram. The result follows by Proposition 1.5. $\square$

Proposition 4.5. *Let $\mathcal{C}$ be a category with pullbacks and $\mathcal{M}$ a class of monomorphisms of $\mathcal{C}$ with the properties*

- (P2) $\mathcal{M}$ is closed under composition;
- (P4) for any pullback projections π_Z, π_X , the monomorphism $\langle \pi_Z, \pi_X \rangle: Z \times_Y X \rightarrow Z \times X$ is in $\mathcal{M}$;
- (P5) $\mathcal{M}$ is stable under pullbacks.

The following conditions are equivalent:

- (i) $\mathcal{C}$ satisfies the property of Proposition 4.4;
- (ii) any relation $\langle d_1, d_2 \rangle: D \rightarrow X \times Y$ in $\mathcal{M}$ is difunctional.

Proof. (i) $\Rightarrow$ (ii) Consider the relations as in (4.ii). We take the following pullbacks, and induced morphism $\lambda: W \rightarrow P$

$$\begin{array}{ccccccc} W & \xrightarrow{\langle \langle u, y \rangle, \langle u, v \rangle \rangle} & & & & & \\ \lambda \searrow & & & & & & \\ & P & \xrightarrow{p_2} & \text{Eq}(d_1) & \xrightarrow{d_{12}} & D & \\ p_1 \downarrow & \lrcorner & \downarrow i_2 & & \downarrow s_1 & \downarrow d_1 & \\ & & \boxed{A} & & & & \\ & \text{Eq}(d_2) & \xrightarrow{d_{22}} & D & \xrightarrow{d_1} & X & \\ d_{21} \downarrow & \lrcorner & \downarrow s_2 & & \downarrow d_2 & & \\ & & D & \xrightarrow{d_2} & Y. & & \end{array}$$

Next we take the pullback of $\langle d_1, d_2 \rangle$ along $\langle d_1 d_{21} p_1, d_2 d_{12} p_2 \rangle$, which is a monomorphism in $\mathcal{M}$ by property (P5), and the induced morphisms $\alpha: \text{Eq}(d_2) \rightarrow Q$, $\beta: \text{Eq}(d_1) \rightarrow Q$

$$\begin{array}{ccc}
 \text{Eq}(d_2) & \xrightarrow{d_{21}} & D \\
 \downarrow \alpha & \searrow & \downarrow \langle d_1, d_2 \rangle \in \mathcal{M} \\
 Q & \xrightarrow{q} & D \\
 \downarrow \mu \in \mathcal{M} & \lrcorner & \downarrow \langle d_1, d_2 \rangle \in \mathcal{M} \\
 P & \xrightarrow{\langle d_1 d_{21} p_1, d_2 d_{12} p_2 \rangle} & X \times Y \\
 \downarrow i_1 & & \\
 P & \xrightarrow{\langle d_1 d_{21} p_1, d_2 d_{12} p_2 \rangle} & X \times Y
 \end{array}
 \quad
 \begin{array}{ccc}
 \text{Eq}(d_1) & \xrightarrow{d_{12}} & D \\
 \downarrow \beta & \searrow & \downarrow \langle d_1, d_2 \rangle \in \mathcal{M} \\
 Q & \xrightarrow{q} & D \\
 \downarrow \mu \in \mathcal{M} & \lrcorner & \downarrow \langle d_1, d_2 \rangle \in \mathcal{M} \\
 P & \xrightarrow{\langle d_1 d_{21} p_1, d_2 d_{12} p_2 \rangle} & X \times Y \\
 \downarrow i_2 & & \\
 P & \xrightarrow{\langle d_1 d_{21} p_1, d_2 d_{12} p_2 \rangle} & X \times Y
 \end{array}$$

Since $\boxed{\text{A}}$ is a pullback of points, then (i_1, i_2) is orthogonal to $\mathcal{M}$ by assumption. From the commutativity of

$$\begin{array}{ccc}
 & Q & \\
 \alpha \nearrow & \downarrow \mu \in \mathcal{M} & \nwarrow \beta \\
 \text{Eq}(d_2) & \xrightarrow{i_1} P \xleftarrow{i_2} & \text{Eq}(d_1)
 \end{array}$$

we conclude that μ is an isomorphism. We then have $\langle d_1, d_2 \rangle q \mu^{-1} \lambda = \langle x, v \rangle$, which proves that $x D v$.

(ii) $\Rightarrow$ (i) Consider the pullback of points below and a monomorphism $m \in \mathcal{M}$ such that $m\alpha = i_Z$ and $m\beta = i_X$ in

$$\begin{array}{ccc}
 M & \xleftarrow{\beta} & X \\
 \downarrow m \in \mathcal{M} & \searrow & \downarrow \pi_X \\
 Z \times_Y X & \xrightarrow{\pi_X} & X \\
 \downarrow \pi_Z & \downarrow i_Z & \downarrow i_X \\
 Z & \xrightarrow{h} Y & \\
 \uparrow \alpha & \uparrow t & \\
 Z & \xleftarrow{t} & Y
 \end{array}$$

By (P2) and (P4) it follows that $\langle \pi_Z m, \pi_X m \rangle: M \rightarrow Z \times X$ is in $\mathcal{M}$; let us denote this relation by M . It is easy to check that

$$\begin{aligned}
 \langle \pi_Z m, \pi_X m \rangle \alpha \pi_Z &= \langle \pi_Z, sh \pi_Z \rangle & \Rightarrow & \pi_Z M sh \pi_Z, \\
 \langle \pi_Z m, \pi_X m \rangle \alpha t \pi_X &= \langle t f \pi_X, sh \pi_Z \rangle & \Rightarrow & t f \pi_X M sh \pi_Z, \\
 \langle \pi_Z m, \pi_X m \rangle \beta \pi_X &= \langle t f \pi_X, \pi_X \rangle & \Rightarrow & t f \pi_X M \pi_X.
 \end{aligned}$$

By difunctionality of M we get $\pi_Z M \pi_X$, from which we conclude that m is an isomorphism. $\square$

Proposition 4.6. *Let $\mathcal{C}$ be a category with pullbacks and $\mathcal{M}$ a class of monomorphisms of $\mathcal{C}$ with the properties*

(P2) $\mathcal{M}$ is closed under composition;

(P4) for any pullback projections π_Z, π_X , the monomorphism $\langle \pi_Z, \pi_X \rangle: Z \times_Y X \rightarrow Z \times X$ is in $\mathcal{M}$;

(P5) $\mathcal{M}$ is stable under pullbacks.

If $\mathcal{C}$ is an $\mathcal{M}$ -protomodular category then it is an $\mathcal{M}$ -Mal'tsev category.

Proof. This follows immediately from Propositions 4.4 and 4.5. $\square$

Remark 4.7. 1. When $\mathcal{M} = \text{Mor}$ or $\mathcal{M} = \text{Mono}$, Proposition 4.4 translates to: any protomodular category is a Mal'tsev category. Indeed, any pullback of points (4.iii) being such that (i_Z, i_X) is jointly extremally epimorphic is equivalent to every $\text{Pt}_Y(\mathcal{C})$ being a unital

category [5], for any object Y of $\mathbb{C}$. This latter is a well-known characterisation of Mal'tsev categories [5, Proposition 10]. Proposition 4.5 recovers the equivalence between two well-known characterisations of Mal'tsev categories: every $\text{Pt}_Y(\mathbb{C})$ is a unital category, for any object Y of $\mathbb{C}$, and every relation $D: X \leftrightarrow Y$ is difunctional.

2. When $\mathcal{M} = \text{SMono}$, Proposition 4.4 translates to: any weakly protomodular category is a weakly Mal'tsev category. Recall from [20] that a *weakly Mal'tsev category* is defined by the property that any pullback of points (4.iii) is such that (i_Z, i_X) is jointly epimorphic. Proposition 4.5 recovers the characterisation of weakly Mal'tsev categories as the finitely complete categories such that every strong relation $D: X \leftrightarrow Y$ is difunctional (see [17]).

5. Ord-PROTOMODULAR CATEGORIES

In this section we explore the notion of $\mathcal{M}$ -protomodular category when the base category $\mathbb{C}$ is Ord-enriched and $\mathcal{M}$ is the class of all full (and faithful) monomorphisms – denoted by *ff-monomorphisms* – of $\mathbb{C}$.

Definition 5.1. Let $\mathbb{C}$ be a finitely complete Ord-category. We call $\mathbb{C}$ an *Ord-protomodular category* when, for any morphism $h: Z \rightarrow Y$, the change-of-base functor $h^*: \text{Pt}_Y(\mathbb{C}) \rightarrow \text{Pt}_Z(\mathbb{C})$ is conservative on the class of ff-monomorphisms of $\mathbb{C}$.

Lemma 5.2. *If $\mathbb{C}$ is a protomodular category, then any Ord-enrichment of $\mathbb{C}$ is an Ord-protomodular category.*

Proof. As mentioned earlier, $\mathbb{C}$ is protomodular if and only if any change-of-base functor is conservative on monomorphisms. In particular, they are conservative on ff-monomorphisms. $\square$

Remark 5.3. Propositions 1.4, 1.5 and 1.6 hold in the context of Ord-categories, where $\mathcal{M}$ is the class of ff-monomorphisms. Propositions 1.7 and 1.8 hold for any (pointed) regular Ord-category with 2-pullbacks, so that 2-pullbacks and pullbacks coincide. Similarly, Proposition 4.4 holds for any Ord-protomodular category and Proposition 4.5 for any Ord-category with 2-pullbacks (again, so that 2-pullbacks and pullbacks coincide). Note that the assumptions (P1)–(P5) on morphisms needed for some of these results from Section 1 and Section 4 to hold, are replaced by the properties (F1)–(F5) on ff-monomorphisms listed below, which are easily seen to be always true in the context of Ord-categories with 2-pullbacks.

Lemma 5.4. *Let $a: A \rightarrow B$ and $b: B \rightarrow C$ be morphisms in an Ord-category $\mathbb{C}$. Then:*

- (F1) *if ba is an ff-monomorphism, then a is an ff-monomorphism;*
- (F2) *if a and b are ff-monomorphisms, then ba is also an ff-monomorphism;*
- (F3) *if a is an ff-monomorphism, then $a \times a$ is an ff-monomorphism;*
- (F4) *for any 2-pullback projections π_Z, π_X , the morphism $\langle \pi_Z, \pi_X \rangle: Z \times_Y X \rightarrow Z \times X$ is an ff-monomorphism;*
- (F5) *ff-monomorphisms are stable under 2-pullbacks.* $\square$

The only properties on ff-monomorphisms concerning the Ord-enrichment of $\mathbb{C}$ are (F4) and (F5), where 2-pullbacks are mentioned instead of pullbacks. However, if the finitely complete category $\mathbb{C}$ admits a 2-pullback of a pair of morphisms (f, g) , then it coincides with the pullback of (f, g) .

The notion of Ord-Mal'tsev category introduced in [15] can also be seen as a particular instance of the notion of $\mathcal{M}$ -Mal'tsev category, when $\mathcal{M}$ is the class of ff-monomorphisms. Hence Proposition 4.6 gives, in particular, that every Ord-protomodular category is Ord-Mal'tsev.

Remark 5.5. The category $\mathbf{Ord}^{\text{op}}$, with its natural $\mathbf{Ord}$ -enrichment given, for morphisms $f, g: X \rightarrow Y$, by $f \leq g$ if $f(y) \leq g(y)$ for all $y \in Y$ is not $\mathbf{Ord}$ -protomodular. As we have seen in Example 2.3, $\mathbf{Ord}^{\text{op}}$ is an $\mathbf{SMono}$ -protomodular category which is not protomodular. In this enriched context, ff-monomorphisms coincide with monomorphisms (that is, surjective monotone maps in $\mathbf{Ord}$), not with strong monomorphisms, hence $\mathbf{Ord}$ -protomodularity fails because protomodularity fails.

Example 5.6. Ord-enrichments of $\mathbf{OrdGrp}$. We start by recalling two different $\mathbf{Ord}$ -enrichment of $\mathbf{OrdGrp}$ studied in [14]. We denote by $\mathbf{OrdGrp}$ the one given, for any morphisms $f, g: X \rightarrow Y$, by $f \leq g$ if, for all $x \in X$, $f(x) \leq g(x)$, while we denote by $\mathbf{OrdGrp}_+$ the other enrichment, where $f \leq g$ if $f(x) \leq g(x)$ for all positive elements x of X . We point out that the former enrichment induces an equivalence relation on the morphisms.

As shown in [11] and in Example 2.4 respectively, $\mathbf{OrdGrp}$ is not a protomodular category but it is $\mathbf{SMono}$ -protomodular. Here we show that neither $\mathbf{OrdGrp}$ nor $\mathbf{OrdGrp}_+$ are $\mathbf{Ord}$ -protomodular. The failure of this property is due to the fact that, for both $\mathbf{Ord}$ -enrichments, ff-monomorphisms do not characterise “initial monomorphisms”, that is, injective homomorphisms whose domain has the initial topology. Indeed:

- a monomorphism $m: X' \rightarrow X$ is an ff-monomorphism in $\mathbf{OrdGrp}$ if and only if, for all $x, x' \in X'$, $x \cong x'$ whenever $m(x) \cong m(x')$, so, in particular, if the order on X' is anti-symmetric m is an ff-monomorphism if and only if it is monomorphic;
- a monomorphism $m: X' \rightarrow X$ is an ff-monomorphism in $\mathbf{OrdGrp}_+$ if and only if, for all $x, x' \in X'$, if $0 \leq x$ and $0 \leq x'$, then $x \leq x'$ whenever $m(x) \leq m(x')$. Hence if the only positive element of X' is 0, m is an ff-monomorphism if and only if it is monomorphic.

Diagrams (5.i) and (5.ii) show, respectively, that $\mathbf{OrdGrp}$ and $\mathbf{OrdGrp}_+$ fail to satisfy the ff-monomorphism version of protomodularity.

$$(5.i) \quad \begin{array}{ccccc} (\mathbb{Z}, \mathbb{N}) & \xrightarrow{\langle 1_{\mathbb{Z}}, 0 \rangle} & (\mathbb{Z} \times \mathbb{Z}, P_{\text{prod}}) & \xrightleftharpoons[\langle 0, 1_{\mathbb{Z}} \rangle]{\pi_2} & (\mathbb{Z}, \mathbb{N}) \\ \parallel & & \downarrow 1_{\mathbb{Z} \times \mathbb{Z}} & & \parallel \\ (\mathbb{Z}, \mathbb{N}) & \xrightarrow{\langle 1_{\mathbb{Z}}, 0 \rangle} & (\mathbb{Z} \times \mathbb{Z}, P_{\text{lex}}) & \xrightleftharpoons[\langle 0, 1_{\mathbb{Z}} \rangle]{\pi_2} & (\mathbb{Z}, \mathbb{N}), \end{array}$$

$$(5.ii) \quad \begin{array}{ccccc} (\mathbb{Z}, \{0\}) & \xrightarrow{\langle 1_{\mathbb{Z}}, 0 \rangle} & (\mathbb{Z} \times \mathbb{Z}, P_{\text{prod}}) & \xrightleftharpoons[\langle 0, 1_{\mathbb{Z}} \rangle]{\pi_2} & (\mathbb{Z}, \mathbb{N}) \\ \parallel & & \downarrow 1_{\mathbb{Z} \times \mathbb{Z}} & & \parallel \\ (\mathbb{Z}, \{0\}) & \xrightarrow{\langle 1_{\mathbb{Z}}, 0 \rangle} & (\mathbb{Z} \times \mathbb{Z}, P_{\text{lex}}) & \xrightleftharpoons[\langle 0, 1_{\mathbb{Z}} \rangle]{\pi_2} & (\mathbb{Z}, \mathbb{N}), \end{array}$$

In both diagrams the order of $\mathbb{Z} \times \mathbb{Z}$ in the top row is the product order, while in the bottom row is the lexicographic order (see [11] for details); that is, in (5.i) the positive cones are given by $P_{\text{prod}} = \mathbb{N} \times \mathbb{N}$ and $P_{\text{lex}} = \{(n, m) \in \mathbb{Z} \times \mathbb{Z}; m > 0\} \cup \{(n, 0); n \in \mathbb{N}\} = \mathbb{Z} \times \mathbb{N}_1 \cup \mathbb{N} \times \{0\}$, while in (5.ii) $P_{\text{prod}} = \{0\} \times \mathbb{N}$ and $P_{\text{lex}} = \mathbb{Z} \times \mathbb{N}_1 \cup \{(0, 0)\}$. In both cases $P_{\text{prod}} \subset P_{\text{lex}}$ and so the identity $1_{\mathbb{Z} \times \mathbb{Z}}$ is a monotone bijection but not an isomorphism.

- Remark 5.7.*
1. As shown in [15], $\mathbf{OrdGrp}_+$ is $\mathbf{Ord}$ -Mal'tsev, hence this is an example of an $\mathbf{Ord}$ -Mal'tsev category which is not $\mathbf{Ord}$ -protomodular.
 2. We do not have an example of an $\mathbf{Ord}$ -protomodular category which is not protomodular.
 3. We point out that the notion of $\mathbf{Ord}$ -protomodular is not directly related to the notion of lax protomodularity introduced in [14]: lax protomodularity deals with the properties of the lax comma change-of-base functor.

Disclosure statement Examples 2.5 and 3.8 were obtained with the help of Anthropic’s Claude Fable 5.1 (via Claude Code), in response to our direct questions whether there were such examples. We did not use any other help from AI to produce this paper.

REFERENCES

- [1] M. Barr, P.A. Grillet, D.H. van Osdol, *Exact categories and categories of sheaves*, Springer Lecture Notes in Math. 236 (1971).
- [2] F. Borceux, D. Bourn, *Mal’cev, protomodular, homological and semi-abelian categories*, Math. Appl., vol. 566, Kluwer Acad. Publ., 2004.
- [3] F. Borceux, M.M. Clementino, *Topological semi-abelian algebras*, Adv. Math. **190** (2005) 425–453.
- [4] D. Bourn, *Normalization equivalence, kernel equivalence and affine categories*, Springer Lecture Notes in Math. 1488 (1991) 43–62.
- [5] D. Bourn, *Mal’cev categories and fibration of pointed objects*, Appl. Categ. Structures **4** (1996) 307–327.
- [6] D. Bourn, G. Janelidze, *Protomodularity, descent, and semidirect products*, Theory Appl. Categ. **4** No. 2 (1998) 37–46.
- [7] D. Bourn, N. Martins-Ferreira, A. Montoli, M. Sobral, *Monoids and pointed S-protomodular categories*, Homology Homotopy Appl. **18** (2016) 151–172.
- [8] A. Carboni, G.M. Kelly, M.C. Pedicchio, *Some remarks on Maltsev and Goursat categories*, Appl. Categ. Structures **14** (1993) 385–421.
- [9] A. Carboni, J. Lambek, M.C. Pedicchio, *Diagram chasing in Mal’cev categories*, J. Pure Appl. Algebra **69** (1991) 271–284.
- [10] A. Carboni, M.C. Pedicchio, N. Pirovano, *Internal graphs and internal groupoids in Mal’cev categories*, Am. Math. Soc. for the Canad. Math. Soc., Providence, 1992, pp. 97–109.
- [11] M.M. Clementino, N. Martins-Ferreira, A. Montoli, *On the categorical behaviour of preordered groups*, J. Pure Appl. Algebra **223** (2019) 4226–4245.
- [12] M.M. Clementino, A. Montoli, *On the categorical behaviour of V-groups*, J. Pure Appl. Algebra **225** (2021), paper 106550, 24 pp.
- [13] M.M. Clementino, A. Montoli, *Right-preordered groups from a categorical perspective*, Algebra Universalis **86** (2025), paper 8, 21 pp.
- [14] M. M. Clementino, A. Montoli, D. Rodelo, *On lax protomodularity for Ord-enriched categories*, J. Pure Appl. Algebra 227(8) (2023) 107348.
- [15] M.M. Clementino, D. Rodelo, *Enriched aspects of calculus of relations and 2-permutability*, J. Algebra Appl., 2025, 2650233.
- [16] M.M. Clementino, W. Tholen, *Metric, topology and multicategory – a common approach*, J. Pure Appl. Algebra **179** (2003) 13–47.
- [17] Z. Janelidze, N. Martins-Ferreira, *Weakly Mal’tsev categories and strong relations*, Theory Appl. Categ. **27** No.5 (2012) 65–79.
- [18] A. Kurz, J. Velebil, *Quasivarieties and varieties of ordered algebras: regularity and exactness*, Math. Struct. Comput. Sci. **27** (2017) 1153–1194.
- [19] G. Janelidze, L. Márki and W. Tholen, *Semi-abelian categories*, J. Pure Appl. Algebra **168** (2002), 367–386.
- [20] N. Martins-Ferreira, *Weakly Mal’tsev categories*, Theory Appl. Categ. **21**(6) (2008) 91–117.
- [21] J. Riguet, *Relations binaires, fermetures, correspondances de Galois*, Bull. Soc. Math. France **76** (1948) 114–155.
- [22] G.J. Seal, W. Tholen, *Monoidal Structures*, in: Monoidal Topology, Encyclopedia Math. Appl. **113**, pp. 19–144.
- [23] V. Aravantinos-Sotiropoulos, *The exact completion for regular categories enriched in posets*, J. Pure Appl. Algebra 226(7) (2022) 106885.

CMUC, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF COIMBRA, 3000-143 COIMBRA, PORTUGAL
 Email address: mmc@mat.uc.pt

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF THE ALGARVE, 8005-139 FARO, PORTUGAL AND CENTER FOR RESEARCH AND DEVELOPMENT IN MATHEMATICS AND APPLICATIONS (CIDMA), DEPARTMENT OF MATHEMATICS, UNIVERSITY OF AVEIRO, 3810-193 AVEIRO, PORTUGAL
 Email address: drodello@ualg.pt