

*Towards a realizability model of homotopy type
theory*

Jonas Frey

joint work (in progress) with Steve Awodey and Pieter Hofstra

CT 2017 Vancouver

Overview

- **Motivation** : construct *realizability model of homotopy type theory*, to show consistency of *impredicative univalent universe*
- **Approach** : internalize *cubical set model* in Hyland's effective topos *Eff*
- **Context** : build on related work by Coquand et al., Orton/Pitts, Gambino/Sattler, Frumin/van den Berg, Rosolini

Homotopy type theory :

Re-reading of Martin-Löf's **dependent type theory** where

- ① **types** are **spaces**
- ② **equalities** are **paths**

... more precisely :

Dependent Type Theory

Dependent type theory comprises:

- simple types $1, X, Y, A \times B, A \Rightarrow B, A + B, \dots$
- dependent types / type families $x:A \vdash B(x)$
- dependent sum types $\Sigma x:A. B(x)$ and product types $\Pi x:A. B(x)$
- inductive types $\mathbb{N}, \text{list}(A), \dots$
- identity types $x:A, y:A \vdash \text{Id}_A(x, y)$
- universes \mathcal{U} which are ‘types of types’, closed under the preceding type constructors

Identity types

In the set-theoretic model, identity types are given by

$$\text{Id}_A(x, y) = \begin{cases} \{*\} & \text{if } x = y \\ \emptyset & \text{else} \end{cases}.$$

In **locally cartesian closed categories**, identity types are modeled by diagonals

$$A \rightarrow A \times A.$$

Interpretation satisfies **uniqueness of identity proofs**

$$\text{(UIP)} \quad \prod a, b : A. \prod p, q : \text{Id}_A(a, b). \text{Id}_{\text{Id}_A(a, b)}(p, q),$$

not provable in type theory (Hofmann-Streicher 1994).

Irritating to classical mathematicians, but leaves room for a **homotopical interpretation**.

Identity types as path objects

Awodey-Warren (2009): interpret **Id**-types by **fibrant replacement** of diagonal, i.e. second part of a trivial-cofibration/fibration factorization

$$\begin{array}{ccc} A & \xrightarrow{\sim} & PA \\ & \searrow & \downarrow \text{Id}_A \\ & & A \times A \end{array}$$

w.r.t. a **weak factorization system / WFS** (possibly part of a model structure).

Intuition: elements of $\text{Id}_A(a, b)$ are **paths** from a to b .

Lifting property of WFS corresponds to **elimination rule** of **Id**-types.

Coherence problem solved by using ‘categories-with-families’ and cloven WFS.

h-levels and equivalences

Types satisfying UIP can be recovered as **0-types** in HoTT.

More generally, **n -types** for $n \geq -2$ are inductively defined as follows:

- A is a **(-2) -type** (or **contractible type**), if

$$\Sigma x:A. \Pi y:A. \text{Id}_A(x, y)$$

is inhabited.

- A is a **$(n + 1)$ -type**, if $\text{Id}_A(x, y)$ is an n -type for all $x, y:A$.

We call **(-1) -types propositions**, and **0-types sets**.

Equivalences

A function $f : A \rightarrow B$ is called an **equivalence**, if its **fibers**

$$\Sigma x : A . \text{Id}_B(fx, y)$$

are contractible for all $y : B$.

$\text{equiv}(A, B)$ is the type of equivalences from A to B .

Universes and univalence

When should two types be considered equal?

Voevodsky's **univalence axiom** asserts that two types are equal iff they are homotopy equivalent.

More precisely, a universe \mathcal{U} is called **univalent**, if the canonical map

$$\text{Id}_{\mathcal{U}}(A, B) \rightarrow \text{equiv}(A, B)$$

is an equivalence for all $A, B : \mathcal{U}$.

Univalence is inconsistent with UIP as soon as a type in \mathcal{U} has a non-trivial automorphism.

Since classical logic implies “proof-irrelevance”, it is inconsistent with univalence.

A model of HoTT with univalent universe in simplicial sets has been described by Voevodsky, written down by Kapulkin-Lumsdaine 2012.

Predicative and impredicative universes

- Ordinary **predicative** universes are closed under **small** products of small types:

$$\frac{\Gamma \vdash A : \mathcal{U} \quad \Gamma, x:A \vdash B(x) : \mathcal{U}}{\Gamma \vdash \prod x : A. B(x) : \mathcal{U}}$$

- **Impredicative** universes are closed under **arbitrary** products of small types:

$$\frac{\Gamma, x:A \vdash B(x) : \mathcal{U}}{\Gamma \vdash \prod x : A. B(x) : \mathcal{U}}$$

- Subobject classifier Ω of a topos models impredicative universe of propositions.
- Impredicative universe \mathcal{U} containing a type $A : \mathcal{U}$ with two distinct elements $x \neq y : A$ inconsistent with classical logic.

Impredicative universes in realizability toposes

The **effective topos** *Eff* (Hyland 1980) models an impredicative universe \mathcal{M} containing non-propositional types.

\mathcal{M} is not univalent (since in topos-models, all types are 0-types)

To get an **univalent, impredicative universe**, need something like

- **homotopical realizability model** or
- **realizability- ∞ -topos**

Constructing the model internally to *Eff*

Observation : Existence of univalent universe in simplicial set model relies on assumption of Grothendieck universe in meta-theory.

Idea : perform model construction internally to *Eff* (containing impredicative universe) to obtain **univalent** impredicative universe.

Working internally to *Eff* imposes restrictions:

- constructive internal logic (no excluded middle)
- no transfinite constructions (no ‘small object argument’)

Coquand et al observed that the simplicial model relies on classical logic, proposed to use **cubical sets** instead.

Cubical sets

Cubical sets are presheaves on a **cube category**

- **Monoidal cube category** \mathbb{C}_m used by Serre, Kan in 50ies
- **Symmetric cube category** \mathbb{C}_s : free symmetric monoidal category on an interval (Bezem, Coquand, Huber 2013)
- **Cartesian cube category** \mathbb{C}_c : free finite-product category on an interval / Lawvere theory with two constants
- **Cartesian cube category with connections** \mathbb{C}_{cc} : Lawvere theory of distributive lattices / full subcat of **Cat** on objects $\mathbb{2}^n$
- **Lawvere theory of de Morgan algebras** \mathbb{C}_{dm} (Cohen, Coquand, Huber, Mörtberg 2016)

Comparison :

- all locally finite & can be internalized in *Eff*
- use \mathbb{C}_c or \mathbb{C}_{cc}
- \mathbb{C}_c much simpler than \mathbb{C}_{cc} :

$$\#\mathbb{C}_c([9], [1]) = 11 \quad \#\mathbb{C}_{cc}([9], [1]) = ? \quad (9\text{th Dedekind number})$$

More on cube categories :

“Varieties of cubical sets” – Buchholtz, Morehouse 2017

(Iterated) path spaces in cartesian cubical sets $\widehat{\mathbb{C}}$

$[0], [1], [2], \dots$ objects of cube category.

Interval : $I = Y([1])$

n -cube : $I^n = Y([n]) = Y([1])^n$

Path object : $PA = A^I = A(- \times [1])$

Iterated path object : $P^n A = A^{I^n} = A(- \times [n])$

(I **tiny object**, $A \mapsto A^I$ has right adjoint – ‘fractional exponent’)

Path space factorization

Awodey 2016 : algebraic weak factorization system (AWFS) on $\widehat{\mathbb{C}}_c$ such that

$$\begin{array}{ccc} A & \xrightarrow{\tilde{\pi}} & A' \\ & \searrow & \downarrow \langle A^\perp, A^\top \rangle \\ & & A \times A \end{array}$$

is an (L, R) -factorization.

Construction uses small objects argument

To avoid this and be able to internalize in *Eff*, restrict to **Kan complexes**.

Uniform normal Kan complexes

Box inclusions analogous to simplicial Horn inclusions :

$$\bigsqcup_j^n \hookrightarrow I^n \quad n \in \mathbb{N}, j \in \{\perp, \top\}$$

Uniform Kan complexes have coherently chosen box fillings :

$$\begin{array}{ccc} X \times \bigsqcup_j^n & \xrightarrow{f} & A \\ \downarrow & \nearrow \tilde{f} & \\ X \times I^n & & \end{array}$$

Normality condition : fillers of 'degenerate boxes' are degenerate

$$\begin{array}{ccc} \bigsqcup_j^n & \longrightarrow & A \\ \downarrow & \nearrow I^{n-1} & \\ I^n & & \end{array}$$

$\mathcal{F}(\widehat{\mathbb{C}}_c) \subseteq \widehat{\mathbb{C}}_c$ category of uniform normal Kan complexes

Cloven weak factorization systems

A **cloven weak factorization system** / **CWFS** (van den Berg, Garner 2010) on \mathcal{C} is a functorial factorization

$$\begin{array}{ccc}
 \begin{array}{ccc}
 A & \xrightarrow{h} & B \\
 \downarrow f & & \downarrow g \\
 X & \xrightarrow{k} & Y
 \end{array} & \mapsto &
 \begin{array}{ccc}
 A & \xrightarrow{h} & B \\
 Lf \downarrow & & \downarrow Lg \\
 P(f) & \xrightarrow{P(h,k)} & P(g) \\
 Rf \downarrow & & \downarrow Rg \\
 X & \xrightarrow{k} & Y
 \end{array}
 \end{array}$$

with specified fillers for all $f : A \rightarrow B$ (no naturality requirement):

$$\begin{array}{ccc}
 \begin{array}{ccc}
 A & \xrightarrow{LLf} & P(Lf) \\
 Lf \downarrow & \nearrow & \downarrow RLf \\
 Pf & \xrightarrow{id} & Pf
 \end{array} & &
 \begin{array}{ccc}
 Pf & \xrightarrow{id} & Pf \\
 LRf \downarrow & \nearrow & \downarrow Rf \\
 P(Rf) & \xrightarrow{RRf} & B
 \end{array}
 \end{array}$$

Theorem

CWFS with stable functorial choice of diagonal factorization gives rise to model of **Id**-types.

A cloven CWFS on $\mathcal{F}(\widehat{\mathbb{C}}_c)$

The mapping-cocylinder factorization on uniform Kan complexes gives a cloven CWFS satisfying the conditions of the theorem :

$$\begin{array}{ccccc}
 & & A & \xrightarrow{f} & B \\
 & & \downarrow Lf \lrcorner & & \downarrow \\
 & & Pf & \xrightarrow{\quad} & B' \\
 & Rf \swarrow & \downarrow \lrcorner & & \downarrow \\
 B & \xleftarrow{\pi_2} & A \times B & \xrightarrow{f \times B} & B \times B \\
 & & \downarrow \lrcorner \pi_1 & & \downarrow \pi_1 \\
 & & A & \xrightarrow{f} & B
 \end{array}$$

The induced WFS

Every CWFS induces a WFS with left maps L -coalgebras and right maps R -algebras.

Theorem

TFAE for $i : U \rightarrow X$ in $\mathcal{F}(\widehat{\mathbb{C}}_c)$:

- ① i is a left map for the mapping-cocylinder CWFS
- ② i is (the section part of) a strong deformation retract

TFAE for $f : A \rightarrow B$ in $\mathcal{F}(\widehat{\mathbb{C}}_c)$:

- ① f is a right map for the mapping-cocylinder CWFS
- ② f is a uniform normal Kan fibration
- ③ f has uniform normal path lifting (1-box filling)

$$\begin{array}{ccc} X & \longrightarrow & A \\ \downarrow & & \downarrow f \\ X \times I & \longrightarrow & B \end{array}$$

Σ -Types and Π -Types

- Σ types are easy
- Π -types are more subtle, so far we only know how to get them using connections (using ideas of Gambino-Sattler and Frumin-vdBerg)

Trivial fibrations and cofibrations

Definition

- $f : A \rightarrow B$ is a **homotopy equivalence**, if there exists $g : B \rightarrow A$ and homotopies $gf \sim \text{id}$ and $fg \sim \text{id}$
- f is a **trivial fibration**, if it is a (normal, uniform) fibration and a homotopy equivalence
- i is a **cofibration**, if it has the lrp wrt all trivial fibrations

Theorem

TFAE:

- f is a trivial fibration
- f is the retract part of a strong deformation retract
- f admits uniform, normal right liftings wrt $\partial I^n \hookrightarrow I^n$

TFAE:

- i is a cofibration
- i is monic and has rlp wrt $\delta : I \rightarrow I \times I$

There is a trivial-fibration/cofibration factorization (see related work by Bourke-Garner, Frumin-van den Berg, Coquand)

Constructing the impredicative universe

In general :

- \mathcal{C} small category, κ inaccessible cardinal
- $F : A \rightarrow B$ in $\widehat{\mathcal{C}}$ called κ -**small**, if $\overline{F} \in \widehat{\int B}$ has κ -small fibers
- **generic** κ -small map $p : \tilde{\mathcal{U}} \rightarrow \mathcal{U}$ given by

$$\mathcal{U}(\mathcal{C}) = \text{hom}(y\mathcal{C}, \mathcal{U}) \cong [(\int y\mathcal{C})^{\text{op}}, \mathbf{Set}_{\kappa}] \cong [(\mathcal{C}/\mathcal{C})^{\text{op}}, \mathbf{Set}_{\kappa}]$$

$$\tilde{\mathcal{U}}(\mathcal{C}) = \coprod_{F \in \mathcal{U}(\mathcal{C})} F(\text{id})$$

- perform construction in cubical sets internal to $\mathcal{E}ff$, with \mathcal{M} for κ
- use arguments of Gambino-Sattler (after Cisinski) to show fibrancy and univalence
- work in progress, connections probably required

Bibliography, related work

- van den Berg, Garner. “Topological and simplicial models of identity types” (2012)
- Bezem, Coquand, Huber. “A model of type theory in cubical sets”, (2014)
- Cohen, Coquand, Huber, Mörtberg. “Cubical type theory: a constructive interpretation of the univalence axiom” (2016)
- Orton, Pitts. “Axioms for Modelling Cubical Type Theory in a Topos” (2016)
- Gambino, Sattler. “The Frobenius condition, right properness, and uniform fibrations” (2017)
- Sattler. “The Equivalence Extension Property and Model Structures” (2017)
- Cisinski. “Univalent universes for elegant models of homotopy types” (2014)
- Frumin, van den Berg. “A homotopy-theoretic model of function extensionality in the effective topos” (2017)
- Rosolini. “The category of equilogical spaces and the effective topos as homotopical quotients” (2016)

Thanks for your attention!